



Fluoride in Groundwater, Its Variation with Seasons and Relationship with Other Water Quality Parameters in Rangareddy District of Telangana, India

M. Ranjith^{a*}, S. Sridevi^{b#}, K. Jeevan Rao^a and T. Ramesh^{c≡}

^a Department of Soil Science, College of Agriculture, Rajendranagar, Professor Jayashankar Telangana State Agricultural University, Telangana State - 500030, India.

^b Agricultural Research Station, Tornala, Siddipet-502114, India.

^c Department of Crop Physiology, College of Agriculture, Rajendranagar-500030, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i1230952

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/83996>

Original Research Article

Received 20 January 2022

Accepted 28 March 2022

Published 02 April 2022

ABSTRACT

Fluorine, which occurs naturally as fluoride in the earth's crust in small amounts, is essential to humans and animals and is known to stimulate plant growth when present in small concentrations. However, higher concentrations of fluoride in drinking water cause dental and skeletal fluorosis in animals and humans and when present in high concentrations in groundwater used for irrigation, it accumulates in the plants and the soil and causes undesirable effects. A study was carried out during *kharif*, 2015 and *rabi* season of 2015-16 in Talakondapalle Mandal of Rangareddy District, Telangana state, India, to measure and map the distribution of fluoride in groundwater in borewells used for irrigation, to observe seasonal variation in fluoride content and to examine the relationship of fluoride with other water quality parameters such as pH, EC, Cl^- , CO_3^- , HCO_3^- , SO_4^{2-} , Ca^{+2} , Mg^{+2} , Na^+ , K^+ , RSC and SAR. Fluoride in groundwater ranged between 0.72 to 4.50 ppm with a mean of 2.84 ppm and standard deviation of 1.30 ppm in *kharif* season and from 1.32 to 4.92 ppm with a mean of 3.29 ppm and standard deviation of 0.24 ppm in *rabi* season. Except for two borewells in *kharif*, fluoride concentration was higher than the permissible limit of 1 ppm. Groundwater fluoride

[#] Principal Scientist & Head;

[≡] Professor & Head;

*Corresponding author: E-mail: ranjithagrico34@gmail.com;

map of the study area was generated using kriging technique. Fluoride concentration was higher in all the borewells in *rabi* over *kharif* with mean fluoride concentration being 15.8% higher in *rabi* season. Fluoride concentration was significantly positively correlated with pH, HCO_3^- , Na^+ , SAR and RSC of ground water and significantly negatively correlated with Ca^{+2} .

Keywords: Fluoride; semi-arid region; irrigation; safe limit; crop production.

1. INTRODUCTION

Fluorine is a very reactive element that occurs naturally as fluoride in the earth's crust in small amounts. The primary source of fluoride in soil and groundwater is weathering and dissolution of fluoride containing minerals like fluorspar, cryolite, fluorapatite, fluorite, fluorophlogopite, etc. [1,2]. Anthropogenic activities such as brick production, production of phosphate fertilizers, cement, and other industrial processes discharge fluoride into the environment. Although fluoride in small amounts is essential to humans and animals and is known to stimulate plant growth, its presence in higher concentrations in drinking water causes dental and skeletal fluorosis in animals and humans and when present in high concentrations in groundwater, it leads to decreased plant growth and productivity and accumulation in plant tissue [3]. In many parts of the world, elevated levels of F contaminated groundwater used for irrigation purposes have been reported to cause considerable adverse effects on crops [4,5]. Physiological processes known to be markedly affected by high concentration of fluoride include chlorosis [6], leaf tip burn and necrosis [7] and decreased plant growth [8]. While Lione *et al.* suggested 10 mg/l as the safe limit of fluoride in irrigation water for all types of plants [8], the Food and Agricultural Organisation [10] prescribed limit is 1.0 mg/l.

Groundwater fluoride concentrations tend to be higher in arid regions, where the groundwater flow is low and the reaction times with rocks are long. The amount of fluoride occurring naturally in groundwater depends mainly on climate, composition of the host rock and hydrogeology [5]. In India, the excessive presence of fluorides in groundwater is reported in nearly 177 Districts covering 20 states and Telangana is one of them [11]. The present study was undertaken to investigate the fluoride content of groundwater used for irrigation in Talakondapalle Mandal (revenue division consisting of several villages, also called Block,

Tehsil, Taluk, etc. elsewhere in India), Rangareddy District, Telangana State, India, to examine seasonal variations in fluoride content and to examine correlations between the concentration of fluoride and other water quality parameters.

2. MATERIALS AND METHODS

Talakondapalle Mandal is located in southern part of Rangareddy District with headquarters at Talakondapalle village, located at 16.88.74 Northlatitude and 78.41.69 East longitude, 72 km from Hyderabad, the capital of Telangana State. The climate of the study area is generally hot semi-arid with mean annual rainfall is 616.1 mm. The predominant crops grown in the area are paddy, maize, sorghum and vegetables. The crops are irrigated with groundwater from borewells. The study area forms a part of the stable Dharwar craton of south Indian shield. Grey granite occupies dominant portion of the area, these rocks are composed of quartz, feldspars, and biotite. These are medium to coarse grained and equigranular in texture. Biotite is the most predominant mineral in these rocks [12].

Groundwater samples were collected from 25 borewells located in 10 villages of Talakondapalle Mandal (Fig. 1) during both *kharif*, 2015 (monsoon) and *rabi* (post monsoon) season of 2015-16. Borewells are vertically drilled wells with depth range from 23 to 60 meters in the study area. Farmers use this groundwater for irrigation purpose. Samples were collected in 1 litre plastic bottles previously thoroughly cleaned with distilled water and subsequently rinsed with sampled ground water before filling. Samples were collected after 10 minutes of pumping from the borewell. The samples were transferred to the laboratory at Professor Jayashankar Telangana State Agricultural University, Hyderabad. The fluoride concentration in groundwater was determined potentiometrically, using fluoride ion selective electrode. Three ml of total ionic strength adjusting buffer grade III (TISAB III) was added to 30 ml of sample groundwater and fluoride

concentration was determined [13]. Other water quality parameters were determined using standard procedures [14]. The pH and EC of water samples were determined using pH meter and EC meter respectively. Chloride (Cl^-) was determined by standard AgNO_3 titration. Carbonate (CO_3^{2-}) and bicarbonate (HCO_3^-) were determined by titration with $0.1 \text{ H}_2\text{SO}_4$. Sulphate (SO_4^{2-}) was determined using UV-visible spectrophotometer. Calcium (Ca^{2+}) and magnesium (Mg^{2+}) were determined titrimetrically using standard EDTA method. Sodium (Na^+) and potassium (K^+) were measured by flame photometry. Sampling, preservation and analysis of water samples were carried out following the method recommended by American Public Health Association (APHA). Residual sodium carbonate (RSC) and sodium adsorption ratio (SAR) were computed as follows:

$$\text{RSC} = (\text{CO}_3^- + \text{HCO}_3^-) - (\text{Ca}^{++} + \text{Mg}^{++})$$

$$\text{SAR} = \text{Na}^+ / \sqrt{(\text{Ca}^{2+} + \text{Mg}^{2+})/2}$$

Through descriptive statistics and correlation analysis the relation between fluoride and other water quality parameters was tested for statistical significance.

3. RESULTS AND DISCUSSION

Fluoride content of irrigation water samples varied from 0.72 to 4.49 ppm with a mean of 2.84 ppm and standard deviation of 1.30 ppm in the *kharif* season (Table 1). The lowest fluoride content was recorded in borewell no. 6 located in Talakondapalle village (0.72 ppm) and the highest, in borewell no. 24 in Venkatrapeta village (4.49 ppm). In the *rabi* season, the average fluoride content of irrigation water samples was 3.29 ppm with a range of 1.32 to 4.91 ppm and a standard deviation of 0.24 ppm (Table 2). Borewell no. 13 in Rampur village (4.91 ppm) exhibited the highest fluoride content and borewell no. 6 in Talakondapalle village, the lowest (1.32 ppm). In the study area, groundwater in only 2 wells (borewell 6 and 7) had fluoride content lower than the permissible limit of 1 ppm (mg/l) prescribed by Food and Agriculture Organization (FAO) in *kharif*, while in *rabi* all the samples had fluoride higher than the permissible limit. 35% of ground water samples in Kalwakurthy area of Mahabubnagar district had high fluoride content [15] and ranged from 0.79 to 4.20 ppm [16].

Maps of groundwater fluoride concentration (Figs. 2 and 3) were prepared for both the

seasons using kriging, a geostatistical interpolation technique that considers both the distance and degree of variation between known data points to estimate values for unknown areas [17]. Data of other water quality parameters are presented in Tables 1 and 2. The groundwater samples were slightly to moderately alkaline. Salinity was within permissible limits for irrigation. Concentrations of anions (chloride, carbonate, bicarbonate and sulphate) and cations (calcium, magnesium, sodium and potassium) were within permissible limits prescribed for irrigation water. RSC and SAR were also within permissible limits. The only major problem with the irrigation water was excess fluoride.

3.1 Seasonal Variation in Fluoride Concentration

Seasonal variation in fluoride content was observed in the borewells of the study area (Tables 1 and 2; Figs. 1 and 2). Fluoride content in irrigation water of all the borewells in all the villages was lower in *kharif* compared to *rabi*. Mean fluoride concentration of the sample borewells in *rabi* was 15.8% higher over *kharif*. The widest seasonal variation was observed in borewell no. 7 in Talakondapalle village, where fluoride concentration in *rabi* was 69.7% higher over *kharif*. The lower fluoride concentration in all the borewells in *kharif* may be attributed to dilution of fluoride due to replenishment of the groundwater by rainfall during the *kharif* season. Seasonal distribution of fluoride is dependent on a variety of factors such as amount of soluble and insoluble F in source rocks, the duration of contact of water with rocks and soil temperature, rainfall and oxidation- reduction process [18].

3.2 Relationships between Fluoride and Other Water Quality Parameters

Pearson correlation coefficients (r) were estimated between the F content and other water quality parameters and are presented in Table 3. Significant positive correlations of fluoride content were observed with pH, bicarbonate, sodium, SAR and RSC in both the seasons and calcium in *rabi* season. A significant positive correlation ($r = 0.543$ in *kharif* and 0.482 in *rabi*) was observed between fluoride concentration and pH of groundwater samples. The pH of irrigation water in the study area was neutral to alkaline, which favours the solubility of fluorine bearing minerals. Reddy *et al.* and Janardhanaraju *et al.* have also indicated that weathering and leaching of fluorine in rock

formations under alkaline environment leads to the enrichment of fluoride in the groundwater [19,20]. Alkaline water activates the processes of dissociation and dissolution of F from weathered rocks [21,22].

Fluoride concentration was significantly positively correlated with sodium, carbonates, RSC and

SAR. This suggests that an increase activity of carbonate and bicarbonate ions coupled with the sodium content increased the solubility and release of F from the F bearing parent material. Water rich in sodium and carbonates in weathered rock formations allows precipitation of calcite from Ca^{2+} and CO_3^{2-} ions and accelerates




Fig. 1. Location map and borewell locations in Talakondapalle mandal




Fig. 2. Fluoride distribution in groundwater during *kharif* season

Table 1. Fluoride and other quality parameters of the water samples in TalakondapalleMandal (*Kharif, 2015*)

Borewell No.	F ⁻ ppm	pH	EC (dS m ⁻¹)	Cl ⁻ me L ⁻¹	CO ₃ ⁻² me L ⁻¹	HCO ₃ ⁻ me L ⁻¹	SO ₄ ⁻² me L ⁻¹	Ca ⁺² me L ⁻¹	Mg ⁺² me L ⁻¹	Na ⁺ me L ⁻¹	K ⁺ me L ⁻¹	RSC me L ⁻¹	SAR
1	1.70	8.20	0.60	1.10	0.00	7.80	0.22	5.80	2.50	2.00	0.14	-0.50	0.98
2	1.40	7.81	0.65	1.30	0.00	7.50	0.19	5.20	2.80	2.30	0.07	-0.50	1.15
3	1.70	7.90	0.80	0.80	0.20	7.70	0.21	5.90	2.70	1.10	0.10	-0.70	0.53
4	1.30	8.25	1.32	1.10	0.00	6.50	0.18	5.70	2.90	1.50	0.12	-2.10	0.72
5	1.80	8.20	0.50	0.90	0.00	8.90	0.28	5.20	3.00	1.60	0.11	0.70	0.79
6	0.72	7.94	1.05	1.30	0.00	7.20	0.11	6.70	3.50	1.20	0.24	-3.00	0.53
7	0.89	7.92	1.32	0.90	0.00	8.90	0.23	6.50	3.70	1.30	0.31	-1.30	0.58
8	1.10	7.83	1.42	1.30	0.00	8.00	0.34	5.80	3.00	0.90	0.17	-0.80	0.43
9	1.70	8.02	0.95	1.00	0.00	7.30	0.33	6.10	2.40	1.10	0.23	-1.20	0.53
10	1.90	7.69	0.63	1.50	0.00	8.10	0.21	5.50	3.10	0.70	0.15	-0.50	0.34
11	4.10	8.53	0.95	0.90	0.30	7.80	0.43	4.50	2.40	3.20	0.11	1.20	1.72
12	3.85	8.28	0.76	1.70	0.00	8.10	0.33	6.60	2.20	2.40	0.10	-0.70	1.14
13	4.49	8.61	1.10	1.80	0.00	8.60	0.51	5.40	2.50	1.80	0.37	0.70	0.91
14	3.91	8.05	0.72	1.50	0.00	8.40	0.16	5.10	2.80	1.40	0.15	0.50	0.70
15	4.20	8.60	0.56	1.20	0.00	9.60	0.24	6.20	3.80	3.10	0.11	-0.40	1.39
16	3.50	7.96	0.85	1.60	0.00	9.30	0.22	5.40	4.00	1.90	0.10	-0.10	0.88
17	4.30	8.16	0.89	1.20	0.00	9.40	0.19	4.70	3.80	3.20	0.21	0.90	1.55
18	4.10	8.34	0.62	1.10	0.40	8.50	0.22	5.40	3.10	2.70	0.17	0.40	1.31
19	3.20	7.90	0.94	1.00	0.00	9.50	0.31	6.80	3.00	2.10	0.09	-0.30	0.95
20	4.40	8.70	1.23	0.80	0.00	8.10	0.18	4.50	2.50	3.60	0.13	1.10	1.92
21	3.00	7.51	0.59	1.50	0.00	9.40	0.26	6.10	4.00	2.10	0.10	-0.70	0.93
22	3.00	8.13	0.75	1.80	0.00	7.60	0.32	4.30	3.10	1.90	0.14	0.20	0.99
23	2.40	7.96	1.02	0.90	0.20	8.30	0.37	4.90	3.50	1.40	0.12	0.10	0.68
24	4.50	7.90	0.97	1.10	0.00	8.50	0.21	5.40	2.80	2.80	0.15	0.30	1.38
25	3.90	8.40	0.90	1.10	0.00	9.10	0.24	5.70	3.20	2.50	0.11	0.20	1.19
Range	0.72-4.49	7.51-8.70	0.50-1.42	0.80-1.80	0.00-0.40	6.50-9.60	0.11-0.51	4.30-6.80	2.20-4.00	0.70-3.60	0.07-0.37	-3.0-1.20	0.34-1.92
Mean	2.84	8.07	0.88	1.20	0.04	8.32	0.26	5.58	3.05	1.99	0.15	-0.26	0.97
SD	1.30	0.30	0.26	0.32	0.11	0.81	0.09	0.70	0.52	0.79	0.07	0.98	0.41


Table 2. Fluoride and other quality parameters of the water samples in TalakondapallyMandal (*Rabi, 2015-16*)

Borewell No.	F ⁻ ppm	pH	EC (dS m ⁻¹)	Cl ⁻ me L ⁻¹	CO ₃ ⁻² me L ⁻¹	HCO ₃ ⁻ me L ⁻¹	SO ₄ ⁻² me L ⁻¹	Ca ⁺² me L ⁻¹	Mg ⁺² me L ⁻¹	Na ⁺ me L ⁻¹	K ⁺ me L ⁻¹	RSC me L ⁻¹	SAR
1	2.23	8.05	0.45	1.3	0.00	9.3	0.51	6.5	3.0	2.1	0.15	-0.20	1.0
2	1.81	7.65	1.60	1.7	0.00	8.8	0.20	7.2	4.7	1.0	0.07	-3.10	0.4
3	2.25	8.03	0.59	0.9	0.00	9.1	0.38	6.9	2.7	1.8	0.09	-0.50	0.8
4	1.46	8.55	0.88	1.6	0.20	5.8	0.32	6.6	4.1	1.1	0.13	-4.70	0.5
5	2.69	8.46	0.61	1.2	0.00	7.1	0.42	5.8	3.3	2.4	0.11	-2.00	1.1
6	1.32	7.42	1.21	3.1	0.00	7.0	0.18	7.4	4.5	2.0	0.24	-4.90	0.8
7	1.51	7.62	1.53	2.6	0.00	8.7	0.24	7.3	3.8	1.8	0.29	-2.40	0.8
8	1.86	7.93	0.72	2.0	0.00	8.1	0.37	6.4	4.2	1.5	0.07	-2.50	0.7
9	2.30	7.82	0.58	1.5	0.50	8.4	0.32	6.1	3.8	1.9	0.12	-1.00	0.9
10	2.82	7.74	0.77	1.8	0.00	8.8	0.21	6.0	4.0	3.1	0.08	-1.20	1.4
11	4.45	8.35	0.83	1.6	0.00	9.1	0.42	6.3	4.3	1.6	0.12	-1.50	0.7
12	4.10	8.41	0.72	2.1	0.00	9.6	0.36	5.9	3.7	1.9	0.43	0.00	0.9
13	4.91	7.95	1.10	2.4	0.60	9.5	0.66	5.5	4.6	2.1	0.18	0.00	0.9
14	4.15	8.02	0.93	1.8	0.00	9.1	0.25	5.8	3.5	3.3	0.07	-0.20	1.5
15	4.55	8.56	1.16	2.3	0.00	9.4	0.31	5.7	3.8	2.5	0.12	-0.10	1.1
16	4.02	8.02	0.83	3.5	0.00	9.6	0.27	7.1	2.9	2.8	0.10	-0.40	1.3
17	4.72	8.26	0.67	1.4	0.00	10.5	0.20	6.4	4.3	1.7	0.08	-0.20	0.7
18	4.23	8.08	0.71	1.5	0.00	9.7	0.31	5.9	3.9	2.1	0.11	-0.10	0.9
19	3.74	8.01	0.79	1.1	0.00	10.1	0.45	6.5	3.7	2.9	0.09	-0.10	1.3
20	4.61	8.15	0.93	2.3	0.00	9.8	0.29	6.7	3.3	3.8	0.14	-0.20	1.7
21	3.29	8.10	0.85	1.9	0.00	9.5	0.35	5.8	4.3	3.4	0.11	-0.60	1.5
22	3.25	8.19	0.78	1.3	0.00	8.8	0.28	6.0	3.5	2.1	0.07	-0.70	1.0
23	3.14	8.05	0.70	2.4	0.00	8.1	0.32	5.1	3.3	2.3	0.09	-0.30	1.1
24	4.61	8.36	1.02	3.6	0.30	9.4	0.20	5.7	3.8	2.3	0.08	0.20	1.1
25	4.15	8.04	0.90	1.1	0.00	9.1	0.26	5.5	3.7	2.8	0.11	-0.10	1.3
Range	1.32-4.91	7.42-8.56	0.45-1.60	0.90-3.60	0.00-0.60	5.80-10.50	0.18-0.66	5.10-7.40	2.70-4.70	1.00-3.80	0.07-0.43	-4.90-0.20	0.41-1.70
Mean	3.29	8.07	0.87	1.92	0.06	8.90	0.32	6.24	3.78	2.25	0.13	-1.07	1.01
SD	0.24	0.29	0.28	0.72	0.16	1.04	0.11	0.61	0.52	0.70	0.08	1.43	0.33

Table 3. Pearson's coefficients of correlation (r) between fluoride concentration and other water quality parameters

S.No	Parameters	'r' value	
		<i>Kharif</i>	<i>Rabi</i>
1	Fluoride vs pH	0.543 [†]	0.482*
2	Fluoride vs EC	-0.196	-0.134
3	Fluoride vs Chloride	0.157	0.119
4	Fluoride vs Sulphate	0.224	0.160
5	Fluoride vs Carbonate	0.168	0.112
6	Fluoride vs Bicarbonate	0.513 [†]	0.709 [†]
7	Fluoride vs Calcium	-0.372	-0.537 [†]
8	Fluoride vs Magnesium	-0.057	-0.043
9	Fluoride vs Sodium	0.754 [†]	0.462*
10	Fluoride vs Potassium	-0.079	-0.091
11	Fluoride vs SAR	0.745 [†]	0.480*
12	Fluoride vs RSC	0.739 [†]	0.774 [†]

Note: * $P < 0.01$, $†P < 0.05$

**Fig. 3. Fluoride distribution in groundwater during *rabi* season**

the dissolution of CaF_2 , thereby releasing fluoride into the groundwater [23] and also found good correlation of fluoride content with carbonates, bicarbonates, sodium, RSC and SAR [16].

A significant negative correlation was observed between F and calcium in groundwater of the study area during *rabi* season. Janardhanaraju *et al.* reported that high fluoride concentration was correlated with low calcium values and high sodium content in the groundwater [20]. High-fluoride ground waters are mainly associated with a sodium-bicarbonate water type and

relatively low calcium and magnesium concentrations and such water types usually have high pH (>7) values [24].

4. CONCLUSION

Analysis of groundwater samples from 25 borewells in Talakondapalle Mandal of Rangareddy district, India, showed that all the borewells except two had fluoride concentration greater than the prescribed limit of 1 ppm in *kharif*, while in *rabi* all the borewells had fluoride concentration greater than the prescribed limit.

Fluoride concentrations were higher in *rabi* season compared to *kharif*. Fluoride content in groundwater was significantly positively correlated with pH, bicarbonate, sodium, SAR and RSC in both the seasons and significantly negatively correlated with calcium in *rabi* season. The groundwater in all the borewells was unsuitable for irrigation, and appropriate remediation measures are essential for successful crop production.

ACKNOWLEDGEMENTS

The authors are thankful to the Professor Jayashanker Telangana State Agricultural University, Hyderabad, India for financial support and cooperation for the completion of this research study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kumar S, Singh M. Effect of fluoride contaminated irrigation water on Ecophysiology, biomass and yield in *Gossypium hirsutum* L. Trop. Plant Res. 2015;2(2):134–142.
2. Choudhary S, Rani M, Devika OS, Patra A, Singh RK, Prasad SK. Impact of fluoride on agriculture: A review on its sources, toxicity in plants and mitigation strategies. Int. J. Chem. Stud. 2019;7(2):1675-1680.
3. Brindha K, Elango, L. Fluoride in groundwater: Causes, implications and mitigation measures. In: Monroy SD. editor. Fluoride properties, applications and environmental management; 2011.
4. Dissanayake CB, Chandrajith R. Introduction to medical geology, erlangen earth conference series. Springer-Verlag Berlin Heidelberg; 2009.
5. Gupta S, Banerjee S, Saha R, Datta JK, Mondal N. Fluoride geochemistry of groundwater in Nalhati-1 block of the Birbhum District, West Bengal, India. Fluoride. 2006;39(4):318-320.
6. McNulty B, Newman DW. Mechanism of fluoride induced chlorosis. Plant Physiol. 1961;30(4):385-388. DOI: 10.1104/pp.36.4.385
7. Hadujue J. Reaction of some relativity resistant plants to sudden increase in the concentration of fluoride exhalation. Biologia. 1966;21:421–427.
8. Elloumi N, Abdallah FB, Mezghani I, Rhouma A, Boukhrisb M. Effect of fluoride on almond seedlings in culture solution. Fluoride. 2005;38:193-198.
9. Lione IA, Brennan EG, Danis RH, Robbins WR. Some effects of fluoride on peach, tomato and buck wheat when absorbed through roots. Soil Sci. 1948;66:259-266.
10. FAO. Irrigation and drainage papers. 1976;1-81.
11. Narsimha A, Sudarshan V. Contamination of fluoride in groundwater and its effect on human health: A case study in hard rock aquifers of Siddipet, Telangana State, India. Appl Water Sci. 2016;7(5):2501-2512. DOI: 10.1007/s13201-016-0441-0
12. Ravikanth P, Sundaraiah R, Sateesh P. Fluoride and nitrate contamination in the groundwater of Kalwakurthy area, Mahabubnagar district, Telangana state, India. Ind J Appl Res. 2015;5(1):469-471.
13. Wedepohl KH. Handbook of geochemistry (Vol. II-1) (Ed.) Berlin: Springer; 1969.
14. APHA. Standard methods for the examination of water and waste water, Washington DC, United States of America; 1985.
15. Sundaraiah R, Sudarshan V, Madhusudhan N, Ashok K, Kumar MR. Geochemistry of groundwater in Kalwakurthy area, Mahabubnagar District of Andhra Pradesh with special reference to fluoride distribution. J Appl Geochem. 2013; 15(2):238-249.
16. Ranjith M, Rao KJ, Sridevi S, Ramesh T, Bhav MHV. Groundwater quality and fluoride contamination in Kalwakurthy Mandal of Mahabubnagar District, Telangana State, India. Int J Pure Appl Biosci. 2017;5(4):1397-1405. DOI:http://dx.doi.org/10.18782/2320-7051.5602
17. Krige DG. A statistical approach to some basic mine valuation on the Witwatersrand. J. South. Afr. Inst. Min. Metall. 1951;52:119-139.
18. Mahapatra MK, Mishra A, Das BP. Fluorosis first reported in Naupada District of Orissa, India. Ecol. Environ. Conserv. 2005;11(2):277-280.
19. Reddy BM, Khan JA, Sunitha V, Reddy MR. Fluoride contamination in groundwater from parts of Anantapur District, Andhra

- Pradesh. Gond. Geol. Mag. 2011;26(2):1-9.
20. Janardhanaraju N, Dey S, Das K. Fluoride contamination in ground waters of Sonbhadra District, Uttar Pradesh, India. Curr. Sci. 2009;96(7):979-985.
21. Gupta SK, Deshpande RD, Agarwal M, Raval BR. Origin of high fluoride in groundwater in the North Gujarat-Cambay region, India. Hydrogeol J. 2005;13:596-605.
DOI: 10.1007/s10040-004-0389-2
22. Khan SMN, Kumar AR. Role of alkalinity for the release of fluoride in the groundwater of Tiruchengode Taluk, Namakkal district, Tamilnadu, India. Chem Sci Trans. 2013;2(S1):302-308.
DOI: 10.7598/cst2013.37.
23. Saxena VK, Ahmed S. Inferring the chemical parameters for the dissolution of fluoride in groundwater. Environ. Geol. 2003;43:731-736.
Available: <https://doi.org/10.1007/s00254-002-0672-2>
24. Brunt R, Vasak L, Griffioen J. Fluoride in groundwater: Probability of occurrence of excessive concentration on global scale, International Groundwater Resources Assessment Center Report. 2004;1-12.

© 2022 Ranjith et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/83996>