



Nafamostat Mesylate in the Prevention of Post-endoscopic Retrograde Cholangiography Pancreatitis: A Meta-analysis

**Hyder Osman Mirghani^{a#}, Abdelwahab Hassan Abdelwahab Hussien^{b*≡}
and Abdullah Suliman Al Atawi^{c#}**

^a Medical Department, Faculty of Medicine, Internal Medicine and Endocrine, University of Tabuk, KSA.

^b Department of Endemic and Infectious Diseases, Faculty of Medicine, Suez Canal University, Egypt.

^c Medical Department, Faculty of Medicine, Gastroenterology, University of Tabuk, KSA.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2022/v34i28A36021

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/84689>

Systematic Review Article

Received 15 January 2022

Accepted 21 March 2022

Published 30 March 2022

ABSTRACT

Previous evidence from observational studies showed that Nafamostat mesylate (NM) was effective in post-ERCP pancreatitis prevention. We aimed to assess full-text prospective studies on the role of NM in post-ERCP pancreatitis prevention. We searched the PubMed, Medline, and Google Scholar databases for relevant articles during the period from 2009 to November 2020 and updated on March 2022. No restriction regarding the language of publication. The keywords nafamostat mesylate, post-ERCP pancreatitis, prevention, and role were used. A total of 113 studies were identified through the database search, and eight studies (all were published in Asia) met the inclusion criteria for the systematic review. There were five control trials (four randomized and one comparative) and three case-control studies, (3186 patients included). In the present meta-analysis, seven studies concluded the benefit of nafamostat mesylate in the prevention of post-ERCP pancreatitis, and one showed no benefit. The overall effect was highly significant, odd ratio, 0.51, 95%CI=0.38-0.70, P-value=0.0001, heterogeneity=0.17%, P-value for heterogeneity=0.30, $I^2=17%$.

[#] Associate Professor;

[≡] Lecturer

*Corresponding author: E-mail: ahussien@ut.edu.sa;

Nafamostat mesylate might be effective in post-ERCP pancreatitis prevention. Larger randomized multi-center studies investigating the effectiveness in combination with other preventive measures are needed.

Keywords: Nafamostat mesylate; post-ERCP pancreatitis; prevention.

1. INTRODUCTION

Endoscopic retrograde cholangiopancreatography (ERCP) is an effective common diagnostic and therapeutic procedure. Post-ERCP pancreatitis (PEP) may be an unavoidable complication of ERCP. Various endoscopic and pharmacological preventive approaches have been tried, but most were ineffective [1]. The European Society of Gastrointestinal Endoscopy recommended rectal non-steroidal anti-inflammatory drugs (NSAIDs) for the prevention of PEP. However, the American Society of Gastrointestinal Endoscopy and the Japanese guidelines emphasized the lack of efficacy of certain pharmacological measures [2-4]. Nafamostat mesylate (NM) (a protease inhibitor) has been used for the treatment of influenza, pancreatitis, and disseminated intravascular coagulation; recently it has been shown to be useful in Covid-19 [5,6]. Literature investigating the role of NM in the prevention of post-ERCP pancreatitis is lacking, thus we conducted this meta-analysis to assess the role of NM in the prevention of PEP.

2. METHODOLOGY

2.1 Literature Search

We searched PubMed, Medline, and the first 100 articles in Google Scholar databases. No restriction to languages was adopted, all the articles published during the period from 2009 to March 2022 were eligible, and bibliographies of relevant systematic reviews were searched manually for relevant articles.

2.2 Eligibility Criteria

Only randomized controlled trials, clinical trials, and case-control studies were included. The studies must assess the role of Nafamostat mesylate on post-ERCP pancreatitis. Conference abstracts, case reports, case-series, and animal and experimental studies were excluded. Conference abstracts were not included

because of the information to measure the outcomes might be inadequate, trials reporting the hyperamylasaemia and not reporting on the PEP risk reduction were excluded.

2.3 Article Review and Data Abstraction

Two reviewers conducted a systematic literature search according to Cochrane guidelines [7], the reviewers independently screened the titles and including any title potentially related to ERCP, then any abstract evaluating the effects of nafamostat mesylate and pancreatitis in the setting of ERCP was included. During the review, any conflict between the reviewers was resolved by consensus. The opinion of a biostatistician and endoscopist were thought when necessary. One reviewer abstracted the data that was confirmed by the second reviewer, the data were transferred to an extraction sheet including the author's name, year of publication, country of origin, the route of administration and dose of NM, the incidence of pancreatitis, the odd ratio/95%CI, P-values. The different phases of the systematic review were reported in Fig. 1 and Tables 1 & 2.

2.4 The Quality and Risk of Bias Assessment

Cochrane risk of bias was used to assess the quality and risk bias of the randomized controlled studies [8].

2.5 Statistical Analysis

RevMan 54 software was used for the meta-analysis. For nafamostat mesylate (binary) risk ratios (RRs) with 95% confidence intervals (CIs) were combined across relevant studies, the fixed effects module was applied unless if substantial heterogeneity was found (A P value ≤ 0.10 for Cochran's Q test or an $I^2 \geq 50\%$ was suggestive). A two-tailed P < 0.05 was considered statistically significant for all analyses except heterogeneity tests.




Fig. 1. Flow diagram through the different phases of the systematic review (PRISMA flowchart)

3. RESULTS

A total of 113 studies were identified through the database search. Among these 113 papers, thirteen full-text articles were assessed for eligibility: Eight studies (all were published in Asia) met the inclusion criteria for the systematic review. There were five control trials (four randomized and one comparative), and three case-control studies, (3186 patients included with 192 events). In the present meta-analysis, seven studies [9,10-15] concluded the benefit of nafamostat mesylate in the prevention of post-ERCP pancreatitis, and one [16] showed no benefit. The overall effect was highly significant,

odd ratio, 0.51, 95%CI=0.38-0.70, P-value=0.0001, heterogeneity=0.17%, P-value for heterogeneity=0.30, I^2 =17%. Fig. 2 and Table 1 & 2.

The present meta-analysis showed that out of the eight studies included, 7 reduced post-ERCP pancreatitis, while one showed no effect.

4. DISCUSSION

The role of nafamostat mesylate in the prevention of PEP is controversial, in the present meta-analysis, seven studies [9,10-15] concluded the benefit of nafamostat mesylate in

Table 1. Nafamostat mesylate and prevention of post-ERCP pancreatitis

Author	Year	Country	Type	Patients/control	result
Choi et al. [9]	2009	South Korea	A randomized controlled trial	354 vs. 350	Reduction in pancreatitis
Kim et al.[16]	2016	South Korea	A randomized comparison trial	191 vs.191	No difference in 6 vs.24 hours infusion
Kwon et al.[10]	2012	Korea	A case-control study	88 vs. 81	No difference between placebo and nafamostat
Matsumoto et al. [11]	2021	Japan	RCT	292 vs. 149	Nafamostat injection into the intrapancreatic duct produced promising results
Ohuchida et al. [12]	2015	Japan	A randomized controlled trial	409 vs. 409	Reduction in pancreatitis
Park et al. [13]	2011	South Korea	A case-control	203 vs. 203	No difference between 20mg and 50mg
Park et al. [14]	2014	South Korea	A case-control study	53 vs. 53	Both ulinastatin and nafamostat reduced pancreatitis
Yoo et al. [15]	2011	South Korea	A randomized controlled trial	143 vs. 143	Prophylactic intravenous nafamostat mesylate reduces the frequency of post-ERCP pancreatitis.

Table 2. Nafamostat mesylate dose and rate of administration

Author	Year	Country	Type	Route of administration	Dose and duration
Choi et al. [9]	2009	South Korea	A randomized controlled trial	Intravenous Infusion	20mg once
Kim et al.[16]	2016	South Korea	A randomized comparison trial	Intravenous Infusion	20mg for six or 24 hours
Kwon et al.[10]	2012	Korea	A case-control study	Intravenous Infusion	Not stated
Matsumoto et al. [11]	2021	Japan	RCT	Intravenous Infusion	20mg before ERCP
Ohuchida et al. [12]	2015	Japan	A randomized controlled trial	Intravenous Infusion	20mg, 2 hours before ERCP
Park et al. [13]	2011	South Korea	A case-control	Intravenous Infusion	20mg and 50mg before ERCP
Park et al. [14]	2014	South Korea	A case-control study	Intravenous Infusion	20mg before ERCP
Yoo et al. [15]	2011	South Korea	A randomized controlled trial	Intravenous Infusion	20mg 60 minutes before ERCP and for 6 hours after ERCP.




Fig. 2. Effects of nafamostat mesylate on post-ERCP pancreatitis

the prevention of post-ERCP pancreatitis, and one [16] showed no benefit. The overall effect was highly significant, odd ratio, 0.51, 95%CI=0.38-0.70, P-value=0.0001, heterogeneity=0.17%, P-value for heterogeneity=0.30, I²=17%. Akshintala et al. [17] in their meta-analysis showed that nafamostat mesylate is the second most efficacious preventive measure only after topical ephedrine regarding PEP prevention and in line with the current findings. However, Akshintala and colleagues reviewed more than 16 agents including NM. Kubiliun et al. [18] found the NM is promising and warranted future confirmation supporting the present observation. Similar findings were reported by Yuhara et al. [19] who showed that NM is efficacious in PEP prevention (RR = 0.41; 95 %CI 0.28-0.59), Yu et al. [20] conducted a meta-analysis and observed the effectiveness of NM in the prevention of PEP (risk ratio [RR], 0.47; 95% confidence interval [CI], 0.34-0.63). Yu and colleagues study was conducted seven years before. Therefore, an update about this important issue is essential.

Mechanism of action of various pharmacological agents used in PEP prevention:

- Protease inhibitors (nafamostat, gabexate, and ulinastatin) had similar anti-secretory effects, but NM also showed higher potency and long duration of action [21,22]. The need to administer intravenously for a prolonged time perioperative limited their use.
- Somatostatin (relaxation of the sphincter of Oddi) and octreotide (constriction of the sphincter of Oddi), otherwise similar for their anti-secretory properties [23,24]
- Non-steroidal anti-inflammatory drugs (anti-inflammatory), rectal administration may be difficult in patients undergoing

ERCP and may be expelled during insufflation [25,26].

- Antibiotics are limited by microbial resistance, a global health challenge [27]
- Ephedrine (relax duodenal musculature and edema reduction). Ephedrine is superior due to its short window of action [28,29]
- Stents are invasive, costly, and need reoperation to remove

The administration of rectal NSAIDs and ephedrine was found to be synergistic [30].

The limitations of this study are the small number of studies included and the fact that we included both case-control and randomized studies.

5. CONCLUSION

The present meta-analysis showed that nafamostat mesylate might reduce the risk of PEP; the need for intravenous administration for a relatively long duration may further limit their use. The availability of ephedrine and NSAIDs, their cost-effectiveness, easy administration, and their few side effects rank these drugs higher, the combination of NSAIDs and ephedrine may be more effective.

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Doi K, Ikeda M, Hayase N, Moriya K, Morimura N; COVID-UTH Study Group. Nafamostat mesylate treatment in combination with favipiravir for patients critically ill with Covid-19: A case series. *Crit Care*. 2020;24(1):392. DOI: 10.1186/s13054-020-03078-z. PMID: 32620147; PMCID: PMC7332736.
2. Anderson MA, Fisher L, Jain R, et al. Complications of ERCP. *Gastrointest Endosc* 2012;75:467–73.
3. Dumonceau JM, Andriulli A, Deviere J, et al. European society of gastrointestinal endoscopy (ESGE) Guideline: Prophylaxis of post-ERCP pancreatitis. *Endoscopy*. 2010;42:503–15.
4. Zhou W, Li Y, Zhang Q, et al. Risk factors for post endoscopic retrograde cholangiopancreatography pancreatitis: a retrospective analysis of 7,168 cases. *Pancreatology*. 2011;11:399–405.
5. Hoffmann M, Schroeder S, Kleine-Weber H, Müller MA, Drost C, Pöhlmann S. Nafamostat mesylate blocks activation of SARS-CoV-2: New treatment option for COVID-19. *Antimicrob Agents Chemother*. 2020;64(6):e00754-20. DOI: 10.1128/AAC.00754-20 PMID: 32312781; PMCID: PMC7269515.
6. Salanti G, Ades AE, Ioannidis JP. Graphical methods and numerical summaries for presenting results from multiple-treatment meta-analysis: an overview and tutorial. *J Clin Epidemiol*. 2011;64(2):163-71. DOI: 10.1016/j.jclinepi.2010.03.016 Epub 2010 Aug 5. PMID: 20688472.
7. Wu SS, Sun F, Zhan SY. [Bias risk assessment: (3) Revised Cochrane bias risk assessment tool for individual randomized, cross-over trials (RoB2.0)]. *Zhonghua Liu Xing Bing Xue Za Zhi*. 2017;38(10):1436-1440. Chinese. DOI:10.3760/cma.j.issn.0254-6450.2017.10.028. PMID: 29060995.
8. Chang JH, Lee IS, Kim HK, Cho YK, Park JM, Kim SW. Nafamostat for prophylaxis against post-endoscopic retrograde cholangiopancreatography pancreatitis compared with gabexate. *Gut Liver*. 2009; 3(3):205-10. DOI: 10.5009/gnl.2009.3.3.205
9. Choi CW, Kang DH, Kim GH, Eum JS, Lee SM, Song GA et al. Nafamostat mesylate in the prevention of post-ERCP pancreatitis and risk factors for post-ERCP pancreatitis. *Gastrointest Endosc*. 2009; 69(4):e11-8. DOI: 10.1016/j.gie.2008.10.046
10. Kwon YH, Kim JY, Lee SJ, Jang SY, Park HW, Yang HM et al. [Could nafamostat or gabexate prevent the post endoscopic retrograde cholangiopancreatography pancreatitis?]. *Korean J Gastroenterol*. 2012;59(3):232-8.
11. Matsumoto T, Okuwaki K, Imaizumi H, Kida M, Iwai T, Yamauchi H, et al. Nafamostat mesylate is not effective in preventing post-endoscopic retrograde cholangiopancreatography pancreatitis. *Dig Dis Sci*. 2021;66(12):4475-4484. DOI: 10.1007/s10620-020-06782-6
12. Ohuchida J, Chijiwa K, Imamura N, Nagano M, Hiyoshi M. Randomized controlled trial for efficacy of nafamostat mesilate in preventing post-endoscopic retrograde cholangiopancreatography pancreatitis. *Pancreas*. 2015;44(3):415-21. DOI: 10.1097/MPA.0000000000000278
13. Park KT, Kang DH, Choi CW, Cho M, Park SB, Kim HW et al. Is high-dose nafamostat mesilate effective for the prevention of post-ERCP pancreatitis, especially in high-risk patients?. *Pancreas*. 2011;40(8):1215-9. DOI: 10.1097/MPA.0b013e31822116d5
14. Park JY, Jeon TJ, Hwang MW, Sinn DH, Oh TH, Shin WC et al. Comparison between ulinastatin and nafamostat for prevention of post-endoscopic retrograde cholangiopancreatography complications: A prospective, randomized trial. *Pancreatology*. 2014;14(4):263-7. doi: 10.1016/j.pan.2014.03.022
15. Yoo KS, Huh KR, Kim YJ, Kim KO, Park CH, Hahn T, Park SH et al. Nafamostat mesilate for prevention of post-endoscopic retrograde

- cholangiopancreatographypancreatitis: A prospective, randomized, double-blind, controlled trial. *Pancreas*. 2011;40(2):181-6.
DOI: 10.1097/MPA.0b013e3181f94d46
16. Kim SJ, Kang DH, Kim HW, Choi CW, Park SB, Song BJ, Nam HS. A Randomized comparative Study of 24- and 6-Hour infusion of nafamostat mesilate for the prevention of post-endoscopic retrograde cholangiopancreatography pancreatitis: A prospective randomized comparison trial. *Pancreas*. 2016; 45(8):1179-83.
DOI: 10.1097/MPA.0000000000000625
 17. Akshintala VS, Hutfless SM, Colantuoni E, Kim KJ, Khashab MA, Li T, Elmunzer BJ et al. Systematic review with network meta-analysis: Pharmacological prophylaxis against post-ERCP pancreatitis. *Aliment Pharmacol Ther*. 2013;38(11-12):1325-37.
DOI: 10.1111/apt.12534.
 18. Kubiliun NM, Adams MA, Akshintala VS, Conte ML, Cote GA, Cotton PB et al. Evaluation of pharmacologic prevention of pancreatitis after endoscopic retrograde cholangiopancreatography: A systematic review. *Clin Gastroenterol Hepatol*. 2015;13(7):1231-9; quiz e70-1.
DOI: 10.1016/j.cgh.2014.11.038.
 19. Yuhara H, Ogawa M, Kawaguchi Y, Igarashi M, Shimosegawa T, Mine T. Pharmacologic prophylaxis of post-endoscopic retrograde cholangiopancreatography pancreatitis: protease inhibitors and NSAIDs in a meta-analysis. *J Gastroenterol*. 2014;49(3):388-99.
DOI: 10.1007/s00535-013-0834-x
 20. Yu G, Li S, Wan R, Wang X, Hu G. Nafamostat mesilate for prevention of post-ERCP pancreatitis: a meta-analysis of prospective, randomized, controlled trials. *Pancreas*. 2015;44(4):561-9.
DOI: 10.1097/MPA.0000000000000310
 21. Mori S, Itoh Y, Shinohata R, et al. Nafamostat mesilate is an extremely potent inhibitor of human tryptase. *J Pharmacol Sci*. 2003;92:420-3.
 22. Keck T. Site-specific therapeutic effects of protease inhibitors: effect of route of administration in experimental pancreatitis. *Pancreatology*. 2001;1:656-61.
 23. Wu SD, Zhang ZH, Jin JZ, et al. Effects of different drugs on the sphincter of Oddi motility: Study with choledochoscope manometry. *Zhonghua yi xue za zhi*. 2005;85:1911-5.
 24. Di Francesco V, Angelini G, Zoico E, et al. Effect of native somatostatin on sphincter of Oddi motility in patients with acute recurrent pancreatitis. A pilot study with ultrasound-secretin test. *Dig Liver Dis*. 2006;38:268-71.
 25. Alvan G, Orme M, Bertilsson L, et al. Pharmacokinetics of indomethacin. *Clin Pharmacol Ther*. 1975;18:364-73.
 26. Helleberg L. Clinical pharmacokinetics of indomethacin. *Clinical Pharmacokinetics*. 1981;6:245-58.
 27. Spellberg B, Bartlett JG, Gilbert DN. The future of antibiotics and resistance. *N Engl J Med*. 2013;368:299-302.
 28. Pezzilli R, Romboli E, Campana D, et al. Mechanisms involved in the onset of post-ERCP pancreatitis. *JOP*. 2002;3:162-8.
 29. Akashi R, Kiyozumi T, Tanaka T, et al. Mechanism of pancreatitis caused by ERCP. *Gastrointest Endosc*. 2002; 55:50-4.
 30. Ding X, Chen M, Huang S, et al. Nonsteroidal anti-inflammatory drugs for prevention of post-ERCP pancreatitis: a meta-analysis. *Gastrointest Endosc*. 2012; 76:1152-9.

© 2022 Mirghani et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/84689>