



## Knowledge and Practice of Pelvic Inflammatory Disease Prevention among Female Students

Akoko Sokiprim <sup>a\*</sup>, Akpeki Ruth <sup>b</sup> and Uahomo Precious Ojo <sup>a</sup>

<sup>a</sup> Department of Pharmacology , Faculty of Basic Clinical Sciences, College of Health Sciences, University of Port Harcourt, Port Harcourt, Nigeria.

<sup>b</sup> Faculty of Nursing Sciences, College of Health Sciences, Niger Delta University, Wilberforce Island, Bayelsa State, Nigeria.

### Authors' contributions

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### Article Information

#### Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/85137>

Received 20 January 2022  
Accepted 30 March 2022  
Published 02 April 2022

Original Research Article

### ABSTRACT

**Introduction:** Pelvic inflammatory disease, also known as pelvic inflammatory disorder (PID), is an infection of the upper part of the female reproductive system, including the uterus, fallopian tubes, and ovaries. PID is a clinical illness caused by the movement of microorganisms from the lower to the upper genital tract, resulting in upper genital tract inflammation. This study assessed the knowledge and practice of pelvic inflammatory disease prevention strategies among the female students of College of Health Sciences, Niger Delta University, Amassoma, Bayelsa State, Nigeria.

**Method:** Descriptive research design was adopted for this study. The design was considered appropriate for the study on the assessment of knowledge and practices of pelvic inflammatory disease prevention among female students in College of Health Sciences, Niger Delta University. A total of 184 females students in the female Hostels of College of Health Sciences, Niger Delta University, Bayelsa State were approached for this study.

**Results:** Data obtained were analyzed using descriptive statistics of frequency tables and simple percentages. The results obtained from the study revealed that majority of the respondents 64(35%) were between the ages of 15-20years. The marital status of the respondents has 104(57%) as single as against 56(30%) as married and 24(13%) as divorced. The result indicated strongly that majority of the respondents 104(57%) were Christians, while 56(30%) and 32(17%) were Muslims and other religions respectively. Also, students in 500level are the highest respondents with a total number of 48(26%).

**Conclusion:** The result demonstrated that 96(52%) of female students in College of Health Sciences of the University had a considerable level of knowledge of pelvic inflammatory disease. More than half (52%) of the respondents had a good level of knowledge about PID prevention; also an overwhelming majority of them held a very positive attitude towards PID prevention.

**Keywords:** *Pelvic inflammatory disease; upper reproductive tract; chlamydia trachomatis neisseria gonorrhoeae; sexually active.*

## 1. INTRODUCTION

Infection-induced inflammation of the female upper reproductive system (endometrium, fallopian tubes, ovaries, or pelvic peritoneum) is known as pelvic inflammatory disease (PID). Endometritis is an intermediate step in the development of illness, with inflammation spreading from the vaginal or cervix to the upper genital tract. Pelvic discomfort mixed with inflammation of the lower genital tract is the hallmark of the diagnosis; women with pelvic inflammatory disease may have very modest symptoms and indications [1]. Many women develop subclinical pelvic inflammatory disease as a result of infection spreading to the upper vaginal tract in a clinically quiet manner. Pelvic inflammatory disease is a major cause for concern because it can lead to long-term reproductive issues such as infertility, ectopic pregnancy, and chronic pelvic pain. After the introduction of laparoscopy in the 1960s, research on pelvic inflammatory disease proliferated through the 1970s, 1980s, and 1990s, leading to major breakthroughs in the understanding of the microbial causes of the disease and its relationship to reproductive disability, as well as enabling the standardization of antimicrobial treatment [1]. According to a national estimate in 2001, more than 750,000 cases of pelvic inflammatory disease occurred in the United States [2]. Over the past two decades, the rates and severity of pelvic inflammatory disease have declined in North America and Western Europe [3]. These decreases coincided with public health measures to combat *Chlamydia trachomatis* and *Neisseria gonorrhoeae* infection. Despite advances, pelvic inflammatory disease continues to be an issue since reproductive results among treated individuals are still inadequate, subclinical pelvic inflammatory disease is poorly controlled, and preventative programmes are not viable in much of the developing world [4]. It is estimated that PID affects millions of women yearly all over the world and it is witnessing an uptick in Nigeria especially among young women [5]. In Port-Harcourt, Rivers State and Bayelsa State of

Nigeria, the rate of PID was put at 21.0% [6]. In Osogbo, a South-Western Nigerian State, the prevalence of PID was 70.0% [7]. Also, in Nguru, North-Eastern Nigeria, the rate of PID was put at 62.8% [8]. Findings from a study conducted by [9] have confirmed the existence of PID among women of child bearing age in Onitsha North; particularly in Anambra State, Eastern Nigeria, with a high prevalence of 60%. Even though there is paucity of data on PID in Nigeria, available literature on the prevalence of PID across Nigerian States and regions implies that the obvious evidence may not capture the actual reality pertaining the problem in the country. Therefore, in preventing PID, it is important that women are equipped with the knowledge and the necessary skill-sets as well as interventions and advocacy [10]. However, it has been observed that some females at the College of Health Sciences, Niger Delta University, suffer from pelvic inflammatory disease, necessitating the conduct of a study to assess pelvic inflammatory disease prevention knowledge and practises among females at the College of Health Sciences, Niger Delta University, Bayelsa State, Nigeria.

## 2. MATERIALS AND METHODS

### 2.1 Research Design

The descriptive research design was used in this investigation. Because the study involves the collecting of data in order to precisely and objectively describe actual occurrences and determine the nature of a situation as it exists at the time of investigation, this design was chosen. Personal facts, opinions, and attitudes are best obtained using the descriptive design. The descriptive design has the advantage of covering a wider scope and at the same time economical. The design therefore was considered appropriate for the study on the assessment of knowledge and practices of pelvic inflammatory disease prevention among female students in college of health sciences Niger Delta University.

## 2.2 Research Setting

The research was conducted in Niger Delta University, College of health sciences which is a tertiary institution of learning in Wilberforce Island, Amassoma, Bayelsa State. The institution (College of Health Sciences) is made up two campuses; the female hostel which is made up just females of approximately 342 students and the second hostel which comprises of both males and females. It also consists of the following departments: Medicine, Pharmacy, Medical Laboratory Science, Nursing Sciences, Anatomy, Physiology and Biochemistry. The research will be carried out in the female hostels of the first campus in College of Health Sciences.

## 2.3 Target Population

The type to be used is accessible population; they are those members of the population who within the reach of the researcher. The target population here is the entire female students living in the hostels.

## 2.4 Sample Size

The sample for this study was calculated to be two hundred (184) female students living in college of health science female hostel using Taro Yamane [11] sample size calculation formula. Taro Yamane is a simple and very frequently uses formula for sample size determination for estimating proportion in a finite population:

$$n = \frac{N}{1 + N(e)^2}$$

Where;

n = sample size

N= population under study

e = margin of error (0.05)

$$n = \frac{342}{1 + 342(0.05)^2}$$

$$n = \frac{342}{1 + 0.855}$$

$$n = \frac{342}{1.855} = 184$$

## 2.5 Sampling Technique

A convenience sampling technique will be adopted. Sampling technique are procedures adopted to systematically select the chosen sample in a specified way under controls. This research work adopted the random sampling

technique in selecting the respondents from the total population. Convenient sampling technique is a type of non-probability sampling method where the sample is taken from a group of people easy to contact or reach.

## 2.6 Instruments for Data Collection

Data collecting entails looking for useful information that can help solve certain problems. Every research endeavour thus focuses on the search for primary or secondary sources of knowledge. But for the purpose of this study, data for this study was gathered through primary sources (questionnaire) which is a self-developed and carefully structured by the researcher with which data was collected. The questionnaire consists of two parts (sections) A and B. Section A was designed to elicit personal data from the respondents and section B focuses on the assessment of knowledge and practices in the prevention of pelvic inflammatory disease which is the subject of the study.

## 2.7 Validity of Instrument

The instrument is said to be valid when it measure what it is intended to measure. The validity of the instrument for data collection will be reviewed and corrected by my supervisor to ensure that the questions are adequate to answer the objectives and the questions of the research are appropriate.

## 2.8 Reliability of Instrument

To ensure the face and the content validity of the items the reliability of the instruments after seen by the project supervisor, a pre-test of the research instrument was carried out prior to the actual study. Copies of the questionnaire will be administered to 10 students who live in main campus female hostel G which is different from the research setting. Using Combat Alpha, the reliability coefficient is 0.84 was gotten indicating good reliability.

## 2.9 Method of Data Collection

The students were approached in the evening hours during their leisure hour. The researcher briefed them about the study and obtained their consent to participate in the study.

## 2.10 Method of Data Analysis

Data were analyzed and presented using descriptive statistics of frequency tables and simple percentage.

### 3. RESULTS

The demographic variables of the respondents were first presented followed by their assessment of knowledge and practices of pelvic inflammatory disease.

Table 1 shows the demographic variables of the respondents. It reveals that 64 (35%) of the respondents were within the age range of 15-20 years and this percentage was the highest number. Considering the marital status of the respondents, it shows that more than half of the respondents 104 (57%) were singles; 56 (30%) were married; concerning the religion of the respondents, it shows that 57% (104) were Christians; 48 (26%) were Muslim and 32 (17%) were of other religion. For department, 32 (17%) were nursing students which was the highest number of students; while 24 (13%) were medical biochemistry students; 24 (13%) were anatomy students; 16 (9%) was the least percentage and were students of physiology. For the academic level, 32 (17.3%) were 100 level students and 48 (26%) were 500 level students.

Table 2 shows the assessment of the level of knowledge of pelvic inflammatory disease (PID) by the respondents. It was reported by 136 (74%) of the respondents have heard of the PID; subsequently, 104 (57%) of the respondents have made effort to learn about PID; it was reported 128 (70%) of the respondents have got tested for PID; it was reported that 120 (65%) of the respondents agreed that their partners have gotten tested for PID; considerably, 96 (52%) of the respondents has learnt about PID; furthermore, 80 (43%) of the respondents are aware that their partners knows about PID; it was also recorded that 96 (52%) of the correspondents knows the cause of PID; it was stated that 40 (21%) of the respondents use toilet roll during menses; furthermore it was stated that 97 (51%) of the respondents douche while washing their vagina; it was stated that 92 (50%) of the respondents bleed between menses; it was stated that 94 (51%) of the respondents have sticky vagina discharge; it was stated that 64 (32%) of the respondents have vagina itching; it was reported that 40 (21%) of the respondents are aware of the health complications of PID.

**Table 1. Personal demographic data (N=184)**

| Items                      | Frequency (n) | Percentage (%) |
|----------------------------|---------------|----------------|
| <b>Age</b>                 |               |                |
| 15-20 years                | 64            | 35             |
| 21-25 years                | 56            | 30             |
| 26-30 years                | 40            | 22             |
| 30 years and above         | 24            | 13             |
| <b>Marital status</b>      |               |                |
| Single                     | 104           | 57             |
| Married                    | 56            | 30             |
| Divorce                    | 24            | 13             |
| <b>Religion</b>            |               |                |
| Christian                  | 104           | 57             |
| Muslim                     | 48            | 26             |
| Others                     | 32            | 17             |
| <b>Department</b>          |               |                |
| Medicine                   | 24            | 13             |
| Pharmacy                   | 24            | 13             |
| Medical laboratory science | 24            | 13             |
| Nursing                    | 32            | 17             |
| Medical biochemistry       | 24            | 13             |
| Anatomy                    | 24            | 13             |
| Physiology                 | 2             | 16             |
| Others                     | 2             | 16             |
| <b>Level</b>               |               |                |
| 100                        | 32            | 17.3           |
| 200                        | 40            | 22             |
| 300                        | 32            | 17.3           |
| 400                        | 32            | 17.3           |
| 500                        | 48            | 26             |

**Table 2. Level of knowledge of pelvic inflammatory disease**

| <b>What is your level of knowledge of pelvic inflammatory disease (PID)?</b> | <b>Yes (%)</b> | <b>No (%)</b> |
|------------------------------------------------------------------------------|----------------|---------------|
| Have you heard of PID?                                                       | 136 (74%)      | 48 (26%)      |
| Have you made effort to learn about PID?                                     | 104 (57%)      | 80 (43%)      |
| Have you get tested for PID?                                                 | 128 (70%)      | 56 (30%)      |
| Have your partner gotten tested for PID                                      | 120 (65%)      | 64 (35%)      |
| Does your partner know about PID?                                            | 80 (43%)       | 104 (57%)     |
| Do you know the cause of PID?                                                | 96 (52%)       | 88 (48%)      |
| Do you use toilet roll during your menses?                                   | 40 (21%)       | 144 (79%)     |
| Do you douche when washing your vagina?                                      | 97 (51%)       | 87 (49%)      |
| Do you bleed between your menses?                                            | 92 (50%)       | 92 (50%)      |
| Do you have sticky vagina discharge?                                         | 94 (51%)       | 90 (49%)      |
| Do you have frequent vagina itching during those discharges?                 | 64 (32%)       | 120 (68%)     |
| Are you aware of the health complications of PID?                            | 40 (21%)       | 144 (79%)     |

**Table 3. The various practices of pelvic inflammatory disease prevention**

| <b>What are the preventive practices you utilize in preventing pelvic inflammatory disease (PID)?</b> | <b>Yes (%)</b> | <b>No (%)</b> |
|-------------------------------------------------------------------------------------------------------|----------------|---------------|
| Use condoms during sexual intercourse?                                                                | 112 (61%)      | 72 (39%)      |
| Drink salt and water immediately after sexual intercourse?                                            | 64 (35%)       | 120 (65%)     |
| Drink warm water immediately after sexual intercourse?                                                | 72 (39%)       | 112 (61%)     |
| Drink alcohol immediately after sexual intercourse?                                                   | 47 (26%)       | 137 (74%)     |
| Douche during vagina washing?                                                                         | 96 (52%)       | 88 (48%)      |
| Wash your vagina with antiseptics?                                                                    | 80 (43%)       | 104 (57%)     |
| Use drugs such as antibiotics for prevention?                                                         | 167 (91%)      | 17 (9%)       |
| Practice good hygiene generally?                                                                      | 156 (85%)      | 28 (15%)      |
| Use candle wax mixed with water to wash your vagina?                                                  | 37 (20%)       | 147 (80%)     |

Table 3 shows the various practices of pelvic inflammatory disease (PID) prevention by the respondents. It was reported that 112 (61%) of the respondents use condoms during sexual intercourse; furthermore it was reported that 64 (35%) of the respondents drink salt and water immediately after sexual intercourse; it was reported that 72 (39%) of the respondents drink warm water immediately after sexual intercourse; it was also reported that 47 (26%) of the respondents drank alcohol immediately after sexual intercourse; it was further reported that 96 (52%) of the respondents douche during vagina wash; it was reported that 80 (43%) of the respondents washed their vagina with antiseptics; it was reported that 167 (91%) of the respondents use drugs such as antibiotics for prevention of PID; it was further reported that 156 (85%) of the respondents practice good hygiene generally; it was reported that 37 (20%) of the respondents use candle wax mixed with water to wash their vagina. Finally, considering the knowledge of the respondents other specification were given as the cause of PID such; wash with soap and warm water after sexual intercourse, sit on hot water after sexual intercourse, wash with palm oil mixed with warm

water after sexual intercourse, get medical check-up every three month, and sit on hot water mixed with salt after sexual intercourse; these was gotten from 40 (22%) of the general respondents.

#### 4. DISCUSSION

This study assessed the knowledge and practice of pelvic inflammatory disease among female students of College Health Sciences, Niger Delta University. The result presented in Table 1 reveals that majority of the respondents 64 (35%) were between the ages of 15-20years. The marital status of the respondents has 104 (57%) as single as against 56 (30%) as married and 24 (13%) as divorced, this is a reflection of the high population of youths in tertiary institutions. The result demonstrated that 96 (52%) of female students in College of Health Sciences had a considerable level of knowledge of pelvic inflammatory disease. This result is in tandem with the study conducted by Obiechina Nworah et al. [12] who reported a good knowledge and awareness of common STDs (HIV/AIDS, gonorrhoea, syphilis, Chlamydia etc) among

Nigeria adolescent girls in a high school at Onitsha, Anambra State of Nigeria. According to the research, 93.6% of the subject was aware of HIV/AIDS, 76.3% had knowledge of gonorrhoea, while 69.1% and 6.6% had knowledge of syphilis and Chlamydia respectively. The highest source of information was through the school (80.6%), followed by television (80.1%) and radio (73.1%). This is not in agreement with Adegun [13] which recorded a high level of knowledge of STIs among patients attending outpatient clinics at University Teaching Hospital; Ado-Ekiti, Ekiti State of Nigeria. The result also shows similarity with Kennedy *et al.* [14] study on the prevalence of *Chlamydia trachomatis* infection among female undergraduates of the University of Port Harcourt. According to the research, 60% of the subject was aware of *C. trachomatis* infection while 40% were ignorant about the infection. Also the result could be compared study by Chioma and Luke [7] who carried out a study on the prevalence pattern of bacterial isolates in cases of pelvic inflammatory disease patients at a tertiary hospital in Osogbo, Nigeria which, results showed that; 73.6% of the subject was aware of bacterial infection, 26.4% had knowledge of pelvic inflammatory disease. From the result of the study, 48 (26%) of the respondents had not heard of PID before. This could be attributed to a positive attitude and practice of PID prevention which in turn could result in the level of knowledge of PID. Also, the most frequent/common signs and symptoms is sticky vagina discharge. This could be as a result of invasion of bacteria (Chlamydia) in the genital tract. The highest level of knowledge of PID is well reflected in the sanitary practice (use of toilet roll during menses) of the respondents which result as 144 (79%) who do not toilet roll during menses. Again, a good knowledge of PID prevention was indicated by the respondents. With a very sound knowledge of its prevention, there was no doubt that the respondents would also be familiar with the preventive measure. Hence, the result showed 128 (70%) have gotten tested for PID; 112 (61%) use condom during sexual intercourse and 167 (91%) use antibiotics for prevention of PID. Furthermore 120 (655) of the respondents showed a positive attitude by confirming that their partners have been tested of PID while 64 (35%) of the respondent confirm that their partners have not been tested. Finally, 40 (22%) of the respondents gave responses other than what was asked on the questionnaire such as; wash with soap and warm water after sexual intercourse, sit on hot water after sexual intercourse, wash with palm oil mixed with warm

water after sexual intercourse, get medical check-up every three month, and sit on hot water mixed with salt after sexual intercourse.

## 5. CONCLUSION

The aim of the study was to investigate the assessment knowledge and attitudes of female students towards pelvic inflammatory disease prevention at college of health science Niger Delta University Wilberforce Island, Bayelsa State, Nigeria. The results of this study indicates that more than half (52%) of the respondents had good level knowledge about PID prevention; also an overwhelming majority of them held a very positive attitude towards PID prevention. Further analyzing showed that level of knowledge through various means did affect their attitude towards prevention. Very importantly, this knowledge go a long way to drastically reduce the incidence of PID in College of Health Sciences, Niger Delta University.

## 6. RECOMMENDATION

Due to the fact that the respondents have a positive attitude towards PID prevention, I recommend that wearing of condoms during sex which helps to protect from sexually transmitted infection. With unprotected sex, the chance of infection rises. Maintaining one sexual partner is important since the risk of infection rises as the number of sexual partners grows. Sexually active women under the age of 25 should be tested for Chlamydia once a year or if their sexual partner changes. Sexual education should be implemented at the university level to raise student understanding of the problem of sexually transmitted diseases, such as pelvic inflammatory disease, and how to prevent them. This research was limited to just the College of Health Sciences, Niger Delta University, therefore, further research work should be done on this topic and it should involve a larger population.

## CONSENT

As per international standard or university standard, respondents' written consent has been collected and preserved by the author(s).

## ETHICAL CONSIDERATION

The researcher collected a letter of introduction from the Dean of Faculty of Nursing Science

which was given to the Dean of Students' Affairs through the Hall-warden. This granted the researcher access to distribute the questionnaire to the female students. Other ethical issues like; confidentiality, voluntary participation and anonymity were considered in the conduct of the research study. In terms of confidentiality, the respondents will be assured that the information they will provide, will not be disclosed to anyone who is not directly involved in the study. As regards voluntary participation the respondents will be fully informed that participation is voluntary. Again, anonymity, privacy and confidentiality of the respondents will also be ensured during the interview. The respondents were assured that they would remain anonymous throughout the study.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Wang Y, Zhang Y, Zhang Q, Chen H, Feng Y. Characterization of pelvic and cervical microbiotas from patients with pelvic inflammatory disease. *J Med Microbiol.* 2018; 67(10):1519-1526
2. Lindsey K. Jennings; Diann M. Krywko. Pelvic Inflammatory Disease;2021. Available:<https://www.ncbi.nlm.nih.gov/books/NBK499959/> Accessed 20<sup>th</sup> August, 2021
3. WHO. WHO Guidelines for the treatment of *Chlamydia Trachomatis*; 2016. Available:<https://apps.who.int/iris/bitstream/handle/10665/246165/9789241549714-eng.pdf> Accessed 20<sup>th</sup> August, 2021
4. Chioma Njoku and Luke Ifeanyi Anorue Dr. Mass Media exposure and its influence on Pelvic inflammatory disease prevention behavior among women of child bearing age in three Southeastern Nigerian States. *Library Philosophy and practice (e-journal).* 2021;5123. Available:<https://digitalcommons.uni.edu/libphilprac/5123>
5. Obokoh Anthonia. Pelvic inflammatory disease on the rise in Nigeria. *Business Day*;2019. Available:<https://businessday.ng/health/article/pelvic-inflammatory-disease-on-the-rise-in-nigeria/> Accessed August, 2021.
6. Kennedy TW, John O, Sunny E. Prevalence of *Chlamydia trachomatis* infection among female undergraduates of the University of Port Harcourt using strand displacement and amplification [SDA] technique. *The Nigerian Health Journal.* 2012;12:35-38.
7. Olowe OA, Alabi A, Akindele AA. Prevalence pattern of bacterial isolates in cases of pelvic inflammatory disease patients at a tertiary hospital in Osogbo, Nigeria. *Environmental Research Journal.* 2012;6:308-311.
8. Okon KO, Ayilara R, Bello, K, Uba A, Aniesona TA. Microbial spectrum of pelvic inflammatory diseases in Nguru, Nigeria. *African Journal of Clinical and Experimental Microbiology.* 2008;9:157-165.
9. Eze EM, Unegbu VN. Ezebialu CU, Nneji IR. Prevalence of microorganisms associated with Pelvic inflammatory disease in reproductive aged women in Onitsha North, Anambra state, Nigeria. *Novel Research in Microbiology Journal.* 2018;2:147-155.
10. Howells IE, Okwudili EO. Acute pelvic inflammatory disease and its long – term sequelae: The University of Port Harcourt experience. *Journal of Advances in Medicine and Medical Research.* 2018;26:1-11.
11. Taro Yamane. *Statistics: An Introductory Analysis, 2<sup>nd</sup> Edition,* New York: Harper and Row;1967.
12. Obiechina Nworah, Diwe Kevin, Ikpeze Okechukwu. Knowledge, awareness and perception of sexually transmitted diseases (STDs) among Nigerian adolescent girls. *Journal of Obstetrics and Gynecology.* 2002;22(3):302-5. DOI: 10.1080/01443610220130634
13. Adegun PT, Solomon OA, Adegoke SA, Ade-Ojo IP, Fape MO. Knowledge of sexually transmitted infections among patients attending outpatient clinics at University Teaching Hospital, Ado-Ekiti, Nigeria. *Journal of Public Health and Epidemiology.* 2013;5(3): 110-114. DOI: 10.5897/JPHE11.117
14. Kennedy Tamunomiebam Wariso, John Odigie, Sunny Eyearu. Prevalence of *Chlamydia trachomatis* infection among

female undergraduates of the amplification (SDA) Technique. The  
University of Port Harcourt using Nigerian Health Journal. 2012;12(2):  
strand displacement and 35–38.

© 2022 Sokiprim et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/85137>*