



Surgical and Functional Outcome of Early Mobilization in Proximal Tibial Fracture

**Amey R. Gursale^{a*‡}, Rohit S. Kaware^{a‡}, Vaibhav U. Mesare^{a#},
Amey M. Borse^{a‡}, Sushant S. Tuse^{a‡}, Aamir Matin^{a†}
and Saurabh S. Dhamdhare^{a‡}**

^a Seth Nandlal Dhoot Hospital, Aurangabad, Maharashtra, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/84480>

Original Research Article

Received 17 January 2022

Accepted 23 March 2022

Published 28 March 2022

ABSTRACT

Background: The major cause of proximal tibial fractures attributes to road traffic accident, dominated by male gender with younger age group from high energy trauma. With advancement in newer surgical techniques, improvement in the quality of implants, better soft tissue handling during surgery, and early mobilization in the form of assisted and supervised physiotherapy; the results of surgical outcome are improving. We aim to study clinical and functional outcome after surgery with introduction of early mobilization in various types of Proximal Tibial fractures in terms of range of motion and its effect on operative construct.

Methods: This is prospective observational study of 68 cases of Proximal Tibia fractures with 53(77.9%) males and 15(22.1%) females using suitable implants and initiating early physiotherapy from day 2. Follow up at 3rd and 6th week was done to assess the outcome.

Results: Early initiation of physiotherapy showed significant association with improvement in range of motion with choice of implant at 3rd week (p value -0.039) and at 6th week (p value-0.000) and type of fracture as well.

[‡]Senior Resident DNB Orthopaedic;

[#]DNB, D-Ortho, MNAMS, FAGE;

[‡]D-Ortho;

[†]Junior Resident DNB Orthopaedic;

*Corresponding author: Email: a.gursale@yahoo.com;

Conclusion: The early mobilization after fixation of proximal tibial fractures helps in improving the range of motion of the knee joint, strength of quadriceps and hamstrings. Since Schatzker type V and VI are more severe than Schatzker type I and II the outcome also varied with improved results seen in later than the former.

Keywords: Proximal tibial fractures; early mobilization; range of motion.

1. INTRODUCTION

Among all the fractures in the body, tibia is commonly involved in injuries. Increase in road traffic accidents, factory accidents and high velocity trauma are the leading cause for proximal tibial fractures. To preserve normal knee function after surgery, it is mandatory to maintain joint congruity, preserve the normal mechanical axis, ensure joint stability and restore full range of motion especially in Indian culture where squatting and sitting cross leg is routine. The meticulous planning with x rays, Computed Tomography scan of the knee joint with three-dimensional (3D) reconstruction should be obtained to identify the size, location, intraarticular extension of fracture line or depression of the fragment. Aim of this study is to assess role of early mobilization in proximal tibial fractures with conventional methods of physiotherapy in improving ROM, in assessing loss of fixation and post operative complications. Tibial plateau fractures result from direct axial compression, usually associated with a valgus (more common) or varus (less common) component, and an indirect shearing force. With the knee in full extension, the force generated by the injury drives the femoral condyle into the Tibial plateau. The direction, magnitude and location of the force, the position of the knee at the time of impact, and the quality of the bone determine the fracture pattern. Isolated avulsion fractures of the anterior cruciate ligament are not considered fractures of the tibial plateau. Conservative management can be an alternative in simple fractures and in those patients who are unfit for surgery [1]. But the complex fractures if treated with cast and traction always has been ineffective in reducing and maintaining the congruency of the articular surface causing multi-axial malalignment leading to mal-union and secondary degenerative arthritis [2]. Also, to add few more are soft tissue contractures, joint stiffness, quadriceps and hamstring weakness and atrophy. Hence the concept of Biological Fixation was developed to avoid above complications [3,4]. The objective of biological fixation is to assist the physiological process of bone healing optimally with a minimal amount of operative intervention. Unfortunately, the effect of

early mobilization on the internal fixation of the fracture construct has been not studied widely for known or unknown reasons. This study projects that "early the rehabilitation started, early is the recovery" in terms of range of motion, decreasing the post operative stiffness and improving the osteogenesis as per Wolff's law. Earlier, the period of immobilization was minimum for 6 weeks in the fear of loss of fracture reduction and implant failure which affected negatively to the patient in terms of increasing hospital stay, financial burden, work absenteeism and overall decreasing the morale of the patient. Problem increases when additional complications like infection steps in, which increases the socioeconomic burden. Hence the concept of early initiation of physiotherapy was considered [1]. There is no structured protocol especially designed for proximal tibial fracture rehabilitation after surgery. However, in our study we followed a set of exercises as described further. Over a period of time there were lots of development in the management of proximal tibial fractures which drastically reduced the intraoperative and post operative complication. Better understanding of biomechanics, quality of implants, principles of internal fixation, soft tissue care, antibiotics and asepsis have all made possible the internal fixation of fractures as an acceptable mode of treatment [5,6]. However, the treatment doesn't end here until surrounding soft tissue are healthy and joints are mobile which eventually helps to bring the affected knee within normal range. After the popularity of Minimally Invasive Percutaneous Plate Osteosynthesis Techniques using conventional implants which were placed through submuscular plane, the soft tissue handling was less with advantage of preservation of periosteal blood supply and hence the acceleration of fracture healing [7]. Minimally invasive percutaneous plate osteosynthesis technique helped to mobilize early and restore joint function [8]. Most of the information fetched here is from the literature available today but still it is very less in terms of quantity and quality. Unfortunately, not much attention had been focused on post operative rehabilitation after proximal tibial fracture fixation and its role to achieve the goal of pre injury level.

2. METHODS

This study is prospective observational hospital-based study conducted from 01 July 2017 to 31st Dec 2018. Total 68 patients were included in the study. Inclusion criteria was a) patient who underwent surgical modality of treatment for particular proximal tibial fractures which were classified as per Schatzker's classification. b) Patients above 18 years of age with proximal tibial fractures with ≥ 5 mm displacement, depression or step. Exclusion criteria was a) age less than 18 years b) open fractures c) displacement, depression or step < 5 mm d) fractures with neurovascular injuries e) floating knee injuries f) compartment syndrome g) Other significant injuries which hampers post operative recovery and ambulation (polytrauma). The study was approved by the local ethical committee and subjects were enrolled in a clinical protocol reviewed and approved by the institution. Before the beginning of the study, written informed consent was obtained from each patient. Patients who were included in the study as per inclusion criteria but who were having tense hemarthrosis, underwent aspiration of the joint under aseptic precautions. For these patients damage control orthopedics was followed. They were operated between day 5 to day 9 depending on the skin condition. Preoperatively all the patients were given cryotherapy, limb elevation on Bohler Braun splint till they are posted for surgery. Neurovascular status was assessed and confirmed with relevant investigation in case of doubt. Post operatively patients were given three doses of intravenous antibiotics and appropriate analgesics. After three doses of intravenous antibiotics patients were shifted to oral formulations. From day 2 to day 5 patients were educated and were subjected to conventional

physiotherapy protocol followed at our center for quadriceps strengthening exercises (Table 1). It was supervised till the patient is in hospital and on discharge (day 4-5) they were educated to practice at their residence with special emphasis on non-weight bearing. Appropriate postoperative x-rays were taken to assess the reduction. First follow up was after 3 weeks post-surgery for suture removal, assessment of improvement in knee range of motion, any post operative complications like gaping of wound, discharge, signs of infection, loosening of implant etc. and second follow up was after 6 weeks post operatively. To allow weight bearing and its effect on Knee biomechanics is out of the scope of this study. We allow weight bearing after 8-10 weeks post operatively depending upon fracture union clinically as well as radiologically; however, there are studies who also claim that early and supervised weight bearing improves postoperative results in terms of proprioception and over all soft tissue healing and osteogenic remodeling. This protocol was started as early as on day 2 – day 3 and as late as day 4 depending on the pain tolerance of the patient.

Note: In all strengthening exercises, each contraction and stretching were held for 10 - 15 seconds followed by a 3 seconds rest and repeated 10 times. The knee flexion was in the initial period is expected less due to surgical trauma and that was well educated to the patient.

Data was tabulated and analyzed using SPSS (statistical packages for social sciences) version 20 software. Data was presented using suitable diagrams and mean, standard deviation, chi square test etc. were applied on the data. Descriptive statistics were done for all the variables.

Table 1. Inclusion and exclusion criteria

Inclusion Criteria:	Exclusion Criteria:
<ul style="list-style-type: none"> • Patients who underwent surgical modality of treatment for the particular proximal tibial fracture classified as per Schatzker's classification. • Patients above 18 years of Tibial plateau fractures with 5mm or more displacement or depression or step 	<ul style="list-style-type: none"> • Age less than 18 years, • All open fractures • Fractures with < 5mm displacement or depression or step • Fractures with neuro vascular injury • All floating knee injuries. • Compartment syndrome • Other injuries which hampers post op recovery and mobilization (polytrauma)

3. RESULTS

There were 68 patients enrolled in the study as per inclusion and exclusion criteria with 53(77.9%) males and 15(22.1%) females. The youngest patient in our study was 20 years old and oldest was 59 years of age with 33% of the patients belonged to age group of 30-39 years with significant male preponderance. Majority of the cases were type V and VI with the total of 29(42.65%) cases and least were type II cases making a total of 6(8.8%). Type 1 were 14(20.6%) and type III and IV were 10(14.7%) and 9(13.2%) respectively. On follow up at 3rd week maximum number to patient have achieved the goal of improvement in range of motion with an average of 10°-15°/ week with the exception of few who were non-compliant. Distribution of patient in terms of range of motion at 3rd and 6th week are shown below (Fig. 1). While at 6th weeks total 13(19.1%) patients have achieved full range of motion and 9(13.2%) patients were able to reach the target of 100° (Fig. 2). We used variety of implants depending on fracture geometry and displacement to achieve and maintain reduction and stable fixation. We also have used combination of implants in 7(10.3%) cases. The association between early mobilization on range of motion at 3rd week and choice of implant was found to be statistically significant with chi square value = 32.435, df = 20 and p value = 0.039 (p value < 0.05). However due to small sample size (n=68) and expected count of <5 was observed in 29 cells. The statistical association may not be the true representation of the real value. (Chi square =50.188, df =25, p value =0.002) (Table 2). At the end of 6th week, the association between choice of implant and range of motion was significant (Chi-square = 63.568, df = 20, p value=0.000). Expected count of <5 was

observed in 26 cells (Table 3). To calculate the association between early mobilization at 3rd week and type of fracture the various cells were clubbed together. Range of motion at 3rd week 40°, 50° and 60° were clubbed in one cell and 70°, 80° and 90° were clubbed in another cell. The type of fracture I, II and III were clubbed in one cell based on lateral side fixation and type IV, V and VI other than only lateral side fixation were clubbed based on the side used for stabilization of the fracture (Table 4). The association between the two was found to be statistically significant with chi square value 10.9487, df = 1 and p value = 0.000937 (<0.05). To calculate the association between early mobilization at 6th week and type of fracture the various cells were clubbed together. Range of motion at 6th week 100°, 110° and 120° were clubbed in one cell and 130° and 140° were clubbed in another cell. The type of fracture type I, II and III were clubbed in one cell based on lateral side fixation and type IV, V and VI other than only lateral side fixation were clubbed based on the side used for stabilization of the fracture (Table 5). The association between the two was found to be statistically significant with chi square value 39.7338, df = 1 and p value = 0.001.

There was no statistically significant association found in following parameters: -

- a) Age and early mobilization
- b) Loss of fixation and early mobilization

The above findings conclude that early mobilization can be safely carried out irrespective of age and without fear of loss of fixation. In our series only one patient had screw back out which led to condylar widening on his follow up at 6 weeks.

Table 2. Physiotherapy protocol

Phase 1	Immediate post op period (0-1 weeks)
Rehabilitation goals	<ul style="list-style-type: none"> • Reduce swelling and oedema • Minimize pain • Start knee flexion as per pain tolerance. • Knee flexion to be gained 100-150/week and strict non weight bearing.
Actions / Interventions	<p>Swelling management –</p> <ul style="list-style-type: none"> • Ice pack application and elevation • Active ankle pumps • Long extension knee brace <p>Range of Motion / mobility</p> <ul style="list-style-type: none"> • Knee passive range of motion and active assisted range of motion • Static quadriceps exercises

Phase 2	1-3 weeks
Rehabilitation goal	<ul style="list-style-type: none"> • Improve knee flexion further with a target of 100-150 /week • Improve strength of quadriceps and hamstrings by resisted exercises and dynamic strengthening exercise by application of weights. • 90° of flexion to be achieved by end of 3 weeks
Action / Interventions	<ul style="list-style-type: none"> • Continue static quadriceps exercise and with weights applied. • VMO exercises • Hip abductor strengthening exercises and resisted quadriceps exercise. • SLR with brace locked in full extension. • Patella and soleus stretching exercises • Continue ice pack application if swelling persist. Continue to remain non weight bearing
Phase 3	3-6 weeks
Rehabilitation goals	Further increase in knee ROM to normal or near normal (up to 140°). Non weight bearing till further advice
Actions / Interventions	Repeat as phase 2 with increase in frequency and repetitions.

Table 3. Assessment of knee ROM at end of 3 weeks post operatively with various implants

ROM at 3rd week (in degrees)	Cancellous Cannulated screws	Dual plating	Lateral plating	Medial plating	Combined	MIPPO	Total
40	0	1	0	0	0	0	1
50	0	4	0	0	0	1	5
60	0	6	0	0	0	2	8
70	1	8	0	2	3	4	18
80	6	2	3	2	1	1	15
90	12	0	3	3	3	0	1
Total	19	21	6	7	7	8	68

Table 4. Assessment of knee ROM at end of 6 weeks post operatively with various implants

ROM at 6th wk (in degrees)	Cancellous Cannulated screws	Dual plating	Lateral plating	Medial plating	Combined	MIPPO	Total
100	0	7	0	0	0	2	9
110	0	7	0	0	0	2	9
120	1	7	0	2	4	4	18
130	10	0	3	4	2	0	19
140	8	0	3	1	1	0	13
Total	19	21	6	7	7	8	68

Table 5. Association between early mobilization in type of fractures and ROM at 3rd week

ROM at 3rd week	Type I , Type II and Type III	Type IV, Type V and Type VI	Total
40 ,50 and 60	01	29	30
70, 80 and 90	14	24	38
Total	15	53	68

Chi square value 10.9487, df = 1 and p value = 0.000937 (<0.05).

Table 6. Association between early mobilization in type of fractures and ROM at 6th week

ROM at 6th week	Type I , Type II and Type III	Type IV, Type V and Type VI	Total
1000,1100 &1200	03	27	30
1300 and 1400	33	5	38
Total	36	32	68

Chi square value 39.7338, df = 1 and p value = 0.001

Table 7. Stretching techniques

Static (most common)	Dynamic	Pre-Contraction
<ul style="list-style-type: none"> • Active • Passive 	<ul style="list-style-type: none"> • Active • Ballistic 	PNF and other Techniques




Fig. 1. Distribution of patients according to (ROM) at 3rd week




Fig. 2. Distribution of patients according to (ROM) at 6th week



Fig. 3. Schatzker's type II, ROM at end of 3 weeks post operative



Fig. 4. Schatzker's type V, ROM at end of 3 weeks

4. DISCUSSION

Tibial plateau fractures are difficult to treat because of their intra-articular nature, cancellous bone involvement, and proximity to a major weight bearing joint. Among the total number of patients males' predominance was noted due to various reasons one of which that the use of vehicles was mostly done by them [9]. This finding was matching with the findings of Rabari Y D, Somani A H et.al. Open reduction and internal fixation are required for depressed or displaced and unstable fractures to regain early and complete range of motion. There is multiple choice of implants which can be used depending on fracture geometry [7]. Proper physiotherapy and compliance of patient are equally important to achieve good results. Keeping these points in mind, the present study was carried out with the purpose to assess the functional outcome of various modalities of surgical treatment in different types of tibial plateau fractures and to compare our results with literature. There is no particular number or figure on the amount of articular depression that can be accepted. However as per Honkonen et.al anything more than 3 mm intra articular depression or more than 5 mm condylar widening is an indication for operative intervention [10,11]. In our study, the indications for the surgery were either 5 mm or more displacement, articular depression or step. Passive and active mobilization has shown to benefit the injured soft tissue (surgical or traumatic insult) by increasing local blood supply of tissue by mobilization [12,2]. This helps in providing nutrition to the tissues necessary for repair. By stretching of the knee joint gradually up to the tolerable limits during early post operative days, the contractile proteins of the muscle fibers also remain viable. Ultimately it leads to suppleness of the joint there by increasing its range of motion and preventing over all stiffness and chances of deep vein thrombosis. On the other side immobilizing a particular joint causes adverse effect on internal milieu like synovial adhesions, disorganization of collagen fibrils and increase in collagen degradation, alteration of matrix turnover with increase in matrix degradation than formation which increases the stiffness of the joints. To add further more it decreases water content, glycosaminoglycans and degree of orientation of matrix collagen fibrils within ligaments.

Prolonged immobilization also increases subperiosteal osteoclastic resorption at fibro-osseous junction. It decreases tensile strength of

the bone ligament interface affecting the healing of the surgical construct and also increases chances of deep vein thrombosis in high-risk patients. Hence by providing early mobilization overcomes the above cascade, thereby improving the chances to achieve pre-injury level. This is particularly for high performing athletes who desires to return to normal sports activity. The type of implant was decided based on fracture geometry, intra articular comminution (step/depression) and mediolateral involvement. Skin condition around the involved knee before surgery was one of the most important factors. As per study conducted by Dr. Ramnath et.al for Functional Outcome of Proximal tibia fractures treated with locking compression plates shown excellent results with less complications. With the use of locking plates, the aim to achieve anatomical intraarticular reduction is easy. Furthermore, it helps in initiation of early mobilization for targeting good functional outcome in tibial condylar fractures [13,14]. This theory was supported by our study and similar results were obtained with respect to fracture reduction, stabilization and which eventually improved range of motion. Similar results were obtained at the retrospective study done at Goa medical college on Outcome of Proximal Tibia Fractures Treated with Dual Plating [15,16]. When a skeletal muscle with an intact nerve supply is stretched, it contracts which is called as normal stretch reflex. During stretch of the muscle, more the muscle fibers that are stretched more the length is achieved. During physiotherapy the rationale behind to hold the stretch for certain period is to recruit more stretch receptors to allow greater lengthening of the muscles. The Golgi tendon organ also helps to relax the stretched muscle through its function. In the initial phases of training of muscle, the mechanical adaptations occur that are followed by neural adaptations. Theoretically stretching techniques are mainly three types (Table 6). Static stretching- application of low force with long duration (30 sec). It is slow controlled movement with importance on postural awareness. Dynamic stretching – application of controlled movement through available range of motion. In ballistic type forcing a part of the body beyond its range of motion. Not practiced due to risk of injury. Precontraction stretching – Proprioceptive neuromuscular facilitation stretching (PNFS) -it is a type where contraction of the muscle being stretched or its antagonist before stretching. It triggers the inverse stretch reflex which prevents injury to the involved muscle.

5. CONCLUSION

Outcome of early mobilization in proximal tibial fractures is strongly depends on thorough understanding fracture geometry, classifying the type of fracture, understanding the biomechanics and choosing the suitable implant. The role of less tissue handling during surgery with stable fixation also helps in stability of the construct and better healing potential. Conventional methods of physiotherapy for quadriceps strengthening exercises, when instituted early has proven to be beneficial in improving range of motion to normal in Schatzker's type I, type II and type III and to near normal for Schatzker's type IV, type V and type VI. Regular physiotherapy advised helps preventing disuse atrophy of quadriceps muscle and also helps in maintaining tone of the muscles.

6. LIMITATIONS

For this study we followed up patients till the end of 6 weeks only so the post operative weight bearing was not considered in our study. For more detailed study longer follow up is required.

CONSENT

It is not applicable.

ETHICAL APPROVAL

The study was approved by the institutional ethics committee, Seth Nandlal Dhoot Hospital, Aurangabad.

ACKNOWLEDGEMENT

Dr. Krishnakant B Patil, MD Physiology, Professor and Dean Smt. Kashibai Navle Medical College, Pune for practical aspects on muscle physiology. Dr. (Lt Col) Shruti Vashisht, MD PSM for helping with statistical inputs and information.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Shrestha R, Kandel M, Gupta HK, Shrestha SK, Dongol S, Hamal RR. A study of conservative management of

- tibial plateau fractures. Journal of College of Medical Sciences-Nepal. 2016;12(1):5–9.
2. Koval KJ, Sanders R, Borrelli J, Helfet D, DiPasquale T, Mast JW. Indirect reduction and percutaneous screw fixation of displaced tibial plateau fractures. J Orthop Trauma. 1992;6(3):340-6. DOI:10.1097/00005131-199209000-00012 PMID: 1403254.
3. Schatzker J, McBroom R, Bruce D. The tibial plateau fracture: the Toronto experience:1968–1975. ClinOrthopRelat Res. 1979;(138):94–104.
4. Tscherner H, Lobenhoffer P. Tibial plateau fractures. Management and expected results. Clin Orthop Relat Res. 1993; 292:87-100.
5. Kulwinder Singh, Manjeet Singh, Rakesh Gautam. Functional Outcome of Surgical Management of Tibial Plateau Fractures: Case Series of 30 Cases, International Journal of Science and Research (IJSR). 2015;4(9):905–909.
6. Reddy RS, Kumar CY, Shah HM, Golla DK, Ganesh DJN, Kumar PA. Evaluation of tibial condyle fractures treated with Ilizarov fixation, a prospective study. J Clin Diagn Res. 2014 Nov;8(11):LC05-7. DOI: 10.7860/JCDR/2014/9564.5117 Epub 2014 Nov 20. PMID: 25584250; PMCID: PMC4290269.
7. Smit Shah, Pulkit Maniar, Niravkumar Moradiya. Management of fractures of proximal tibia by various treatment modalities: A study of 56 patients. Int J Orthop Sci. 2017;3(1):346-350. DOI: 10.22271/ortho.2017.v3.i1f.51
8. Arumugam S, Arumugam V, Raviraman V. To analyze the functional outcome of proximal tibial fractures treated with minimally invasive percutaneous plate osteosynthesis technique. Int JH Res Orthop. 2017;3:1205-10.
9. Rabari YB, Somanni AM, Prasad DV, Thadeshwar KH. Various modalities of treatment of proximal tibial fractures: A prospective longitudinal study. Int J Res Orthop. 2017;3:259-65.
10. Bowes DN, Hohl M. Tibial condylar fractures. Evaluation of treatment and outcome. Clin Orthop Relat Res. 1982 Nov-Dec;(171):104-8. PMID: 7140056.

11. Honkonen SE. Indications for surgical treatment of tibial condyle fractures. Clin Orthop Relat Res. 1994 May;(302):199–205.
12. Iliopoulos E, Galanis N. Physiotherapy after tibial plateau fracture fixation: A systematic review of the literature. SAGE Open Med. 2020 Oct 14;8: 2050312120965316.
DOI: 10.1177/2050312120965316
PMID: 33133602; PMCID: PMC7576901.
13. Ramnath DS, et al. A study of functional outcome of proximal tibial fractures treated with L.C.P IOSR Journal of Dental and Medical Sciences (IOSR-JDMS). 2016;15(6):01-02.
e-ISSN: 2279-0853, p-ISSN: 2279-0861.
14. Segal D, Mallik AR, Wetzler MJ, Franchi AV, Whitelaw GP. Early weight bearing of lateral tibial plateau fractures. Clin Orthop. 1993;(294):232-37.
15. Nagendra Singh, et al. Outcome of Proximal Tibia Fractures Treated With Dual Plating - A Retrospective Study 3IOSR Journal of Dental and Medical Sciences (IOSR-JDMS). 2018;17(8 Ver. 13):61-71.
e-ISSN: 2279-0853, p-ISSN: 2279-0861.
16. Rohra N, Suri HS, Gangrade K. Functional and Radiological Outcome of Schatzker type V and VI Tibial Plateau Fracture Treatment with Dual Plates with Minimum 3 years follow-up: A Prospective Study. J Clin Diagn Res. 2016 May;10(5):RC05-10.
DOI: 10.7860/JCDR/2016/18732.7855
Epub 2016 May 1.
PMID: 27437315; PMCID: PMC4948491

© 2022 Gursale et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/84480>*