



Evaluation of DSSAT-CROPGRO Module for Spatial Yield Estimation of Chickpea in Vidisha (Madhya Pradesh) and Nagaur (Rajasthan), India

A. Ragul ^a, S. Pazhanivelan ^{b*}, A. P. Sivamurugan ^b,
K. Boomiraj ^a, K. P. Raghunath ^b and Kancheti Mrunalini ^{b,c}

^a Agro Climate Research Centre, Tamil Nadu Agricultural University, Coimbatore – 641003, India.

^b Centre for Water and Geospatial Studies, Tamil Nadu Agricultural University, Coimbatore – 641003, India.

^c Indian Institute of Pulses Research, Kanpur, Uttar Pradesh-208024, India.

Authors' contributions

This work was carried out in collaboration among all authors. Authors AR and SP carried out verification of output, and wrote the protocol and the first draft of the manuscript. Authors APS and KB generated the outputs. Authors KPR and KM defined the methodology of the research and verification part. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i92461

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/103827>

Original Research Article

Received: 23/05/2023

Accepted: 25/07/2023

Published: 29/07/2023

ABSTRACT

Crop simulation models are often used to characterize, develop and assess field crop production practices. The present study was carried out for chickpea spatial yield estimation at Vidisha of Madhya Pradesh and Nagaur district of Rajasthan employing the DSSAT model. In this study, the DSSAT-CROPGRO module was used to estimate chickpea yield during *rabi* 2022. To simulate the yield, DSSAT required datasets of crop growth and management, daily weather data, and soil data were provided. The simulated yield was validated using the observed yield through CCEs from

*Corresponding author: E-mail: pazhanivelans@gmail.com;

farmers' fields. When the observed and simulated yields were compared, their deviation was found to be less than 20 percent for all varieties at experimental locations of the Vidisha and Nagaur districts. The observed yield of Chickpea matched well after calibration which showed that model could simulate the yield with high accuracy as it showed R^2 , d , and MAPE of 0.87, 0.92, and 7.30 for calibration and 0.88, 0.90, and 7.60 for the validation, respectively. The model has been successfully calibrated and validated for the chickpea at spatial level and it can be taken for further applications in natural resources management and climate change impact studies.

Keywords: Chickpea; DSSAT; CROPGRO; spatial yield; Vidisha; Nagaur.

1. INTRODUCTION

“Chickpea (*Cicer arietinum* L.) is grown over a wide range of agroclimatic environments. It is traditionally grown in the northern hemisphere mostly between 20°N and 40°N latitude. Most of the desi (with yellow to brown seed testa) chickpea is grown between 20°N and 30°N, while kabuli (with cream-colored seed testa) types are grown above 30°N. In addition, there is a small area between 10°N and 20°N at relatively high elevations in India and Ethiopia where it is grown” [1].

“Crop simulation models involve the mathematical function of various crop physiological factors such as photosynthesis, respiration, and relative growth rate to describe the crop growth changes under various climatic and environmental conditions. The model at times becomes complicated as it needs several detailed inputs for simulation and makes the calibration process tedious to perform” [2]. “Crop simulation models are key components to test the advances in agricultural technology and to predict crop responses to present and future climate forcing. These models are being used widely to estimate the crop production potential, transfer Agro-technologies, assist strategic decisions, and forecast real-time yield” [3]. “DSSAT model is one of the crop simulation models used to simulate the growth and development of a crop by integrating soil, crop phenotype, weather, and management options” [4]. “DSSAT has modules that allow users to build model input files for spatial simulations across predefined management zones, and calibrate the model to simulate historic spatial yield variability and crop response to environmental and management variations” [5]. “The capability of the DSSAT model in simulating crop responses and the sensitivity of the model output to input parameters with spatial attention to the determinants of the model response to the practice of conservation agriculture were analyzed. The results showed that the

phenological cultivar parameters were the most influential model parameters. The correlation between the input parameters and output variables was stable over a wide range of seasonal rainfall conditions” [6].

CERES-Maize module was employed to estimate maize yield spatially during *kharif*, 2017 at Ariyalur and Perambalur districts which indicates that the module can be used to estimate the maize yield spatially at different weather, and soil conditions [7]. Similar to CERES-Maize, the CROPGRO-Chickpea module is a dynamic simulation module, which replicates the growth and yield of a variety of leguminous crops, including chickpea, soybean, peanut, and groundnut. In India, Singh and Virmani *et al.*, (1996) created “the CHIKPGRO (CROPGRO-Chickpea) model, which can be used to predict the potential and water-limited yields of chickpeas” [8]. “Using the DSSAT-CROPGRO model in the CDR, the future climate change impact of black gram yield has been evaluated, the results show that the simulation of black gram yield change for RCP 4.5 might be around 34%, 52%, and 25% during the near-century, mid-century, and end-century, respectively” [9]. “Calibration and validation were done for DSSAT (v.4.6) CROPGRO – chickpea model was used to study the impact of climate change by its at Anand. Thus, the higher temperature regimes resulted in a gradual decrease in grain yield whereas lower temperature showed an increase in grain yield” [10].

The present study utilizes the CROPGRO-chickpea module for estimating the yield of chickpeas at spatial levels at different locations, weather, soil, and management conditions.

2. METHODOLOGY

2.1 Study Area

Geographically, the experiment was carried out for the rabi season (2022) in the Vidisha district

of Madhya Pradesh and Nagaur of Rajasthan (Figs. 1 & 2). Vidisha is situated at 23° 31' N latitude, 77° 49' E Longitude, and at an altitude of 429 meters above mean sea level (MSL) in the eastern part of the fertile Malwa Region. Similarly, Nagaur is situated at 27° 12' N latitude, 73° 44' E Longitude, and at an altitude of 302 meters above mean sea level (MSL) in the northwestern Marwar region of Rajasthan. Vidisha district enjoys a subtropical climate and receives an average annual rainfall of about 1299 mm. On average, about 85 percent of the total rainfall is received during the South-West monsoon period i.e., from June to September. However, occasionally 5 to 10 percent showers occur during the winter season, whereas Nagaur district enjoys the desert climate and receives an

average annual rainfall of about 307 mm. The experimental study was carried out in different locations in the Vidisha and Nagaur districts. In each of the districts, around 70 ground truth points and Crop Cutting Experiments (CCE) data were collected at random locations from which 20 monitoring sites for Vidisha and 17 monitoring sites for Nagaur were selected for the experimental study. The package of practices for the cultivation of chickpeas was followed as per the recommendation. The crop parameters such as yield and yield attributes, LAI, harvest index, and phenology were used for calibration of the DSSAT v.4.8 model. Three varieties were chosen for Vidisha (JG 16, RVG 202, JAKI 9218) and three varieties were chosen for Nagaur (RSG 44, RSG 896, RSG 807).




Fig. 1. Study area location of Vidisha (Madhya Pradesh)




Fig. 2. Study area location of Nagaur (Rajasthan)

2.2 CROPGRO Module Input

The pertinent daily weather, soil, and crop management data for all the sowing dates and irrigation levels were used as input and experiment performance data files. The daily weather data on minimum and maximum temperature (°C), solar radiation (MJm⁻²day⁻¹), and rainfall (mm) were collected for the study area. The weather input files for crop simulation were generated using the weatherman tool in DSSAT for monitoring sites and the soil files are gathered from the International Research Institute for, Climate Society, Michigan State, University Harvest Choice, International Food Policy Research Institute at 1:10,000 scale with a 5-min resolution. These files were the inputs to the model for the monitoring fields for the study area to be simulated.

2.3 Calibration and Validation of the CROPGRO Chickpea Model

For calibration, the yield data from the three different monitoring sites for three different varieties (JG 16, RVG 202, JAKI 9218) in Vidisha and three different monitoring sites for three different varieties (RSG 44, RSG 896, RSG 807) in Nagaur were used for *rabi* season 2022 - 23. The remaining 17 monitoring sites for three varieties in Vidisha and 14 monitoring sites for three varieties in Nagaur were used to validate the model using the yield data collected from farmers' fields through CCEs (observed yield data) in the study area.

2.4 Statistical Approach of Model Evaluation

The model's performances were evaluated by using the statistical indices including the coefficient of determination (R²), Mean Absolute Percentage Error (MAPE), Nash-Sutcliffe efficiency (NSE), and Index of Agreement (d) [11].

2.4.1 Pearson's correlation coefficient (r) and coefficient of determination (R²)

$$R^2 = 1 - \frac{RSS}{TSS}$$

R² = Coefficient of determination
 RSS = Sum of squares of residuals
 TSS = Total sum of squares

2.4.2 Index of agreement (d)

$$d = 1 - \frac{\sum_{i=1}^n (n_i - s_i)^2}{\sum_{i=1}^n (|M_i - \bar{M}| + |S_i - \bar{M}|)^2}$$

where,

M_i and S_i are the observed and simulated values, respectively.

n = the number of observations

M = the mean of n measured values

2.4.3 Mean Absolute Percentage Error (MAPE)

$$M = \frac{1}{n} \sum_{t=1}^n \left| \frac{A_t - F_t}{A_t} \right|$$

M = mean absolute percentage error

N = number of times the summation iteration happens

A_t = actual value

F_t = forecast value

3. RESULTS AND DISCUSSION

3.1 Chickpea Genetic Coefficients

The calibration of the CROPGRO - Chickpea module, data on plant growth and development, soil characteristics, weather, and crop management were collected as required for determining the cultivar coefficients of JG 16, RVG 202, JAKI 9218, RSG 44, RSG 896 & RSG 807 following the procedures described in International Benchmark Sites Network for Agrotechnology Transfer (IBSNAT). "These coefficients allow the model to simulate the performance of diverse genotypes under different soil, weather, and management conditions" [12].

To determine the genetic coefficients (Table 1) of chickpeas, the calibrated values are obtained by changing their values to determine the variation in the magnitude of output manually. Then, those values of the genetic coefficients that were found most realistically simulated the growth and yield of chickpeas were selected. The data set for genetic coefficients calculations include days to anthesis, days to the first pod, days to physiological maturity, days to harvest maturity, seed yield, by-product leaf area, and harvest index. The procedure for determining genetic coefficients involved in running the model using a range of values of each coefficient, until the desired level of agreement between simulated and observed values was reached.

Table 1. Calibrated genotypic coefficients for rabi Chickpea cultivar

Coefficient code	Description	Genetic coefficient					
		JG 16	RSG 44	RSG 896	JAKI 9218	RVG 202	RSG 807
CSDL	Critical Short-Day Length below which reproductive development progresses WITH daylength effect (for long day plants) (hour)	11	11	10.9	10.9	10.9	11
PPSEN	The slope of the relative response of development to photoperiod with time (negative for long-day plants) (1/hour)	-0.143	-0.143	-0.32	-0.32	-0.143	-0.43
EM-FL	The time between plant emergence and flower appearance (R1) (photothermal days)	31	42	35	40.8	32.8	36
FL-SH	The time between the first flower and the first pod (R3) (photothermal days)	5.5	10.5	7.7	8	5	6.5
FL-SD	The time between the first flower and the first seed (R5) (photothermal days)	13	13.5	14.5	13	9	11.1
SD-PM	The time between the first seed (R5) and physiological maturity (R7) (photothermal days)	30.5	50	45	41	35	39.7
FL-LF	The time between the first flower (R1) and end of leaf expansion (photothermal days)	34	60	53	49	45	45
LFMAX	Maximum leaf photosynthesis rate at 30 C, 350 vpm CO ₂ , and high light (mg CO ₂ /m ² /s)	0.95	1.3	1	0.95	1.2	1.2
SLAVR	Specific leaf area of cultivar under standard growth conditions (cm ² /g)	200	200	200	200	220	200
SIZLF	Maximum size of the full leaf (three leaflets) (cm ²)	10	10	10	10	10	10
XFRT	The maximum fraction of daily growth that is partitioned to seed + shell	0.96	1.05	1	0.95	0.95	0.95
WTPSD	Maximum weight per seed (g)	0.17	0.32	0.18	0.18	0.22	0.19
SFDUR	Seed filling duration for pod cohort at standard growth conditions (photothermal days)	22	28	22	20	20	20
SDPDV	Average seed per pod under standard growing conditions (#/pod)	1.2	1.6	1.3	1.3	1.5	1.2
PODUR	Time required for cultivar to reach final pod load under optimal conditions (photothermal days)	18	19	18	18	16	16
THRSH	The maximum ratio of (seed/ (seed + shell)) at maturity.	82	85	85	85	85	85
SDPRO	Fraction protein in seeds (g(protein)/g(seed))	0.216	0.216	0.216	0.216	0.216	0.216
SDLIP	Fraction oil in seeds (g(oil)/g(seed))	0.48	0.048	0.048	0.048	0.048	0.048

3.2 Yield Analysis

The observed yield and simulated yield are compared, and their deviation is calculated. It indicates that the deviation is less than 20 per cent for all varieties of chickpea in 20 experimental locations of Vidisha district (Table 2). In Nagaur district, the deviation is less than 18 per cent for all varieties of chickpea in 17 experimental locations (Table 3). For all varieties in Vidisha and Nagaur districts, the simulated yield fitted well with the observed yield although the simulated yield was slightly higher than the observed values. Choudhury et al., [13] also stated that the simulated biomass yield was slightly higher than that of the observed biomass yield in wheat while using the DSSAT model.

3.3 Calibration and Validation of Model

The observed yield of Chickpea matched well after calibration which showed that model could simulate the yield with high accuracy, as it showed R^2 , d and MAPE of 0.87,0.92 and 7.30 for calibration and 0.88,0.90 and 7.60 for the validation respectively, which indicates a good prediction efficiency. The R^2 values are good and significant, the 1:1 line graph was drawn showing observed yield in X-axis and simulated yield in Y-

axis. The regression line of grain yield was near the 1:1 line, indicating that the model was performing well under the test environment, thus model simulated the yield perfectly (Figs. 3 & 4).

The index of agreement (d) ranges around 0.90, which indicates all varieties have a high index of agreement (i.e., perfect match). Thus, The MAPE is very good for all varieties ($< 10\%$) thus, the mean absolute percentage errors between the predicted and actual values in the calibration and validation of the model were less and the model predicted the yield well and which indicates a very good score for the simulation. The MAPE was good for evaluating the yield and growth of wheat in Algeria [14].

3.4 Correlation Matrix of Weather Variables with Yield

The climatic conditions of the experimental area are taken and analyzed for predicting their correlation matrix. The Minimum dataset of weather parameters that are required for DSSAT (Maximum Temperature, Minimum Temperature, Solar Radiation and Rainfall) is chosen. The correlation matrix using R-Studio indicates that the yield when correlated with the TMax (Maximum Temperature) indicated a positive

Table 2. Observed and simulated yield for Vidisha monitoring sites

Vidisha				
Latitude	Longitude	Observed Yield (kg/ha)	Simulated Yield (kg/ha)	Deviation (%)
Variety I - JG 16				
23.51063	78.06736	1464	1536	4.9
23.71673	77.86417	1655	1996	19.6
23.72553	77.83993	1995	2011	0.8
23.68868	78.00718	1765	1904	7.9
23.70251	77.99177	1318	1366	3.6
24.06780	77.91728	1427	1537	7.7
23.63285	77.84866	1185	1238	4.5
Variety II - JAKI 9218				
23.71113	77.90449	1750	1809	3.4
23.48960	78.05553	1834	1916	4.5
23.47606	78.03633	1812	1922	6.1
23.70830	77.93528	1969	2041	3.7
24.02634	77.86156	1703	1863	9.4
23.39478	78.01640	1684	1790	6.3
Variety III - RVG 202				
23.67509	77.97400	1492	1633	9.5
23.73018	77.86488	1585	1689	6.6
23.58353	77.98783	1612	1715	6.4
23.53913	78.02460	1505	1670	11.0
24.00687	78.14264	1320	1550	17.4
24.06878	77.74187	1386	1570	13.3
23.54101	77.98305	1198	1260	5.2

Table 3. Observed and simulated yield for Nagaur Monitoring Sites

Nagaur				
Latitude	Longitude	Observed Yield (kg/ha)	Simulated Yield (kg/ha)	Deviation (%)
Variety I - RSG 44				
26.79998	74.23855	1788	1907	6.7
26.64958	74.27593	1518	1559	2.7
26.82605	74.65694	1645	1897	15.3
26.99160	74.77227	1377	1476	7.2
Variety II - RSG 807				
26.62767	74.19618	1809	1840	1.7
26.64373	74.00105	1695	1994	17.6
26.86517	74.76031	1126	1290	14.6
Variety III - RSG 896				
26.74230	74.40154	1492	1666	11.7
26.78795	74.22688	1259	1341	6.5
26.77226	74.43233	1380	1520	10.1
26.77273	47.43463	1210	1308	8.1
26.75381	74.43324	1380	1459	5.7
26.66367	74.31238	1579	1617	2.4
26.59892	74.17314	1412	1559	10.4
26.61882	74.16225	1477	1559	5.6
26.77100	74.46893	1421	1474	3.7
26.82490	74.65146	1462	1582	8.2




Fig. 3. Simulated and observed yield for calibration

value and significant relationship. It is also noted that the yield when correlated with rain shows a negative and non-significant correlation for Vidisha. For Nagaur, when the yield is correlated with all the parameters it shows a negative value and is non-significant. The correlation with the Tmin (Minimum Temperature) showed a positive correlation (Figs. 5 & 6).

Studies on zoning and agricultural potential of tropical crops in Latin America have relevance

and importance similar to the evaluation of DSSAT-CROPGRO for chickpeas in India. However, there are key differences and possible areas of comparison. The geographical and climatic conditions in Latin America differ significantly from those in Vidisha and Nagaur [15]. Comparing the performance of crop models like DSSAT-CROPGRO across diverse regions can provide insights into their generalizability and adaptability [16]. Latin America hosts a wide range of tropical crops [17,18], each with unique responses to environmental conditions.

Evaluating zoning studies for multiple crops can offer a comprehensive understanding of how well different crops are suited to specific regions compared to a focused evaluation of a single crop like chickpea [19]. Comparing the methodologies and data sources used in Latin American studies with the evaluation in India can highlight the importance of accurate and consistent data for robust crop modeling and decision-making [20]. The socio-economic context in Latin America may differ from that in India, influencing the implementation of agricultural policies and the adoption of technology [21]. Understanding these differences

can aid in tailoring interventions to specific regional needs [22,23].

As the evaluation of the DSSAT-CROPGRO module for spatial yield estimation of chickpeas in Vidisha and Nagaur is highly relevant and important for improving agricultural practices, climate change adaptation, and policy formulation. By comparing these findings with studies on tropical crop zoning in Latin America, we can gain valuable insights into the transferability of crop models across regions and the broader implications of precision agriculture and climate-resilient farming practices.




Fig. 4. Simulated and observed yield for Validation




Fig. 5. Correlation matrix for Vidisha




Fig. 6. Correlation matrix for Nagaur

4. CONCLUSION

The validated outcomes of the DSSAT-CROPGRO Chickpea module revealed that this model simulates the yield attributes more than that of observed crop data. The DSSAT model has proved to be a valuable tool for predicting chickpea yield. Therefore, the validated DSSAT can be further used for applications such as prediction of crop growth, phenology, potential and actual yield, performance of chickpeas under climate change study, etc., The model may also be used to improve and evaluate the current practices of chickpea growth and its management as well as the CROPGRO module can be used to simulate the yield spatially for different conditions of soil, weather and management practices to enhance chickpea production. It could be concluded that the model works well for rainfed growing environments and further, it can be taken for application in natural resource management and climate change impact analysis studies.

ACKNOWLEDGEMENT

The authors are thankful to the Agro Climate Research Centre and Centre for Water and Geospatial Studies for providing the fund through DCM – GIZ scheme to carry out the research work in a project mode and like to extend our sincere thanks to all staff members, seniors and my batchmates for their valuable comments and constructive suggestions on the manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Khanna-Chopra R, Sinha SK. Chickpea: physiological aspects of growth and yield. *The chickpea*. 1987;163-87.
2. Boote KJ, Jones JW, Pickering NB. Potential uses and limitations of crop models. *Agronomy Journal*. 1996 Sep; 88(5):704-16.
3. Bannayan M, Crout NM. A stochastic modelling approach for real-time forecasting of winter wheat yield. *Field Crops Research*. 1999 Jun 1;62(1):85-95.
4. Jones JW, Hoogenboom G, Porter CH, Boote KJ, Batchelor WD, Hunt LA, Wilkens PW, Singh U, Gijsman AJ, Ritchie JT. The DSSAT cropping system model. *European journal of agronomy*. 2003 Jan 1;18(3-4):235-65.
5. Thorp KR, DeJonge KC, Kaleita AL, Batchelor WD, Paz JO. Methodology for the use of DSSAT models for precision agriculture decision support. *Computers and Electronics in Agriculture*. 2008;64(2): 276-285.
6. Marc Corbeels, Guillaume Chirat, Samir Messad, Christian Thierfelder, Performance and sensitivity of the DSSAT crop growth model in simulating maize

- yield under conservation agriculture, European Journal of Agronomy. 2016; 76:41-53.
7. Venkatesan M, Pazhanivelan S. Estimation of Maize yield at spatial level using DSSAT crop simulation model. Madras Agricultural Journal. 2018 Dec 1;1:105.
 8. Singh P, Virmani S. Modeling growth and yield of chickpea (*Cicer arietinum* L.). Field Crops Research. 1996 Jun 1;46(1-3):41-59.
 9. SN AI, KP. Assessing the black gram (*Vigna mungo*) yield due to climate change impacts and its adaptation strategies in the Cauvery delta region of South India. Theoretical and Applied Climatology. 2023 Jun 20;1-4.
 10. Patil DD, Karande BI, Satpute SB, Patel HR. Sensitivity analysis to study the impact of climate change on chickpea using DSSAT (4.6) CROPGRO model over middle Gujarat Agroclimatic region. A Peer-Review. Multi-Disciplinary Int. J. 2018;223.
 11. Willmott CJ. Some comments on the evaluation of model performance. Bulletin of the American Meteorological Society. 1982 Nov;63(11):1309-13.
 12. Hunt LA, Pararajasingham S, Jones JW, Hoogenboom G, Imamura DT, Ogoshi RM. Agron. J. 1993;85:1090–1094.
 13. Jahan MA, Sen R, Ishtiaque S, Choudhury AK, Akhter S, Ahmed F, Biswas JC, Maniruzzaman M, Miah MM, Rahman MM, Kalra N. Optimizing sowing window for wheat cultivation in Bangladesh using CERES-wheat crop simulation model. Agriculture, Ecosystems & Environment. 2018 Apr 15;258:23-9.
 14. Waffa Rezzoug, Benoit Gabrielle, Suleiman A, Benabdeli K. Application and evaluation of the DSSAT-wheat in the Tiaret region of Algeria. African Journal of Agricultural Research. 2008;3(4):284-296.
 15. Olivares Campos, BO. Identification of the Main Soil Variables Correlated with Banana Productivity. In: Banana Production in Venezuela. The Latin American Studies Book Series. Springer, Cham; 2023.
 16. Paredes F, Olivares B, Rey J, Lobo D, Galvis-Causil S. The relationship between the normalized difference vegetation index, rainfall, and potential evapotranspiration in a banana plantation of Venezuela. SAINS TANAH - Journal of Soil Science and Agroclimatology. 2021;18(1):58-64
 17. Hernández R, Olivares B, Arias A, Molina JC, Pereira Y. Identification of potential agroclimatic zones for production of onion (*Allium cepa* L.) in Carabobo, Venezuela. Journal of the Selva Andina Biosphere. 2018;6(2):70-82.
 18. Hernandez R, Olivares B, Arias A, Molina JC, Pereira Y. Eco-territorial adaptability of tomato crops for sustainable agricultural production in Carabobo, Venezuela. Idesia. 2020;38(2):95-102.
 19. Olivares B, Hernández R. Ecoterritorial sectorization for the sustainable agricultural production of potato (*Solanum tuberosum* L.) in Carabobo, Venezuela. Agricultural Science and Technology. 2019;20(2):339-354.
 20. Olivares, B., Hernández, R. Application of multivariate techniques in the agricultural land's aptitude in Carabobo, Venezuela. Tropical and Subtropical Agroecosystems, 2020. 23(2):1-12.
 21. Olivares B, Lobo D, Cortez A, Rodríguez MF, Rey JC. Socio-economic characteristics and methods of agricultural production of indigenous community Kashaama, Anzoátegui, Venezuela. Rev. Fac. Agron. (LUZ). 2017;34(2):187-215.
 22. Zingaretti ML, Olivares B, Demey Zambrano JA, Demey JR. Typification of agricultural production systems and the perception of climate variability in Anzoátegui, Venezuela. Revista FAVE - Ciencias Agrarias. 2016;15(2):39-50.
 23. Olivares B. Banana production in Venezuela: Novel Solutions to Productivity and Plant Health. Springer Nature. 2023;176.

© 2023 Ragul et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/103827>