



***In silico* Analysis of *SINAC1* Transcription Factor, a Positive Modulator for Tomato Leaf Curl Virus**

**Nimbal Shivraj Ramchandra ^a, Williams M. ^a,
Debasmita Behera ^a, Sakthi, A. R. ^a, Veera Ranjani ^a,
Saraswathi T. ^b, Varanavasiappan S. ^a, Raveendran M. ^c
and Sudha M. ^{a*}**

^a Department of Plant Biotechnology, Centre for Plant Molecular Biology and Biotechnology, TNAU, Coimbatore, Tamil Nadu, India.

^b Department of Medicinal and Aromatic Crops, HC & RI, TNAU, Coimbatore, Tamil Nadu, India.

^c Director of Research, Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i203819

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/107016>

Original Research Article

Received: 17/07/2023

Accepted: 21/09/2023

Published: 22/09/2023

ABSTRACT

NAC proteins, part of the largest plant transcription factor family, play pivotal roles in plant development, and biotic and abiotic stresses. Among them, the *SINAC1* protein in tomato serves as a Negative regulator, enhancing tolerance to cold, bacterial, and viral diseases. However, our knowledge of *SINAC1* protein *in silico* analysis, evolutionary relationships, and protein interactions remains limited. We used *in silico* tools to study the *SINAC1* protein in tomato. Phylogenetic analysis showed it is closely related to other *Solanaceae* family members. Motif analysis revealed seven conserved motifs shared among *Solanaceae* members. The presence of NLS confirms its abundance in the nucleus. Protein-protein interaction analysis predicted ten potential nuclear partners, including five uncharacterized proteins and others involved in plant development.

*Corresponding author: E-mail: sudhatamil@gmail.com;

In silico expression, analysis showed abundant SINAC1 presence in both fruit and roots. This study provides the role of SINAC1 protein in tomato, shedding light on its evolutionary context and protein interactions.

Keywords: SINAC1; tomato leaf curl virus; phylogenetic analysis; *In silico* expression analysis.

1. INTRODUCTION

Transcription factors (TFs) are proteins that control gene expression by binding to specific regulatory elements in gene promoters, either by activating or repressing transcription [1]. In plants, this regulation is essential for processes like cellular development, signal transduction, and responses to environmental stress and pathogens [2,3]. Identifying and understanding the relationship between TF and target genes is crucial for unraveling the intricate transcription network in plants [4]. Among the 58 TF families, six Transcription factor families play a prominent role in stress responses. Among these, NAC TFs actively participate in the response to biotic and abiotic stresses. A limited number of cases have provided a comprehensive account of the molecular mechanisms underlying these processes [5].

NAC proteins constitute the largest protein family and are distinguished by their distinctive structures, featuring a highly conserved N-terminal NAC domain and a notably diverse C-terminal region. The NAC domain was initially identified in four proteins: NAM, ATAF1, ATAF2, and CUC2 [6]. In contrast, the C-terminal domain of NAC proteins exhibits significant variations in both sequence and length, having a pivotal role in specifying the precise functions and activities of NAC proteins in the species [7].

NAC proteins are majorly involved in stress conditions because they influence the plant's response to stress. The alteration of NAC gene expression through overexpression or silencing leads to heightened or diminished resistance against pathogens [8,9], and Overexpression of ANAC019, ANAC055, ANAC72 in *Arabidopsis* [10], SNAC2 [11], and OsNAC4 [12] in *O. Sativa* leads to drought and salt tolerance. Loss of function in ataf1-1 [13] and rim1-1 [14] leads to attenuated penetration resistance toward Bgh attack in *A. thaliana* and resistance to rice dwarf virus (RDV) in *O. sativa*, respectively. overexpression of HbNAC1, PtrNAC72 leads to the drought tolerance in a *h. brasiliensis* and *Arabidopsis* [15,16] and overexpression of HvSNAC1, and HvNAC6 leads to the resistance

against *barley ramularia leaf spot* (RLS) disease and powdery mildew pathogen *Blumeria graminis f. sp. hordei* (Bgh) respectively in barley [17,18] and the knockdown of the TaNAC1 and enhances the resistance against Puccinia stripe rust in wheat [19]. This indicates that the NAC Transcription factor especially the NAC1 Transcription factor may exert both positive and negative regulatory effects on plant defence responses in biotic as well as abiotic stresses [20] The significant reduction in tomato production due to Tomato Leaf Curl Virus (TYLCV) necessitates the development of resistant tomato varieties, achievable through the application of genome editing tools by targeting recessive gene [21]

In tomato, SINAC1 gene expression increases exponentially by the replication enhancer (RE) of the *Tomato Leaf Curl Virus* (TYLCV), which helps the virus to accumulate more in the plant and is also enhanced by *Pseudomonas* infection, indicating the role of the SINAC1 protein as a positive modulator of TYLCV and *Pseudomonas* infection in plants [8,22]. Knocking down this gene resulted in Cold and Drought tolerance and increased fruit shelf life in the tomato [23,24]. This TF upon knocking down might harbour multi-stress tolerance in the plant, but unfortunately, *in silico* analysis was little explored.

In the present study, to understand the structural, functional, and evolutionary relationship of the SINAC1 protein in tomato, we conducted an *in silico* analysis of the SINAC1 protein in tomato in comparison with seven different model plants. This analysis will provide a reference point for the SINAC1 gene in tomato, as we have studied its interaction with other proteins and its molecular aspects.

2. MATERIALS AND METHODS

2.1 Sequence Retrieval, Chromosomal Location Identification, MSA, and Phylogenetic Analysis of NAC1 Transcription Factors

The NAC1 transcription factor sequence from *Solanum lycopersicum* was obtained from the

Plant Transcription Factor Database (PlantTFDB v5.0) (<http://planttfdb.gao-lab.org>) [25]. In addition, sequences from various model plants, such as *Arabidopsis thaliana* and *Oryza sativa* subsp. *Japonica*, *Solanum melongena*, *Zea mays*, *Capsicum annuum*, *Glycine max*, and *Sorghum bicolor*, were retrieved from the National Center for Biotechnology Information (NCBI) (<https://www.ncbi.nlm.nih.gov>) and UniProt databases (<https://www.uniprot.org>) [26]. Once these sequences were obtained, they were imported into InterProScan (<https://www.ebi.ac.uk/interpro/search/sequence/>) [27] to check for the presence of the NAC Domain.

Following the sequence analysis, a multiple sequence alignment was carried out using the Clustal Omega online software [<https://www.ebi.ac.uk/Tools/msa/clustal/>] [28]. Subsequently, a phylogenetic tree was constructed using the Maximum Likelihood algorithm in the MEGA 11 software [29]. The resulting data were visualized and represented in a more informative graphical format using the Interactive Tree Of Life (iTOL) online tool (<https://itol.embl.de/>) [30].

2.2 Possible Motif Prediction and Physiochemical Analysis and Protein-protein Interaction Study of NAC1 Transcription Factor

Several essential analyses were conducted to study the NAC1 Transcription factor. Initially, tools such as TBtool (<https://github.com/CJ-Chen/TBtools>) [31] and MG2C (http://mg2c.iask.in/mg2c_v2.1/) [32] were used to precisely locate the NAC1 gene within the tomato genome. Subsequently, motif prediction was carried out using the MEME online tool (<https://meme-suite.org/meme/tools/meme>) [33], with a focus on sequences from different model plants. The maximum and minimum motif widths were 50 and 6, respectively, while limiting the maximum number of motifs to 10. The physiochemical properties of the SINAC1 protein were analyzed. Parameters such as the theoretical isoelectric point (pI), molecular weight, aliphatic index, and Grand Average of hydropathicity (GRAVY) were calculated using the ExPASy Prot-Param tool (<https://web.expasy.org/protparam/>) [33,34], with the UniProt sequence as the reference. Finally, the protein-protein interaction network of the SINAC1 protein was explored using the STRING online software tool (<https://string-db.org/>), shedding light on potential functional

associations and interactions with other proteins.

2.3 Secondary Structure Prediction, Subcellular Location Identification, and In silico Expression Analysis of SINAC1 Transcription Factor Protein

The SOPMA server [35] was used for the prediction of secondary structures of SINAC1 proteins with the output width set to 70, the number of conformational states to 4, similarity threshold to 8, and window width to 17. The subcellular location of the SINAC1 Transcription factor protein was identified by using the TargetP2.0 online tool (<https://services.healthtech.dtu.dk/services/TargetP-2.0/>) [36], this tool particularly predicts the presence of N-terminal sequences: signal peptide (SP), mitochondrial transit peptide (mTP), chloroplast transit peptide (cTP) or thylakoid luminal transit peptide (ITP) this helps to determine the subcellular location. The tomato plant Electronic Fluorescent Pictograph Browsers (Tomato eFP browsers) (http://bar.utoronto.ca/eplant_tomato/) were used for the *In silico* expression analysis of the SINAC1 protein

3. RESULTS AND DISCUSSION

3.1 Identification of Evolutionary Relationships

In this study, putative NAC1 transcription factor (NAC1) genes were identified in the *Solanum lycopersicum* genome using *in silico* tools, revealing one gene, SINAC1 (Solyc04g009440.2.1), whose protein product contains the NAC conserved domain of 127 amino acids and its chromosomal location is represented in Fig. 1a.

To explore the expansion of NAC1 genes across different plant species, 18 full-length amino acid sequences from seven model plants (*Arabidopsis thaliana*, *Zea mays*, *Oryza sativa* subsp. *Japonica*, *Sorghum bicolor*, *Capsicum annuum*, *Solanum melongena*, and *Glycine max*) were used to construct a phylogenetic tree using the maximum likelihood method (Fig. 1b). A study conducted by Bhattacharjee et al. utilized *in silico* tools to identify a total of 99 NAC transcription factors (NAC TFs) in the *Solanum lycopersicum* genome. These NAC TFs were found to be distributed across various regions of the genome

[37]. Across various species, different numbers of *NAC1* genes were found, including three in *Arabidopsis thaliana* and five in *Zea mays*. These results indicated that these transcription factors could be classified into three primary groups. One group consisted of *Arabidopsis* proteins with distinct motifs, whereas all *Solanaceae* plants fell into another main group. Additionally, all cereals and pulse crops were grouped together, sharing similar motifs with slight variations that led to the formation of smaller subgroups. The research conducted by Zhu et al. explored the evolutionary relationships among NAC transcription factors across nine major land plant lineages, revealing a high degree of conservation in their NAC domains across species, and proposed the existence of 21 subfamilies [38]. Furthermore, a study by Cenci et al. performed a phylogenetic analysis of NAC domain sequences from *A. thaliana*, *V. vinifera*, *O. sativa*, and *M. acuminata*. Their analysis identified 26 clusters that encompassed all NAC members of orthologous groups, as determined by expert comparison [39]. Bengoa Luoni et al. reported that the C-terminal region of NAC transcription factors is highly variable and species-specific, with this variability playing a critical role in stress responses [40]. Phylogenetic analysis demonstrated that *Solanaceae* plants share a common evolutionary history, making them somewhat distinct from other plant classes. However, they share some common motifs that connect them.

3.2 Physiochemical and Motif Analysis

The Physiochemical properties of *SINAC1* were thoroughly examined using the ExPASy Prot-Param tool, and the resulting data are presented in Table 1. The *SINAC1* Transcription factor protein sequence contained 39 and 38 positive and negative amino acids, respectively. The isoelectric point was 7.6. It has been observed that the isoelectric point value of any protein has a significant impact on its subcellular localization and stability [41]. The research conducted by Nasrabadi et al. (2021) demonstrated that altering the physiochemical characteristics of plant proteins can influence both their technological functionality and biological properties [42]. Given that the *SINAC1* protein functions as a suppressor, it is crucial to evaluate the consequences of modifying physiochemical properties in genome-edited crops when this factor is knocked down. There is the total of 10 different motifs were identified (Fig. 2), and the first three motifs are in Table 2. and belongs to the highly conserved NAC domain. *Solanum Lycopersicum*, *capsicum annum*, and *solanum melongena*, three share a close association and collectively possess 7 motifs, these can be easily identified by the phylogenetic tree. A similar study conducted by Du et al. in *Kandelia obovate* revealed the total of 20 conserved motifs in the *KoNAC* protein. Additionally, through a comprehensive phylogenetic analysis, they were able to categorize the 79 *KoNAC* proteins into 16 distinct subgroups [43].




Fig. 1. a) Chromosomal location of *NAC1* in *S. Lycopersicum* b) Phylogenetic map of *SINAC1* Transcription factor in different model plants

By examining protein motifs and tracing their evolutionary relationships, valuable insights can be obtained. It is evident that *SINAC1* shares its evolutionary history with members of the *Solanaceae* family and shares common conserved regions with other crop families. There is a total of four motifs (Table 2) which were conserved in all the sequences of the model plant. Of these four motifs, the first three motifs are part of the NAC domain. With the exception of the *Solanaceae* family, motifs 8 and 10 were lost in the other studied species. This information can serve as a starting point to delve into the mysterious role of the *NAC1* transcription factor in various crops. In some crops, it may confer resistance, whereas in others, it may act as a susceptibility factor [44]. This divergence could be influenced by specific motifs with distinct plant-specific roles that require closer examination.

3.3 Secondary Structure Prediction, Subcellular Location Identification, and *In silico* Expression Analysis

The process involved utilizing the SOPMA tool with a retrieved sequence as input to predict the secondary structure, which is presented in Table 3. Subsequently, subcellular localization of the Transcription Factor was determined, predominantly in the nucleus, without the

presence of mitochondrial, thylakoid lumen, or chloroplast transfer peptides in the protein sequence. Transcription factors play a crucial role in the nucleus by facilitating transcription processes and other essential biological functions. A study reported that the NAC domain typically contains the Nuclear Localization Signal (NLS) and rarely other extra-nuclear signal peptides [45]; however, in the case of the *SINAC1* protein, it is missing, which indicates its limited role up to the nucleus. To confirm the *in silico* subcellular localization predictions, GFP tags were fused with the KoNAC proteins, and subsequent examination was conducted using confocal microscopy. The fluorescent signals obtained delineated the subcellular localization of the NAC proteins within the nucleus [43]. Additionally, the results obtained from the Tomato eFP browsers indicated significant *SINAC1* gene activity across various tissues, with the highest expression observed in fruits, followed by roots, vegetative meristem, developing fruit, leaves, stems, and unopened flowers, as illustrated in Fig. 3. An *in vivo* expression study of NAC1 TF found that it was highly expressed in the root and ripened fruits of crops such as tobacco and wheat [20]. This analysis provides valuable insights into optimal plant tissues for studying the expression of this specific protein, which can be particularly useful for investigating protein product mutants.

Table 1. Physiochemical properties of *SINAC1* protein

Properties	<i>SINAC1</i> Protein
Amino acids	301
MW(DA)	34775.2
Positively charged amino acids	39
Negatively charged amino acids	38
Theoretical Pi	7.6
Extinct coefficient	54235
Instability index	38.46
Aliphatic index	58.7
GRAVY	-0.824

Table 2. Conserved motifs discovered in *SINAC1*

Sr. No.	Motifs	Width
1	VDLYKFDPWDLPERALFGEKEWYFFSPRDRKYPNGSRPNRAAGNGYWKAT	50
2	RGRTLGIKKALVFYAGKAPRGVKTWDWIMHEYR	32
3	ELNLPPGFRFHPTDDELVEHYLCRKAAGQ	29
4	RLDDWVLCRLYNKKNEWKMQ	21
5	DMATSHSHSHSQSHSHSWGETRTPESIVDNDPPELDSFPAFQDPAMMM	50
6	DGCSAKSSBLFVDLSYDDIQGMYSGLDMLPP	31
7	GEDFYSSLFASPRVKGNQPA	20
8	DFQHNYLDGLLNPPFETQMQQQNCNIDQFNSFQDMFJPMQK	41
9	JVPPIAE	8
10	TRMHTTNSSSGSEHVLSPCDKEVQSAPKW	29



Fig. 2. Motif identification in the seven-model plant NAC1 Transcription factor and NAC1 Transcription factor of *S. Lycopersicum*

Table 3. Secondary structure prediction of *SINAC1* transcription factor

Secondary structure element	Percentage
Alpha helix (%)	19.60%
Beta turn (%)	2.99%
Extended strand (%)	9.63%
Random coil (%)	67.77%

3.4 In silico Study of Protein-protein Interactions of the *SINAC1* Transcription Factor

The protein sequence was input into the String software, which then identified the sequence with the highest similarity, *A0A3Q7FYL7*, exhibiting a remarkable 94.7% identity, a bit score of 474.6, and an e-value of 5.2e-134. This analysis generated a network graph (Fig. 4) with 11 nodes and 48 edges. Predictions also identified 10 functional partners, of which five were proteins with uncharacterized functions, while the rest are detailed in Table 4. *ZmABI19*, a protein

containing the transcription factor-B3 domain found in *Zea mays*, plays a direct role in governing essential factors that coordinate endosperm and embryo development and grain filling in maize [46]. Auxin response factor protein is a transcription factor that plays an active role in plant growth and development by regulating the expression of auxin response genes. An interaction study conducted for the *CaNAC* genes in pepper, interaction network revealed its interaction with the 48 genes which have a significant role in biotic, abiotic stress tolerance and the growth and development in plants [47]. Similarly, in the case of the *SINAC47*

transcription factor, a similar interaction study unveiled its interactions with proteins that are involved in hormonal responses [48]. This significantly simplifies our ability to anticipate the outcomes of genome editing

experiments aimed at knocking down target protein molecules. As the CRISPR-Cas9 system has OFF-Target as a limitation, it is very important to predict which proteins might be affected.

Plant eFP: Solyc04g009440




Fig. 3. *In silico* expression characterization of tomato *SINAC1* transcription factor

Table 4. List of Functional Partners of *SINAC1* Transcription Factor

Primary accession	Name
A0A3Q7J310	TF-B3 domain-containing protein.
ARF6	Auxin_resp domain-containing protein.
A0A3Q7H9Z3	Auxin response factor
A0A3Q7I5Z7	PHD_Oberon domain-containing protein.
A0A3Q7EAX8	Phytoeyanin domain-containing protein.




Fig. 4. Network graph

4. CONCLUSION

The present study unveiled the evolutionary relationship between our target gene and members of the *Solanaceae* group, as evidenced by the sharing of all seven predicted motifs. Additionally, through physicochemical analysis and *in silico* expression profiling, we explained the chemical properties of this gene and identified potential target sites for further qualitative and quantitative protein studies. Furthermore, our investigation into protein-protein interactions revealed crucial associations with proteins that play pivotal roles in plant developmental processes. These interactions hold the promise of providing insights into the potential consequences of knocking down this gene in plants using the CRISPR-Cas9 technology.

ACKNOWLEDGEMENTS

The authors are thankful to the Centre for Plant Molecular Biology and Biotechnology, Tamil Nadu Agricultural University for providing access to conducting *the in silico* analyses.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Schwechheimer C, Zourelidou M, Bevan M. Plant transcription factor studies. Annual Review of Plant Biology. 1998;49(1):127-50.
- Riechmann JL, Ratcliffe OJ. A genomic perspective on plant transcription factors. Current Opinion in Plant Biology. 2000;3(5):423-34.
- Zhang C-Q, Xu Y, Lu Y, Yu H-X, Gu M-H, Liu Q-Q. The WRKY transcription factor OsWRKY78 regulates stem elongation and seed development in rice. Planta. 2011;234:541-54.
- Shang H, Li W, Zou C, Yuan Y. Analyses of the NAC transcription factor gene family in *Gossypium raimondii* Ulbr.: Chromosomal location, structure, phylogeny, and expression patterns. Journal of Integrative Plant Biology. 2013;55(7):663-76.
- Meraj TA, Fu J, Raza MA, Zhu C, Shen Q, Xu D, et al. Transcriptional factors regulate plant stress responses through mediating secondary metabolism. Genes. 2020;11(4):346.
- Olsen AN, Ernst HA, Leggio LL, Skriver K. NAC transcription factors: Structurally distinct, functionally diverse. Trends in plant science. 2005;10(2):79-87.
- Ooka H, Satoh K, Doi K, Nagata T, Otomo Y, Murakami K, et al. Comprehensive analysis of NAC family genes in *Oryza sativa* and *Arabidopsis thaliana*. DNA research. 2003;10(6):239-47.
- Huang W, Miao M, Kud J, Niu X, Ouyang B, Zhang J, et al. SI NAC 1, a stress-related transcription factor, is fine-tuned on both the transcriptional and the post-translational level. New Phytologist. 2013;197(4):1214-24.
- Wang Xe, Basnayake BVS, Zhang H, Li G, Li W, Virk N, et al. The Arabidopsis ATAF1, a NAC transcription factor, is a negative regulator of defense responses against necrotrophic fungal and bacterial pathogens. Molecular Plant-Microbe Interactions. 2009;22(10):1227-38.
- Tran L-SP, Nakashima K, Sakuma Y, Simpson SD, Fujita Y, Maruyama K, et al. Isolation and functional analysis of Arabidopsis stress-inducible NAC transcription factors that bind to a drought-responsive cis-element in the early responsive to dehydration stress 1 promoter. The Plant Cell. 2004;16(9):2481-98.
- Sindhu A, Chintamanani S, Brandt AS, Zanis M, Scofield SR, Johal GS. A guardian of grasses: specific origin and conservation of a unique disease-resistance gene in the grass lineage. Proceedings of the National Academy of Sciences. 2008;105(5):1762-7.
- Zheng X, Chen B, Lu G, Han B. Overexpression of a NAC transcription factor enhances rice drought and salt tolerance. Biochemical and biophysical research communications. 2009;379(4):985-9.
- Jensen MK, Rung JH, Gregersen PL, Gjetting T, Fuglsang AT, Hansen M, et al. The Hv NAC6 transcription factor: A positive regulator of penetration resistance in barley and Arabidopsis. Plant molecular biology. 2007;65:137-50.
- Yoshii M, Shimizu T, Yamazaki M, Higashi T, Miyao A, Hirochika H, et al. Disruption of a novel gene for a NAC-domain protein in rice confers resistance to Rice dwarf virus. The Plant Journal. 2009;57(4):615-25.

15. Cao Y, Zhai J, Wang Q, Yuan H, Huang X. Function of *Hevea brasiliensis* NAC1 in dehydration-induced laticifer differentiation and latex biosynthesis. *Planta*. 2017;245:31-44.
16. Wu H, Fu B, Sun P, Xiao C, Liu J-H. A NAC transcription factor represses putrescine biosynthesis and affects drought tolerance. *Plant Physiology*. 2016;172(3):1532-47.
17. McGrann GR, Steed A, Burt C, Goddard R, Lachaux C, Bansal A, et al. Contribution of the drought tolerance-related Stress-responsive NAC 1 transcription factor to resistance of barley to *Ramularia leaf spot*. *Molecular Plant Pathology*. 2015;16(2):201-9.
18. Chen Y-J, Perera V, Christiansen MW, Holme IB, Gregersen PL, Grant MR, et al. The barley HvNAC6 transcription factor affects ABA accumulation and promotes basal resistance against powdery mildew. *Plant molecular biology*. 2013;83:577-90.
19. Wang F, Lin R, Feng J, Chen W, Qiu D, Xu S. TaNAC1 acts as a negative regulator of stripe rust resistance in wheat, enhances susceptibility to *Pseudomonas syringae*, and promotes lateral root development in transgenic *Arabidopsis thaliana*. *Frontiers in plant science*. 2015;6:108.
20. Bian Z, Gao H, Wang C. NAC transcription factors as positive or negative regulators during ongoing battle between pathogens and our food crops. *International Journal of Molecular Sciences*. 2020;22(1):81.
21. Pramanik D, Shelake RM, Park J, Kim MJ, Hwang I, Park Y, et al. CRISPR/Cas9-mediated generation of pathogen-resistant tomato against tomato yellow leaf curl virus and powdery mildew. *International journal of molecular sciences*. 2021;22(4):1878.
22. Selth LA, Dogra SC, Rasheed MS, Healy H, Randles JW, Rezaian MA. A NAC domain protein interacts with tomato leaf curl virus replication accessory protein and enhances viral replication. *The Plant Cell*. 2005;17(1):311-25.
23. Niu X, Lu H, Fan Y, Wang W, Yuan Y, Hawkins M, et al. Manipulation of the transcription factor SINAC1 for improved tolerance to abiotic stress in tomato. *Plant, Cell & Environment*. 2022;45(12):3537-50.
24. Meng C, Yang D, Ma X, Zhao W, Liang X, Ma N, et al. Suppression of tomato SINAC1 transcription factor delays fruit ripening. *Journal of Plant Physiology*. 2016;193:88-96.
25. Tian F, Yang D-C, Meng Y-Q, Jin J, Gao G. Plant Reg Map: Charting functional regulatory maps in plants. *Nucleic Acids Research*. 2019;48(D1):D1104-D13.
26. Consortium TU. UniProt: the Universal Protein Knowledgebase in 2023. *Nucleic Acids Research*. 2023;51(D1):D523-D31.
27. Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, et al. InterProScan 5: genome-scale protein function classification. *Bioinformatics*. 2014;30(9):1236-40.
28. Madeira F, Pearce M, Tivey ARN, Basutkar P, Lee J, Edbali O, et al. Search and sequence analysis tools services from EMBL-EBI in 2022. *Nucleic Acids Research*. 2022;50(W1):W276-W9.
29. Tamura K, Stecher G, Kumar S. MEGA11: molecular evolutionary genetics analysis version 11. *Molecular Biology and Evolution*. 2021;38(7):3022-7.
30. Letunic I, Bork P. Interactive Tree of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation. *Nucleic Acids Research*. 2021;49(W1):W293-W6.
31. Chen C, Chen H, Zhang Y, Thomas HR, Frank MH, He Y, et al. TBtools: An integrative toolkit developed for interactive analyses of big biological data. *Molecular Plant*. 2020;13(8):1194-202.
32. Chao J, Li Z, Sun Y, Aluko OO, Wu X, Wang Q, et al. MG2C: A user-friendly online tool for drawing genetic maps. *Molecular Horticulture*. 2021;1(1):1-4.
33. Bailey TL, Johnson J, Grant CE, Noble WS. The MEME Suite. *Nucleic Acids Research*. 2015;43(W1):W39-W49.
34. Gasteiger E, Hoogland C, Gattiker A, Duvaud Se, Wilkins MR, Appel RD, et al. Protein identification and analysis tools on the ExpASY Server: Springer; 2005.
35. Geourjon C, Deleage G. SOPMA: Significant improvements in protein secondary structure prediction by consensus prediction from multiple alignments. *Bioinformatics*. 1995;11(6):681-4.
36. Emanuelsson O, Nielsen H, Brunak S, Von Heijne G. Predicting subcellular localization of proteins based on their N-terminal amino acid sequence. *Journal of Molecular Biology*. 2000;300(4):1005-16.
37. Bhattacharjee P, Das R, Mandal A, Kundu P. Functional characterization of tomato membrane-bound NAC transcription factors. *Plant molecular biology*. 2017;93:511-32.

38. Zhu T, Nevo E, Sun D, Peng J. Phylogenetic analyses unravel the evolutionary history of nac proteins in plants. *Evolution*. 2012;66(6):1833-48.
39. Cenci A, Guignon V, Roux N, Rouard M. Genomic analysis of NAC transcription factors in banana (*Musa acuminata*) and definition of NAC orthologous groups for monocots and dicots. *Plant molecular biology*. 2014;85(1-2):63-80.
40. Bengoa Luoni SA, Cenci A, Moschen S, Nicosia S, Radonic LM, Sabio y García JV, et al. Genome-wide and comparative phylogenetic analysis of senescence-associated NAC transcription factors in sunflower (*Helianthus annuus*). *BMC Genomics*. 2021;22(1):893.
41. Tokmakov AA, Kurotani A, Sato K-I. Protein pi and intracellular localization. *Frontiers in Molecular Biosciences*. 2021;8:775736.
42. Nasrabadi MN, Doost AS, Mezzenga R. Modification approaches of plant-based proteins to improve their techno-functionality and use in food products. *Food Hydrocolloids*. 2021;118:106789.
43. Du Z, You S, Yang D, Tao Y, Zhu Y, Sun W, et al. Comprehensive analysis of the NAC transcription factor gene family in *Kandelia obovata* reveals potential members related to chilling tolerance. *Frontiers in Plant Science*. 2022;13:1048822.
44. Nuruzzaman M, Sharoni AM, Kikuchi S. Roles of NAC transcription factors in the regulation of biotic and abiotic stress responses in plants. *Frontiers in microbiology*. 2013;4:248.
45. Mohanta TK, Yadav D, Khan A, Hashem A, Tabassum B, Khan AL, et al. Genomics, molecular and evolutionary perspective of NAC transcription factors. *PloS one*. 2020;15(4):e0231425.
46. Yang T, Guo L, Ji C, Wang H, Wang J, Zheng X, et al. The B3 domain-containing transcription factor ZmABI19 coordinates expression of key factors required for maize seed development and grain filling. *The Plant Cell*. 2020;33(1):104-28.
47. Diao W, Snyder JC, Wang S, Liu J, Pan B, Guo G, et al. Genome-wide analyses of the NAC transcription factor gene family in pepper (*Capsicum annuum* L.): Chromosome location, phylogeny, structure, expression patterns, cis-elements in the promoter, and interaction network. *International Journal of Molecular Sciences*. 2018;19(4):1028.
48. Huang Y, Li T, Xu Z-S, Wang F, Xiong A-S. Six NAC transcription factors involved in response to TYLCV infection in resistant and susceptible tomato cultivars. *Plant Physiology and Biochemistry*. 2017;120:61-74.

© 2023 Ramchandra et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/107016>