



Effect of Improved Hive Cover Designs on Internal Microclimate and Colony Establishment of West African Honeybees (*Apis mellifera adansonii* L.) in Awka, Nigeria

B. U. Ononye ^{a*}, C. E. Akunne ^a, E. O. Ogbuefi ^b, T. E. Okeke ^c,
K. P. Okafor ^a, E. I. Azaka ^d, G. E. Obiyo ^d,
C. O. Aniefuna ^b, P. C. Akwuaka ^d and C. A. Chidi ^a

^a Department of Zoology, Honeybee Research Centre, Nnamdi Azikiwe University, Awka, Nigeria.

^b Department of Parasitology and Entomology, Nnamdi Azikiwe University, Awka, Nigeria.

^c Department of Biology, Alex Ekwueme, Federal University, Ndufu-Aliko Ikwo, Ebonyi, Nigeria.

^d Department of Zoology, Nnamdi Azikiwe University, Awka, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JALSI/2023/v26i5614

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/104464>

Original Research Article

Received: 05/06/2023

Accepted: 08/08/2023

Published: 22/08/2023

ABSTRACT

Hive cover designs majorly affects the microclimate and colony establishment of African honeybee colonies in Nigeria. It became pertinent therefore to assess the effect of modified hive cover designs on the internal microclimate and colony establishment of West African honeybee (*Apis mellifera adansonii* L.) Colonies in Awka, Nigeria. Four insulating materials were used to construct

*Corresponding author: E-mail: bu.ononye@unizik.edu.ng;

the covers of twelve modified frame bar hives and designated: Control (plywood); T1 (warped boards); T2 (PVC) and T3 (foam). Each hive treatment was replicated three times. Data on ambient and hive microclimate as well as the colonization of honeybees' colonies were observed and reported. The results indicated that the control had the highest temperature (31.86 °C) while T1 (30.40°C) had the least. The hive temperature was higher in March, 2023 (32.72°C) while August (29.87°C) had the least. The highest relative humidity (73.18%) was recorded in T1 while the control had the least (64.38%). The hive temperature and relative humidity was significantly affected by hive cover designs and different months of the study period ($P < 0.05$). The highest colonization by African honeybees was observed in the control (100%) but none occurred in T2 (0.00%). Colonization of the Control hives occurred in July, October, and November, 2022 while that of T1 and T3 occurred in December, 2022. Colony establishment of African honeybees was not significantly affected by hive cover designs ($p = 0.26$) and different months of the study period ($p = 0.24$). It was concluded that beekeepers should adopt the insulation of hives covers with plywood wood since it had 100% colonization of West African honeybees'.

Keywords: *Hive cover designs; internal microclimate; colony establishment; African honeybees.*

1. INTRODUCTION

Thermoregulation is one of the most remarkable characteristics of the honey bee, particularly the *Apis mellifera* colony, which can maintain a temperature range of 33-36°C while the ambient temperature falls between -40 and 40°C [1]. Honeybee colonies exposed to high external temperatures may experience a rise in hive temperature, which can have a negative impact on honeybee growth, metabolism, and physiological responses [2-4]. More so, temperature and relative humidity have a significant impact on the honeybee colony's homeostasis in the hive [5]. The level of honeybee activity in the hives and its surroundings is highly influenced by ambient temperature and relative humidity, which, if consistently observed, might be used to determine the harvest season [6,7]. Previous research revealed that maintaining the optimal hive temperature range (33 to 36°C) is crucial to beekeeping due to the vulnerability of juvenile bee development cycles to high temperatures [8]. When the ambient temperature is excessively hot or cold, honeybees maintain a temperature of roughly 32°C of the hive by fanning their wings to either circulate air and cool the hive or create friction and warm the internal air [9]. However, Park et al. [10] found that, despite the shift in ambient temperature, the internal temperature remained above 31°C. According to Abd-Elmawgood et al. [11], honeybees can survive in temperatures ranging from -20 to +60 degrees Celsius, however they perform best in temperatures ranging from +21 to +35 degrees Celsius.

Maintaining a constant temperature and relative humidity is critical for bee colony activities, this is

why some researchers have redesigned hives in the past. For instance, the modifications of hive entrances was previously reported to significantly affect the temperature and relative humidity variations of hives [4,12]. Tuo et al. [13] used pan traps (UV- brilliant blue, yellow and white) to analyse the impact of different agroecosystems on bee activity in hives and found that bee activity evolves with temperature (34°C and 35°C) up to a certain point and inversely with relative humidity (56% and 65%). Furthermore, pollination in natural and agricultural systems is affected by bees' responses to changing environmental temperatures, with rising average temperatures resulting to pollinator population declines [14]. Studies by Lepkova et al. [15] revealed that the microclimate in hives composed of various materials was substantially altered, however ceramic hives with high kaolin content had better thermal insulation capabilities than timber hives. Some authors earlier reported that foam polystyrene is one of the most recent alternatives utilised for hive design, and has proven to be harmless to organisms when used under normal settings, thus it is recommended for future bee breeding because its qualities are superior to those of wood [16-19]. Furthermore, hives made of foam polystyrene have strong thermal insulation capabilities, preventing the hive from being heated or cooled from the outside [16]. St. Clair et al. [20] recently reported that the new insulating materials with superior qualities to wood could provide better choices for the beekeeping industry and it was reported that polyurethane hives maintained a much higher overall temperature than wooden hives, and that polyurethane hives had significantly higher ideal relative humidity (52.05%) compared to wooden hives (62.50%). Furthermore, they discovered that the hive temperature patterns of the wooden

hives fluctuated significantly more than those of the polyurethane hives. Another study by Meikle et al. evaluated the effects of hive entrance direction on honey bee colony activity and temperature and concluded that hive orientation should be considered in the design of field research involving colony activity monitoring.

Developing strategies for keeping a stable brood hive temperature despite the external environment has allowed western honey bees to colonise much of the globe and are indeed successful organisms [1]. Akinbi et al. [21] studied the performance of wood hive and plywood hive types in three different vegetation types and noted an improvement in the colonisation performance at different months of the study period after installation of hives. They also reported that plywood hives installed in the cocoa farmland and plantation accounted for higher colonisation percentage than natural forest. Studies by Bykov and Zaitsev [16] revealed that honeybee colonies in wooden hives colonised at a rate of 91.67%, with slower development due to the hive's periodic compression throughout the cold season. Dupleix et al. [22] perceived that the reasons wooden habitat supports bees' well-being include heat insulation, humidity regulation, "naturalness," and fragrance. Adedeji et al. [23] reported that season has a significant impact on hive colonisation in Nigeria's South-South region, and it has also been stated that the best period to build hives is between August and September, because colonisation usually occurs in October. Different times for colonization of honeybees have been reported ranging from February to April [23] and October to February [23]. Anumba et al. [24] after the evaluation of the influence of hive colours on the performance of African honeybee colonies in Awka reported that hive colour did not significantly enhance colonisation. Studies by Ononye and Akunne, [12] on the impact of time and beehive entrance on colony development of African honeybees indicated that the colonisation months in the research area were August to October.

Other independent studies [25,26] attributed variation in the rate of colonisation of hive types to a number of environmental and anthropogenic

factors while the high colonisation was attributed to differences in vegetation cover. However, delay in hive colonisation of honeybees was attributed to unfavorable microclimatic during rainy season under a natural forest ecosystem [21]. Studies on the effects of temperature and relative humidity on colonisation of honey bee hives within FUTA community by Oluwaseyi et al. [27] showed that hive located where the environmental conditions were favorable was observed to colonised quickly when compared to other locations. However, earlier studies by Ononye and Akunne [28] on the effect of modified hive entrances on colony establishment found that hive entrances had no effect on African honeybee colonisation during the study period. Poor hive and seasonal management were cited as some of the causes responsible for the annual reduction in honey bee colony establishment. The paucity of information on the influence of microclimate and hive cover designs on the performance of West African honeybee colonies in Nigeria necessitated this study which aimed to improve on the existing hive cover designs using various insulating materials such as plywood, warped wood, polyvinyl chloride, PVC ceilings and foam.

2. MATERIALS AND METHODS

2.1 Study Area and Duration

This study was carried out at the Honeybee Research Centre (Fig. 1) of the Department of Zoology, Nnamdi Azikiwe University, Awka from January, 2022 to March, 2023. Awka lies within geographical coordinates of 6°14'53" N and 7°05'34" E. Awka is located in Nigeria's tropical rain forest zone and has two distinct seasons: wet and dry, which occur between April and October and are followed by five months of dryness (November to March). The Harmattan is a dry and dusty breeze that blows through the area from late December to January and is characterised by a grey haze that limits visibility and blocks the sun's beams [29]. The temperature in the study area is generally 27-30°C between June and December but rises to 32-34°C between January and April, with the month of March experiencing the hottest temperature 36-38°C. The relative humidity in the rainy season is 82.37% and 74.25% in the dry season.




Fig. 1. Map of Nnamdi Azikiwe University, Awka showing the Honeybee Research Centre

Source: Researcher's Field Work and GIS Mapping, 2023

2.2 Experimental Design

The experiment was a one-factor experiment with three replications in a Completely Randomised Design (CRD). (Fig. 2). The four hive cover designs constructed for this study varied only in the type of insulating materials used for their hive covers and were assigned the following: Control: hive covers insulated with plywood; T1: hive covers insulated with warped boards; T2: hive covers insulated with PVC (Polyvinyl chloride) ceiling and T3: hive covers insulated with foam. The hives were labelled

according to the hive cover designs which they are designated and scattered randomly in the experimental area.

2.3 Sources of Materials

The hard wood species, *Milicia excelsa* (Iroko) used for the hive construction were obtained from the timber market, Umuokpu, Awka, Anambra State, Nigeria. The different insulating materials namely: plywood, warped wood, PVC ceiling and foam used for the various hive cover designs were purchased from Building materials section of Eke Awka market, Awka.




Fig. 2. Illustration of the experimental set up

2.4 Hive Construction and Installation

A total of twelve (12) modified frame bar hives constructed with *M. excelsa* grouped into four hive cover designs namely: Control, T1, T2 and T3 were constructed for this study with each hive treatment replicated three times and labeled as: Control_{I-III}, T1_{I-III}, T2_{I-III} and T3_{I-III}. The timber boards used for the hive construction were piled with spacers in well aerated timber shed to dry for a period of four months [30]. Each hive consisted of the bottom board (length = 53.34cm, width = 35.56cm), main cover board (length = 55.88cm, width = 38.1cm), brood chamber (length = 52.07cm, width = 36.83cm, depth = 24.9cm). All the hive cover designs were made of wood covered with sheets of corrugated iron. Each hive comprised 10 frame bars of 2cm width with two hive entrances of 1cm in diameter [31] with a bee space of 3mm between the frame bars.

Each hive was cleaned and baited with beeswax. Between 6:00 and 8:00 a.m., the hives were baited with forty grammes (40 g) of beeswax, which was melted and applied with a brush over the hive entrances, frame bars, and main cover boards. The hives were allowed to dry after which they were installed [12]. In the month of January 2022, each of the baited hives was kept on metallic supports (46 cm high) within the apiary. The hives were randomly arranged at a distance of 10 metres apart, with the hive entrances oriented to the east [32,12]. During this experiment, the hives environment were kept clean before and after colonisation [23].

2.5 Data Collection

2.5.1 Determination of the influence of different hive cover designs on the microclimate of hives during the study period

Temperature and relative humidity of the four hive cover designs labelled Control_{I-III}, T1_{I-III}, T2_{I-III} and T3_{I-III} were measured and recorded once a week and average taken per month for a period of 15 months from January, 2022 to March, 2023 with the aid of a hand-held digital thermo hygrometer {Model EUROLAB 411TH} [12]. The thermo-hygrometer was placed outside and inside of each hive for 2 minutes, in order to record the ambient and hive temperature and relative humidity respectively. This was carried out between 7:00 am to 10:00am on each day before and after colonisation of hives [28]. The

temperature and relative humidity ranges of the hive cover designs were determined using the formula adopted by Mogbo [33]:

Temperature range = Maximum temperature - Minimum temperature

Relative humidity range = Maximum relative humidity - Minimum relative humidity

2.5.2 Ascertaining the rate of colonisation of African honeybees as influenced by hive cover designs at various periods

The modified frame bar hives labelled Control_{I-III}, T1_{I-III}, T2_{I-III} and T3_{I-III} with different cover designs were monitored two times per week for one hour between 8:00 am to 9:00 am while the number of hives colonised in each hive cover design and the date of colonisation were recorded. The percentage rate of colonisation per hive cover design was computed using the formula adopted by Babarinde et al. [32] and Ononye and Akunne, [12].

Colonisation rate per hive cover design = (Number of colonised hives per hive cover design / Number of hives installed) X 100

2.6 Data Analysis

Data collected from hive temperatures, relative humidity, colonisation, were subjected to One-Way Analysis of Variance (ANOVA) while sample means were separated using Tukey-HSD Test at 5% significant level (P<0.05) using SPSS computer package (version 25) (IBM Corp., 2011). However, Microsoft Excel, 2019 were used to plot the graphs.

3. RESULTS

Table 1 revealed that the Control had the highest temperature (31.86°C) followed by T3 (31.76°C) while T1 (30.40°C) had the least. There was significant difference in the hive temperatures recorded among the four hive cover designs (P=0.00). Fig. 3 revealed that the hive temperature was higher in the month of March, 2023 (32.72°C) followed by the month of February, 2023 (32.56°C) while August (29.87°C) had the least. There was a significant difference in the hive temperature among the various months (P=0.00). The interaction between hive cover designs and different months of the study period had significant effect on the hive temperature (P=0.01).

Table 1. Average minimum and maximum hive temperature, temperature variation, ambient temperature and mean hive temperatures

Hive cover designs	Min. average temp. (°C)(a)	Max. Average temp. (°C) (b)	Range (b-a) (°C)	Min. average ambient temp. (°C)(a)	Max. Average ambient temp. (°C) (b)	Mean ambient temperature	Mean hive temperature (°C) ± SD
Control	23.81	38.44	14.63	22.13	36.76	29.87	31.86 ^b ±3.495
T1	27.21	32.16	4.95	22.13	36.76	29.87	30.40 ^a ±1.351
T2	28.25	34.37	6.12	22.13	36.76	29.87	31.23 ^b ±1.490
T3	29.21	34.09	4.88	22.13	36.76	29.87	31.76 ^b ±1.156

KEY:
 Control (Plywood)
 T1 (Warped wood)
 T2 (PVC)
 T3 (Foam)




Fig. 3. Mean hive temperature of hive cover design at different months of the study period

Table 2. Average minimum and maximum hive humidity, humidity variation, mean ambient humidity and mean humidity of the four hive cover designs

Treatment	Min. average humidity (%) (a)	Max. Average humidity (%) (b)	Range (b-a) (%)	Min. average ambient humidity (%)	Max. Average ambient humidity (%)	Mean ambient hive relative humidity (%)	Mean hive relative humidity (%) ± SD
Control	40.00	84.00	44.00	94.00	48.00	70.69	64.38 ^a ±15.448
T1	47.00	92.00	45.00	87.00	45.00	70.69	73.18 ^b ±14.834
T2	45.00	89.00	44.00	86.00	44.00	70.69	70.91 ^{ab} ±14.670
T3	46.00	89.00	43.00	83.00	43.00	70.69	70.62 ^{ab} ±14.621

*a, b = superscripts for sample mean separation using Tukey-HSD Multiple Range Test
Columns sharing similar superscripts are not significantly different at P>0.05*




Fig. 4. Mean relative humidity of hive cover designs at different months of the study period

Table 3. Colonisation of African honeybees as influenced by different hive cover designs at different months of the study period

Hive Cover Designs	No. of Hives Installed	Months of colonisation															Total hives colonised (%)
		2022												2023			
		Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	
Control	3	0	0	0	0	0	0	1	0	0	1	1	0	0	0	0	3(100.00)
T1	3	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1(33.33)
T2	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0(0.00)
T3	3	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1(33.33)
Total (%)	12	0	0	0	0	0	0	1	0	0	1	1	2	0	0	0	5(41.67)

Table 2 revealed that T1 had the highest relative (73.18%) humidity followed by T2 (70.91%) while the control had the least (64.38%). There was a significant difference in the hive relative humidity among the four hive cover designs ($p=0.04$). Fig. 4 revealed that the hive relative humidity was highest in the month of March, 2023 (32.72°C) followed by February, 2023 (32.56°C) while August, 2022 (29.87°C) had the least. There was a significant difference in the hive relative humidity among the various months ($P=0.00$). The interaction between hive cover designs and different months of the study period had significant effect on the hive relative humidity ($P=0.01$).

Table 3 revealed that the overall colonisation of the African honeybees was 41.67% based on hive cover designs. The colonisation of the African honeybees was observed to be highest in the Control (100%) followed by T1 and T3 (33.33%) but no colonisation was observed in T2 (Plate 1) (0.00%) during the study period. The hive cover designs did not significantly affect colonisation rates of African honeybees ($p=0.26$). Fig. 2 shows the monthly colonisation of African honeybees during the study period. The colonisation of the Control (Plate 2) was observed in the months of July, October, and November, 2022 while that of T1 and T3 (Plate 3 and 4) occurred in December, 2022. However, there was no significant difference in the colonisation of African honeybees among the various months ($p=0.24$).

4. DISCUSSION

The result of this study revealed that the insulating materials used for hive cover designs significantly affected the microclimate (temperature and relative humidity) of hives. This indicates that the use of different insulating materials altered the internal microclimate and this supports previous studies by Lepkova et al. [15] who reported that the microclimate in hives made out of different materials used to design hives greatly influenced the thermal insulation properties the hives. The mean hive temperature values (30.40 - 31.86°C) recorded during this study fall within the recommended hive temperature range for proper development of honeybees. Earlier studies suggested that ensuring optimum hive temperature range (33 to 36°C) is crucial to beekeeping because of the sensitivity of immature bees' cycles to high temperature [8]. This observation is in line with the study by Abd-Elmawgood et al. [11] who reported that hive temperature values of 30 - 36°C are suitable for brood development of honeybee. This study also corroborates with the study by Park et al. [10] who reported that despite the change in external temperature, the internal temperature was maintained above 31°C. Contrastingly, the hive temperature values was slightly higher than that of Tuo et al. [13] and Floris et al. (2020) who reported that bee's activity evolves within adequate temperature range of 34°C and 35°C.



Plate 1. T2 showing no colony formation



Plate 2. Colonised Control hive showing honeybees at the hive entrances



Plate 3. Colonised T1 showing honeybees in the hive

Hive cover designs also affected the internal relative humidity of the hives significantly and the range of values (64.38 - 73.18%) fall short of the optimal relative humidity (52.05%) reported by St. Clair et al. This range falls below the relative humidity values (75.04- 77.94%) earlier reported in the study area by Ononye and Akunne [12] who compared the effect of entrances on hive microclimate. The reason for difference in the relative humidity values could be the variation in

the effect of the various insulating materials used for hive cover design. The result also revealed that hive cover insulated with plywood had the lowest relative humidity when compared to those of polyvinyl chloride ceiling and foam. This is similar to the observation by St. Clair et al. [17] who recently reported that the development of modern insulation materials with higher proprieties could serve as better alternatives wood in the beekeeping sector.



Plate 4. Colonised T3 showing honeybees on the frame bars

The hive cover designs used in this study did not significantly affect colonisation which indicates that colonisation rate may not be dependent on hive cover designs. The wood based designs (plywood and warped wood) recorded the highest colonization by African honeybees when compared with those of foam and PVC during the study period. This supports the observation by Akinbi et al. [21] who also reported that plywood based hives recorded higher percentage colonization when mounted in cocoa farmland and plantation than natural forest. This study agrees with that of Bykov and Zaitsev [16] who reported 91.67% colonisation by honeybee colonies housed in wooden hives. The highest colonization in Control hives could be attributed to the fact that honeybees preferred wood based designs which mimics their natural nest in the wild. Dupleix et al. [22] earlier noted that wooden habitat benefits the bees' well-being because they can analyse heat insulation, humidity control, naturalness and smell. This supports the work by Sahin et al. [34] who stated that the natural surface appearance and color of wood are very important quality criteria in the utilization, especially for outdoor applications [35,36].

Based on the findings of this study, the months of colonisation by the African honeybees housed in control hives was July, October, November while T1 and T3 colonized in December, 2022. The colonization of African honeybees was not significantly different among the various months and this differs from the work by Adedeji et al.

[23] who reported that season has a substantial impact on hive colonisation in the South-South region of Nigeria. Earlier reports on different months of hive colonisation by African honeybees have been given to fall within February – April [23] and October – February [23]. The result of this study is also in contrast with those of Ononye and Akunne [31] who reported August to October as the colonisation months of African honeybees in the study area. Studies by Goulson et al. [25] and Cham [26] attributed variation in the rate of colonisation of hive types to some environmental and anthropogenic factors while the high colonisation was attributed to differences in vegetation cover. The delay in colonization observed in T2 could imply that the hive cover designs was unfavourable and was not preferred by the honeybees. This corroborates with the work report by Akinbi et al. [21] who attributed delay in hive colonisation of honeybees to unfavorable microclimatic conditions in a natural forest ecosystem. This study supports the work by Oluwaseyi et al. [27] who posited that favorable environmental conditions promoted rapid colonization of by honeybees.

5. CONCLUSION

Hive cover designs significantly affected the microclimate of hives but did not significantly affect colonisation of African honeybees. The months of July, October, November may be suitable for the colonisation of African honeybees in the study area. The use of wood-based hive

cover designs should be adopted in the study area but more durable woods should be selected.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ellis J. Colony level thermoregulation and the honey bee dance language. *American Bee Journal*. 2016;147–154.
2. Fasasi KA. Comparative seasonal yield of colonies of *Apis Mellifera adansonii* (Hymenoptera: Apidae) in Response to some Environmental Variables. *Journal of Entomology*. 2016;13(1):11–18. Available: <https://doi.org/10.3923/je.2016.11.18>
3. Poot-Báez V, Medina-Hernández R, Medina-Peralta S, José Javier G, Quezada-Euán. Intranidal temperature and body size of Africanized honey bees under heatwaves (Hymenoptera: Apidae). *Apidologie*. 2020;51:382–390. Available: <https://doi.org/10.1007/s13592-019-00725-5>
4. Li X, Ma W, Shen J, Long D, Feng Y, Su W, Xu K, Du Y, Jiang Y. Tolerance and response of two honeybee species *Apis cerana* and *Apis mellifera* to high temperature and relative humidity. *Plos One*. 2019;14(6):1–18.
5. Abou-shaara HF, Owayss AA, Ibrahim YY, Basuny NK. A review of impacts of temperature and relative humidity on various activities of honey bees. *Insectes Sociaux*. 2017;1–9. Available: <https://doi.org/10.1007/s00040-017-0573-8>
6. Brito ET. Application of phase change material heat storage device for control of *Varroa* mites in beehives. University of Nottingham; 2022.
7. Supandi FA, Gusa RF, Jumnahdi M, Kurniawan R. Design of system for setting the temperature and monitoring bees in and out the hive. *IOP Conference Series: Earth and Environmental Science*. 2020;1–6. Available: <https://doi.org/10.1088/1755-1315/599/1/012050>
8. Gajardo-rojas M, Muñoz AA, Barichivich J, Klock-barr K, Lucas CM. Declining honey production and beekeeper adaptation to climate change in Chile. *Progress in Physical Geography*. 2022;0(0):1–20. Available: <https://doi.org/10.1177/03091333221093757>
9. Brown IL. Beegin Beehive Materials Research Summary. Begin, November; 2018.
10. Park B, Kim G, Kim J, Kim D, Choi Y, Lee M, Kang E. Analysis of termination timing of over-wintering in honey bee (*Apis mellifera*) using digital sensors. *Journal of Apiculture*. 2021;36(3): 97–104. Available: <https://doi.org/10.17519/apiculture.2021.09.36.3.97>
11. Abd-Elmawgood BH, Al-Rajhi MA, El-Ashhab AO. Effect of the internal size and thermal insulation of the hive on bee colonies. *Egypt Journal of Agric Research*. 2015;93(1):185–196.
12. Ononye BU, Akunne C. Microclimate, colonization, colony strength and absconding of *Apis mellifera adansonii* reared in hives with modified entrances. *Journal of Apiculture*. 2019;34(2): 117–123. Available: <https://doi.org/10.17519/apiculture.2019.06.34.2.117>
13. Tuo Y, Coulibaly D, Coulibaly T, Bakayoko S, Hervé Koua K. Role of two agrosystems (mango and cashew trees orchards) in bees' activity increasing within beehives in Korhogo, Northern Ivory Coast (West Africa). *Entomology and Applied Science Letters*. 2020;6(3):48–54.
14. Jaboor SK, Rose C, Da B, Kellermann V. The effect of environmental temperature on bee activity at strawberry farms. *Austral Ecology*. 2022;47:1470–1479. Available: <https://doi.org/10.1111/aec.13228>
15. Lepkova T, Lakov L, Aleksandrov M. Studing the temperature microclimate in beehives made out of defferent materials. *International Scientific Journal Innovations*. 2022;92(2):91–92.
16. Bykov AR, Zaitsev GA. Impact of the design and material of the hives on honey production of bees. *IOP Conference Series: Earth and Environmental Science Paper*. 2021;839:1–6. Available: <https://doi.org/10.1088/1755-1315/839/2/022011>
17. Hassan AH, Kamel SM, Osman MA, Mahmoud MF, Bedeir EH, Shebl MA. Conservation of the mason bees *Osmia latrellei* Spinola (Hymenoptera

- Megachilidae) in Egypt. Egyptian Journal of Applied Science. 2016; 31(6):73-84
18. Prendergast KS. Scientific note: mass-nesting of a native bee *Hylaeus* (Euprosopoides) *ruficeps* kalamundae (Cockerell, 1915) (Hymenoptera: Colletidae: Hylaeinae) in polystyrene. *Apidologie*. 2020; 51(1):107-11.
 19. Wang K, Li J, Zhao L, Mu X, Wang C, Wang M, Xue X, Qi S, Wu L. Gut microbiota protects honey bees (*Apis mellifera* L.) against polystyrene microplastics exposure risks. *Journal of Hazardous Materials*. 2021;402:123828.
 20. St. Clair AL, Beach NJ, Dolezal AG. Honey bee hive covers reduce food consumption and colony mortality during overwintering. *PLoS One*. 2022;17(4):1–17. Available:<https://doi.org/10.1371/journal.pone.0266219>
 21. Akinbi O, Adeduntan S, Toyinbo E, Alamu O. Assessment of Gmelina, Danta Solid Wood and Plywood Hive Types for Beekeeping. *Journal of Applied Science Environment Management*. 2021; 25(4):625–629.
 22. Dupleix A, Julliena D, Moity-Maizi P, Schatzc B. Practices and knowledge of beekeepers and beehive suppliers regarding the wood material in the South of France Practices and knowledge of beekeepers and beehive suppliers regarding the wood material in the South of France. *Journal of Rural Studies*. 2020; 77:11–20. Available:<https://doi.org/10.1016/j.jrurstud.2020.04.005>
 23. Adedeji GA, Aiyeloja AA, Larinde SL, Omokhua GE. Effect of Seasons on Colonisation and Suitability of *Triplochiton scleroxylon* K. Schum. Wood for Beekeeping in Rivers State, Nigeria. *Nature and Science*. 2014;12(8):117–122. Available:<https://doi.org/10.1017/CBO9781107415324.004>
 24. Anumba IA, Akunne CE, Ononye BU. Assessment on Colonization, Absconding and Honey Yield by African Honeybee Colonies Reared in Hives with Different Colours in Awka,. *Journal of Apiculture*. 2020;35(4)199–204. Available:<https://doi.org/10.17519/apiculture.2020.11.35.4.199>
 25. Goulson D, Nicholls E, Botías C, Rotheray EL. Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science*. 2015;347(6229):1255957.
 26. Cham DT. Diversity of honey bee *Apis mellifera* Subspecies (Hymenoptera: Apidae) and Their Associated Arthropod Pests in Cameroon. Ph. D Thesis Submitted to the University of Nairobi. 2017;132.
 27. Oluwaseyi FO, Mustapha MA, Oluwaseyi IG. Effects of temperature and relative humidity on colonization of bees. *Journal of Research in Forestry, Wildlife & Environment*. 2022;14(2):18–25.
 28. Akunne CE, Mogbo TC, Ewuim SC, Ufele AN, Abajue MC, Okafor KC. Relationship between ambient and top-bar hive temperature in relation to optimal production of honey in fallow and cassava plots at science village, Nnamdi Azikiwe University Awka, Nigeria. *Journal of International Academic Research for Multidisciplinary*. 2015;3(3):95–100.
 29. National Statistical Book (NSB). National Stational Book for Anambra State. 2010; 23-342.
 30. Mbobua S. Response to Warre experience in Africa. *Bees for Development Journal*. 2013;106:12.
 31. Ononye BU, Akunne CE. The effects of time and beehive entrance on colony establishment of african honeybees, *Apis mellifera* Adansonii. *Journal of Apiculture*. 2017;32(1):19–26. Available:<https://doi.org/10.17519/apiculture.2017.04.32.1.19>
 32. Babarinde SA, Odewole AF, Oyegoke O, Amao O. Impact of hive dimension and flight entrance on hive colonization, pest infestation and hive weight gain in *Apis mellifera* Adansonii (Hymenoptera: Apidae). *Munis Entomology & Zoology*. 2012;7(1):634–641.
 33. Mogbo TC. Comparison of the growth performance of *Achatina fulica* and *Achatina achatina* as influenced by housing types. *Direct Res J Agric Food Sci*. 2014;2(1):1–7.
 34. Sahin CK, Topay M, Var AA. A study on suitability of some wood species for landscape applications: Surface color, hardness and roughness changes at outdoor conditions. *Wood Research*. 2020; 65(3):395–404. Available:<https://doi.org/10.37763/wr.1336-4561/65.3.395404>

35. Ande AT, Oyerinde A, Jibril MN. Comparative study of the efficacy of six different baiting materials on bee colony performance in traditional and modern hives. *Advances in Biological Research*. 2008;2(1–2):13–16.
36. Meikle WG, Weiss M, Beren E, Meikle WG. Effects of hive entrance orientation on honey bee colony activity. *Journal of Apicultural Research*. 2023;62(3):444–449. Available:<https://doi.org/10.1080/00218839.2023.2165769>

© 2023 Ononye et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/104464>