



Evaluation of the Influence of oral administration of Aqueous Leaf Extract of *Sida acuta* on Lipid Profile and Body Weight of Wistar Rats

Ebere Lolly Mbanaso ^{a*}, Elizabeth Obioma Nweke ^b,
Dominic Chinedu Ejiofor ^c, Abanni Samson ^d,
Iheukwumere Barry Chinedu ^a
and Ngwu Ernest Emeka ^c

^a Department of Human Physiology, Faculty of Basic Medical Sciences, Abia State University, Uturu, Abia State, Nigeria.

^b Department of Human Anatomy, Chukwuemeka Odumegwu Ojukwu University Uli, Anambra State, Nigeria.

^c Department of Human Physiology, Imo State University Owerri, Imo State, Nigeria.

^d Department of Medical Biochemistry, David Nweze Umahi Federal University of Medical Sciences, Uburu, Ebonyi State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJBGMB/2024/v16i1351

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/108420>

Original Research Article

Received: 22/08/2023
Accepted: 28/10/2023
Published: 03/01/2024

*Corresponding author: E-mail: lolymba101@gmail.com;

ABSTRACT

Abnormalities in serum lipid profiles play a central role in endothelial functional abnormality which is a major risk factor for atherosclerosis including coronary artery disease, cerebrovascular disease and peripheral vascular diseases. Hepatic damage disturbs hepatic functions such as lipid metabolism which can translate to a distorted lipid profile. Therefore, the aim of this study was to evaluate the influence of oral administration of *Sida acuta* on the lipid profile and body weight of Wistar rats. Freshly harvested leaves of *Sida acuta* was dried under room temperature and afterwards ground to powder which was subsequently processed into extract. Thirty adults male Wistar rats were divided into five groups of five rats each. Group I (normal control) was administered with 2 ml of distilled water, groups II–VI were induced hepatic damage. While group II was left untreated, groups III-V were administered with 100, 200 and 300 mg/kg of aqueous extract of *Sida acuta* (AESD) respectively. Group VI was administered with silymarin (standard drug). Treatment lasted for 28 days after which animals were humanely sacrificed and blood sample collected for analysis. Treatment with extract significantly ($P < 0.05$) reduced the activities of the liver enzymes; aspartate aminotransferase (AST), alanine transaminase (ALT), alkaline phosphatase (ALP). While the serum levels of TC, TG, HDL and LDL reported for group II were significantly ($P < 0.05$) higher than those reported for the normal control. Oral administration of extract significantly ($p < 0.05$) reduced the levels of TC, TG, HDL and LDL reported for groups III-V. Oral administration of extract did not adversely affect the body weight of animals. In conclusion, it can be deduced from this study that oral administration of *Sida acuta* reversed a distorted lipid profile and did not result in weight loss.

Keywords: Lipid; liver; enzymes; silymarin; lemongrass.

1. INTRODUCTION

“The liver is not only the largest organ of the body but also one the organs critical to the human organism. It plays cardinal roles in lipid synthesis and transportation among numerous functions” [1]. Thus, it is not in doubt that impaired liver function can translate to abnormal lipid profile [2,3].

Human exposure to pollutants such as carbon black resulting from incomplete combustion of fossil fuel as well as diesel exhaust particles discharged by the diesel-powered trucks and automobiles which constitute a better part of the atmospheric particulate matter in urban centers has tremendously increased and have been positively correlated with increased human mortality through various diseases notably hepatic damage among others through oxidative stress [4]. Conventionally, treatment of liver damage involves the use of synthetic drugs which may further hamper hepatic health [5].

The earth is endowed with diverse plants of therapeutic values globally utilized by an estimated 80% of the world's population in

pursuance of health care needs [6]. *Sida acuta* commonly referred to as the ironweed is abundantly around improved pastures, wastes, disturbed places and roadsides [7].

Although native to Mexico and Central Africa, it can now be seen throughout the tropics and subtropics [8]. In addition to the fact that it has been used successfully in the treatment of diseases such as fever, headache, skin infections, diarrhea and dysentery [1] numerous pharmacological activities such anti-plasmodial, cytotoxic and antioxidant properties [9] are attributed to the said plant.

2. MATERIALS AND METHODS

2.1 Collection of Plant Material

Fresh leaves of *Sida acuta* were collected from a bush within a residential area in Amaokwe Uturu, Isiukwuato Local Government Area of Abia State in the Southeastern part of Nigeria. The leaves were transported in a dark polythene bag to the herbarium unit of the Department of Forestry, Michael Okpara University of Agriculture, Umudike Abia State, and Southeast Nigeria for identification.

2.2 Preparation of Extract

The leaves of *Sida acuta* were properly washed with the aid of clean tap water in order to remove dirt. This was followed by drying at room temperature. Afterwards, the leaves were ground and sieved to fine powder. Exactly 500 g of powdered plant sample was subsequently steeped in 3 L of distilled water for 24 h. The mixture was filtered with a clean sieve and was concentrated to dryness in a water bath for 3 days at 50°C.

2.3 Animals

Adult male Wistar rats weighing 120-150 g were bought from the animal house of the Department of Science Laboratory Technology, Akanu Ibiam Federal Polytechnic Unwana, Afikpo, Ebonyi State. The rats were housed in plastic cages under standard laboratory conditions. They were allowed access to food and water *ad-libitum*. Animals were acclimatized for 14 days.

2.4 Median Lethal dose 50% (LD50)

The LD₅₀ determination was conducted in two phases. In the first phase, nine (9) adult male rats were divided into three groups of three rats each. The groups were separately administered 10, 100 and 1000 mg/kg of extract orally. Afterwards, animals were observed closely for 24 h. for possible signs of toxicity. In the absence of mortality at the first phase, the second phase was initiated and three groups of one rat per group were involved. The various groups of the second phase were separately administered 1600, 2900 and 5000 mg/kg of extract, after which signs of toxicity were looked out for on the animals for a period of 48 h Lorke [10].

2.5 Animal Grouping

Group I (Normal Control): animals were administered distilled water.

Group II (Negative Control) animals induced hepatic damage by oral administration of 600 mg/kg/day of paracetamol without treatment

Group III: Rats with hepatic damage were administered 100 mg/kg of extract

Group IV: Rats with hepatic damage were administered 200 mg/kg of extract

Group V: Rats with hepatic damage were administered 300 mg/kg of extract

Group VI: Rats with hepatic damage administered standard drug (silymarin)

Administration of extract lasted for 28 days after which animals were sacrificed and blood sample collected for analysis.

2.6 Liver Function Test

The activities of liver enzymes; aspartate aminotransferase (AST), alanine transaminase (ALT), alkaline phosphatase (ALP) were determined by introducing 2 mL of blood into an EDTA tube and afterwards centrifuged at 4,000 rpm for 15 min. The resulting plasma was analyzed with the aid of kits.

2.7 Determination of Lipid Profile

Cholesterol, HDL and triacylglyceride levels were estimated from serum by CHOD-PAP. LDL and HDL were calculated, while the atherogenic index was calculated using the method described by Muruganandan et al. [11].

2.8 Body Weight Measurement

Determination of animal weight was performed at the commencement of the study (day 0) and before termination of experiment.

2.9 Statistical Analysis

Data generated from the study were expressed as mean \pm standard deviation using SPSS (Ver. 23). Data were analysed using one way analysis of variance (ANOVA). Variation in mean values was compared using Turkey Test. *P-values* less than 0.05 was considered statistically significant.

3. RESULTS AND DISCUSSION

Cellular damage either orchestrated naturally or artificially is mediated mainly by free radical generation. Table 1 shows the activity of liver enzymes in rats with damaged hepatocytes treated with aqueous extract of *Sida acuta* indicating that the activity of AST, ALT and ALP in rats induced hepatic damage without treatment was significantly ($p < 0.05$) higher than that reported for the normal control. However, following oral administration of 100, 200, and 300 mg/kg of *Sida acuta* extract, a significant reduction in the activity of the aforementioned enzymes was observed which however was significantly ($p < 0.05$) lower than that reported for the group administered standard drug which in turn was not significantly ($p > 0.05$) different from

that reported for the normal control. This could be attributed to the influence of flavonoids and zinc which are appreciably present in the leaf of *Sida acuta* (Nwankpa et al., [12]. This result is consistent with the finding of Nwankpa et al. [12] which reported the antioxidant properties of *Sida acuta* extract. Damage to the liver will undoubtedly impair hepatic functions including hepatic lipid metabolism which can translate to a distorted lipid profile. Table 2 shows the lipid profile of Wistar rats administered aqueous extract of *Sida acuta* indicating that Total Cholesterol (TC), Triacylglycerol (TG), High Density Lipoprotein (HDL), and Low Density Lipoprotein levels in group II were significantly

($p < 0.05$) higher than those reported for the normal control which in turn were not significantly ($p < 0.05$) different from those reported for the groups administered *Sida acuta* extract as well as the group administered the standard drug (silymarin). “The potential of the said extract to restore a normal lipid profile of the animals could be attributed to the presence of flavonoids reportedly present in the leaf of the plant” Nwankpa et al. [12]. This is consistent with the finding of Ogbodo et al. [13] which demonstrated the anti-hyperlipidemic effect of the leaf extract of *Sida corymbosa* a member of the malvaceae family to which *Sida acuta* belongs. Liver damage can cause symptoms such as poor

Table 1. Liver Enzyme activity in Rats with Hepatic Damage treated with Aqueous Extract of *Sida acuta* (Ironweed)

Treatment	AST (U/l)	ALT(U/l)	ALP(U/l)
Group I (Normal CTRL)	9.12±1.20 ^a	14.00±2.67 ^a	75.80±2.24 ^a
Group II Negative CTRL	34.00±4.10 ^c	27.01±4.10 ^d	116.01±4.21 ^c
Group III (100 mg/kg Ext.)	11.00±3.66 ^b	21.06±3.27 ^b	84.00±2.00 ^b
Group IV (200 mg/kg Ext.)	11.80±4.34 ^b	22.07±4.36 ^{bc}	82.00±1.43 ^b
Group IV (300 mg/kg Ext.)	11.00±5.67 ^b	21.30±4.66 ^b	81.02±1.65 ^b
Group IV (Std. silymarin)	9.08±2.20 ^a	14.04±4.26 ^a	76.00±6.52 ^a

Results are expressed as mean ± standard deviation of three determinations. Values with different superscripts in a column are significantly different at $P < 0.05$

Table 2. Lipid Profile of Wistar Rats administered Aqueous Extract of *Sida acuta* (Ironweed)

Treatment	TC (mg/dl)	TG (Mg/dl)	HDL (Mg/dl)	LDL (Mg/dl)
Group I (Normal CTRL)	203±11.53 ^a	63±10.56 ^a	71±4.56 ^d	127±42.56 ^a
Group II Negative CTRL	222±2.06 ^b	73±7.06 ^b	84±5.46 ^c	136±45.56 ^c
Group III (100 mg/kg Ext.)	206±6.31 ^{ab}	64±5.01 ^{ab}	75±4.65 ^b	130±2.51 ^d
Group IV (200 mg/kg Ext.)	203±4.21 ^a	62±2.36 ^a	70±2.46 ^c	130±1.16 ^b
Group IV (300 mg/kg Ext.)	204±5.05 ^a	63±21.56 ^a	72±3.52 ^b	131±5.32 ^b
Group IV (Std. silymarin)	202±6.30 ^a	62±1.59 ^a	73±12.46 ^d	128±5.16 ^a

Results are expressed as mean ± standard deviation of three determinations. Values with different superscript in a column are significantly ($p < 0.05$) different

Table 3. Body weight of wistar rats administered Aqueous Extract of *Sida acuta* (Ironweed)

Treatment	Initial wt. (g)	Final wt. (g)
Group I (Normal CTRL)	125.23±8.72	157.20±6.00
Group II Negative CTRL	135.64±4.50	130.40±5.63
Group III (100 mg/kg Ext.)	167.32±5.27	175.65±2.40
Group IV (200 mg/kg Ext.)	147.23±6.19	165.30±5.60
Group IV (300 mg/kg Ext.)	175.03±6.27	187.03±5.76
Group IV (Std. silymarin)	167.30±6.39	178.21±6.50

Results are expressed as mean ± standard deviation of three determinations. Values with different superscripts in a column are significantly different at $P < 0.05$

appetite which can translate to weight loss. Table 3 shows the body weight of rats administered aqueous extract of *Sida acuta*, showing that there was a significant increase in the weight of rats over treated periods. However, a contrary observation was made on group II which was induced with hepatic damage without treatment. Weight increase observed on treated rats could be attributed to the reversal of damaged liver cells to healthy ones and hence improved consumption of meal.

4. CONCLUSION

Through this research effort, it can be deduced that oral administration of *Sida acuta* reversed hepatic damage, distorted lipid profile and did not result in weight loss.

ETHICAL APPROVAL

Animal Ethic committee approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ozougwu JC, Eyo JE. Hepatoprotective effects of *Allium cepa* extracts on paracetamol-induced liver damage in rat. *Afr. J. Biotechnol.* 2014;13(26):2679-2688.
2. Allen SE. *The liver: Anatomy, Physiology, Disease and Treatment.* North Eastern University Press, USA; 2002.
3. Halsted CH. Nutrition and alcoholic liver disease. *Semin Liver Dis.* 2004;24(3):289-304.
4. Son JY, Lee JT, Kim KH, Jung K, Bell ML. Characterization of fine particulate matter and associations between particulate chemical constituents and mortality in Seoul, Korea. *Environ. Health Perspect.* 2012;120:872-878.
5. Ostapowicz G, et al. Results of a prospective study of acute liver failure at 17 tertiary care centers in the United States. *Ann Intern Med.* 2002;137:947-54.
6. Newman, DJ, Cragg GM. Current Status of Marine-Derived Compounds as Warheads in Anti-Tumor Drug Candidates *Mar. Drugs.* 2017;15:99.
7. Mann A, Gbate M, Umar AN. *Sida acuta* subspecies *acuta*. Medicinal and economic plant of Nupeland, Jube Evans Books and Publication. 2003;241.
8. Holm LG, Plucknett DL, Pancho JV, Herberger JP. *The world's worst weeds: distribution and biology.* University Press of Hawaii, Honolulu, USA; 1977.
9. Karou D, Dicko MH, Sanon S, Simporé J, Traore SA. Anti-malarial activity of *Sida acuta* BURMF L. (Malvaceae) and *Pterocarpus erinaceus* POIR (Fabaceae) J. *Ethnopharmacol.* 2003;89:291-294.
10. Lorke DA new approach to practical acute toxicity testing. *Arch of Toxicol.* 1983;54(4): 275-287.
11. Muruganandan S, Scrivinasan K, Gupta S, Gupta PK, Lal J. Effect of mangiferin on hyperglycemia and atherogenicity in streptozotocin diabetic rats. *J Ethnopharmacol.* 2005;97:497-501.
12. Nwankpa P1, Chukwuemeka OG, Uloneme GC, Etteh CC, Ugwuezumba P, Nwosu D. Phyto-nutrient composition and antioxidative potential of ethanolic leaf extract of *Sida acuta* in wistar albino rats. 14(49):3264-326.
13. Ogbodo EC, Ezeugwunne IP, Otuokwu BC, Onyegbule OA, Analike RA, Amah AK. Oguaka VN. Effect of *Sida corymbosa* leaf extract on the serum lipid profile levels of alloxan-induced diabetic albino wistar rats in college of health sciences and technology, World J. Pharm. Res. 7(11):51-59.

© 2024 Mbanaso et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/108420>