



Microbial Populations of Agricultural Soil Polluted With Crude Oil

Soludo, O. C. ^{a*}, Orji, M. U. ^a, Anaukwu, C. G. ^a,
Anyaocha, V. I. ^a, Ajogwu, T. M. C. ^a and Eze, H. C. ^a

^a Department of Applied Microbiology and Brewing, Nnamdi Azikiwe University, Awka, P.M.B., 5025, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JABB/2024/v27i2706

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/107825>

Original Research Article

Received: 24/08/2023

Accepted: 29/10/2023

Published: 28/02/2024

ABSTRACT

Crude oil pollution of the soil has become a public health and environmental concern. It impacts soil microbial diversity and population. In this study, the microbial population of crude oil polluted soil and unpolluted were determined. The potential for the indigenous organisms to utilize hydrocarbon as a source of carbon was also determined. A soil sample was obtained from an agricultural field and polluted with a defined amount of crude oil for 12 weeks. The microbial population in the polluted soil and unpolluted soil were estimated and characterized following standard microbiological methods. The heterotrophic bacterial and fungal counts in the unpolluted and polluted soil were 6.5×10^4 CFU/g and 4.7×10^4 CFU/g, and 6.3×10^4 CFU/g and 3.5×10^4 CFU/g respectively. Total hydrocarbon utilizing bacterial and fungal counts in the polluted soil was 4.6×10^4 CFU/g and 2.8×10^4 CFU/g. The genera of bacteria and fungi identified in the soil samples were *Bacillus*, *Enterococcus*, *Micrococcus*, *Escherichia*, *Pseudomonas*, *Staphylococcus*, *Serratia*, *Proteus*, *Klebsiella*, *Arthrobacter*, *Aspergillus*, *Candida*, *Penicillium*, *Fusarium*, *Trichoderma*, *Cladosporium* and *Hyphopichia*. The preliminary screening for hydrocarbon utilization shows that

*Corresponding author: E-mail: oc.soludo@unizik.edu.ng;

Bacillus cereus, *Bacillus subtilis*, *Aspergillus niger* and *Hypopichia burtonii* were excellent hydrocarbon utilizers and can be used in bioaugmentation.

Keywords: Hydrocarbon; degradation; crude oil; bacterial count; fungi count.

1. INTRODUCTION

“Soils are complex mixtures of minerals, water, air, organic matter and countless organisms that are the decaying remains of once-living things” [1]. “The Natural Resource Conservation Service (NRCS) defines soil as a natural body comprised of solids (minerals and organic matter), liquid, and gases that occur on the land surface, occupy space, and are characterized by one or both of the following: horizons, or layers, that are distinguishable from the initial material as a result of additions, losses, transfers, and transformations of energy and matter or the ability to support rooted plants in a natural environment” [2]. “Soil plays a vital role in sustaining life on the planet and is also a habitat for many microorganisms. Nearly all of the food that humans consume, except for what is harvested from marine environments, is grown in the Earth’s soils” [3]. “Soil is responsible for nutrient cycling, plant growth, gas exchange, carbon storage, and waste disposal” [4].

“The contamination of the soil by frequent crude oil spillages renders the soil toxic and agriculturally unproductive. The oil reduces the soil’s fertility to the extent that most of the essential nutrients are no longer available to plants and for crop utilization” [5]. “Crude oil as a pollutant, decreases the nitrogen and phosphorus contents of the soil and promotes excessive hydrocarbon which affects soil enzymatic activities, due to the inability of soil microbes to degrade the excess hydrocarbons” [6]. “The contamination of soil by crude oil and petroleum products has become a serious problem that represents a global concern for its potential consequences on the ecosystem and human health” [7,8]. Crude oil contaminations on land affect the physicochemical properties of the soil which causes deleterious effects on plant germination and growth. Soil properties such as; soil texture, moisture content, pH and bulk density are degraded [9-11]. This leads to poor root and leaf development, which consequently affects plant growth and yield. Contaminated land can present a risk to human health, the immediate ecosystem, and the environment. Oil spills cause soil pollution and eventual

environmental degradation [1,12]. “Leakages and accidental spills commonly occur during the exploration, production, refining, transport, and storage of petroleum and petroleum products” [8].

“However, crude oil-polluted soil can be remediated by the actions of indigenous microorganisms with an inherent ability to utilize hydrocarbon as a carbon source. Microorganisms are capable of degrading a wide range of hydrocarbon molecules in the environment, especially the lithosphere, where they attack and digest petroleum hydrocarbons; this thus forms the emergence of the field of bioremediation” [13]. Indigenous microorganisms in hydrocarbon-polluted sites have been reported to be excellent hydrocarbon utilizers and can be employed for the reclamation of polluted environments.

The present study aimed at identifying the microbial populations in crude oil-polluted and unpolluted soil, and to determine their potential for hydrocarbon utilization.

2. MATERIALS AND METHODS

2.1 Collection of Samples

An agricultural soil sample (unpolluted soil) was obtained from an agricultural field in Nnamdi Azikiwe University, Awka, Nigeria in a sterile sample bag using a sterile hand trowel and digging 15cm depth in the soil. The sample was conveyed to the laboratory for analysis.

The crude oil used in the experiment was obtained from a Petrochemical Company situated at Eleme Oil field, Eleme Local Government Area in Port Harcourt, Rivers state of Nigeria.

2.2 Sample Preparation

One hundred grams of the soil sample was polluted with 100 ml of crude oil and evenly mixed to obtain a homogenous mixture. The mixture was allowed to stand at room temperature for 12 weeks, during which there was periodic turning of the soil for aeration.

2.3 Isolation of Microorganisms from the Soil Samples

The method described by Onuorah et al. [13] was used for the isolation of bacterial and fungal organisms from the polluted and unpolluted soil samples. Ten-fold serial dilutions were carried out using 1g each of the samples in 10 ml of sterile distilled water. 1 ml of the solvent was introduced into test tubes containing 9 ml of sterile water and serially diluted from 10⁻¹ to 10⁻¹⁰.

2.3.1 Enumeration of total heterotrophic bacteria

From the serially diluted samples, one millilitre of 10⁻⁴ was inoculated onto a sterile nutrient agar plate by pour plating technique in triplicate. The plates were incubated at 25°C and colonies that developed after 24 hours were counted in the triplicate plates, and mean values were recorded and used to calculate the colony forming unit per gram (CFU/g). Distinct colonies were sub-cultured and stored at 40C for further identification.

2.3.2 Enumeration of total fungi

From the serially diluted samples, one millilitre of 10⁻⁴ was inoculated onto a sterile Sabouraud dextrose agar plate fortified with chloramphenicol to suppress bacterial growth by pour plating technique in triplicate. The plates were incubated at 25°C and colonies that developed after 72 hours were counted in the triplicate plates, and mean values were recorded and used to calculate the colony forming unit per gram (CFU/g). The colonies that developed were thereafter sub-cultured to obtain pure cultures. The pure cultures obtained were transferred into agar slants for identification.

2.4 Isolation of Hydrocarbon-utilizing Bacteria and Fungi from the Soil

The method described by Olukunle [14,15] was used. The crude oil-polluted soil sample was diluted tenfold at a series of 10⁻¹ to 10⁻¹⁰. An aliquot of 0.5ml from 10⁻⁴ was inoculated using the pour plate method into a mineral salt agar for bacteria and Sabouraud dextrose agar fortified with chloramphenicol for fungi. A Whatman filter paper impregnated with crude oil was placed on the inside cover of the petri dish, covered and incubated for 24h - 48h at 25°C in an inverted position for bacterial isolates and 72h at 25°C for fungi isolates. The crude oil-impregnated filter

paper supplied hydrocarbons by vapour phase transfer to the inocula. The test was done in triplicates. Developed colonies on the culture plates were counted and mean values were used to calculate the colony-forming unit per gram (cfu/g).

2.4.1 Identification of bacterial and fungal Isolates

The identification of bacteria was done based on morphological characteristics and biochemical tests. Morphological characteristics observed for each bacteria colony after 24-48 hours of growth include colony appearance; shape, elevation, edge, optical characteristics, consistency, colony surface and pigmentation. Biochemical tests carried out for the identification of bacterial isolates were Gram stain test, motility test, catalase test, coagulase test, citrate test, indole test, oxidase test, methyl red test, spore stain test, Voges Proskauer test, glucose fermentation, sucrose fermentation, lactose fermentation and mannitol fermentation. The results were compared to Bergey's manual of determinative bacteriology [16].

The fungal isolates were identified based on morphological and cultural characteristics, lactophenol cotton blue stain and slide culture test.

2.4.2 Lactophenol cotton blue staining technique

A drop of Lactophenol cotton blue solution was placed on a clean grease-free slide. A sterilized straight wire was used to transfer, tease and properly mix the organism with the solution on the slide. A cover slip with the help of a pair of forceps was placed on the mycelial materials. Excess stain was wiped with blotting paper and the slide was viewed under a microscope using x40 lens magnification for the presence of hyphae, mycelia and spores [17].

2.4.3 Slide culture technique

A sterile Sabouraud dextrose agar medium was prepared and inoculated with a mould isolate. A cover slip was inserted beside the mould and incubated in an inverted position for 4-7 days. The cover slip was placed on a slide containing a drop of lactophenol blue and was examined with 10x and 40x microscope objectives for the presence of fungal features. Fungal isolates were identified following the Scheme of Oyeleke and Manga, [18].

2.5 Preliminary Screening for Hydrocarbon degradation by the Microbial Isolates from the soil samples

The method described by Ramaraj [19] was used to screen the bacterial and fungal isolates for hydrocarbon degradation. The microorganisms isolated from the crude oil-polluted soil sample were used for the test. Mineral salt medium (MSM) supplemented with 1% crude oil was prepared and sterilized in an autoclave at 121°C for 15 minutes. A loopful of a 24-hour-old culture of the isolate was inoculated into 50 ml MSM in a 100 ml Erlenmeyer flask. The flask was incubated in a rotary shaker for 14 days. Triplicate flasks were used and an uninoculated flask served as control. Bacteria and fungi growth in the medium were indicated by turbidity and clarity respectively.

3. RESULTS

The heterotrophic bacterial and fungal counts of the crude oil-polluted and unpolluted soil samples are shown in Table 1. The total heterotrophic bacteria and fungi are 6.5×10^4 CFU/g and 4.7×10^4 CFU/g respectively in the unpolluted soil and 6.3×10^4 CFU/g and 3.5×10^4 CFU/g in the polluted soil.

The petroleum hydrocarbon utilizing bacterial and fungal counts was estimated in the polluted soil using the vapour phase transfer method [20]. The bacterial colonies were identified to their species level using conventional microbiological biochemical tests as described by [16,21] using Bergey's Manual of Determinative Bacteriology. The total hydrocarbon utilizing bacteria and fungi are 4.6×10^4 CFU/g and 2.8×10^4 CFU/g respectively (Table 2).

Table 3 shows the morphological and biochemical characteristics of the heterotrophic bacteria isolated from the unpolluted soil, and the organisms isolated are *Bacillus cereus*,

Enterococcus faecalis, *Micrococcus luteus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Serratia marcescens* and *Bacillus subtilis*. Table 4 shows the morphological and biochemical characteristics of hydrocarbon utilizing bacteria isolated from the crude oil-polluted soil. The organism isolated are *Bacillus cereus*, *Pseudomonas aeruginosa*, *Serratia marcescens*, *Bacillus subtilis*, *Micrococcus luteus*, *Proteus vulgaris*, *Klebsiella aerogenes* and *Arthrobacter oxydans*.

Table 1. Heterotrophic bacterial and fungal counts in the soil samples

Sample	Count (CFU/g)
Unpolluted soil	
Total heterotrophic bacteria	6.5×10^4
Total heterotrophic Fungi	4.7×10^4
Crude oil-polluted soil	
Total heterotrophic bacteria	6.3×10^4
Total heterotrophic Fungi	3.5×10^4

The colonial and microscopic features of the heterotrophic fungi isolated from the unpolluted soil are shown in Table 5. The organisms isolated are *Aspergillus niger*, *Candida utilis*, *Penicillium expansum*, *Fusarium oxysporium*, *Trichoderma herbarum* and *Cladosporium resinae*. Table 6 shows the colonial and microscopic features of the hydrocarbon utilizing fungi isolated from crude oil-polluted soil. The isolates are *Aspergillus niger*, *Penicillium expansum*, *Hyphopichia burtonii*, *Candida utilis*, *Fusarium oxysporium*, *Cladosporium resinae* and *Candida tropicalis*.

Table 7 shows the result of the screening test for hydrocarbon utilization by the microbial isolates from the crude oil-polluted soil. *Bacillus cereus*, *Aspergillus niger*, *Hyphopichia burtoni* and *Bacillus subtilis* had heavy turbidity (+++), *Candida tropicalis* had moderate turbidity (++) while *Candida utilis* and *Penicillium expansum* had minimal turbidity (+).

Table 2. Petroleum Hydrocarbon Utilizing Bacterial and Fungal Count

	Count (CFU/g)
Petroleum hydrocarbon utilization bacteria	4.6×10^4
Petroleum hydrocarbon utilization fungi	2.8×10^4

Table 3. Morphological and Biochemical characteristics of the heterotrophic bacteria isolated from the unpolluted soil

Isolates	Gram stain	Form	Spore stain	Motility	Catalase	Oxidase	Citrate	Indole	Methyl red	Voges Proskauer	Coagulase	Glucose	Sucrose	Lactose	Mannitol	Identity
A	+	Coccus	-	-	-	-	+	-	+	+	-	+	+	+	+	<i>Enterococcus faecalis</i>
B	+	Coccus	-	-	+	+	-	-	-	-	-	-	-	-	-	<i>Micrococcus luteus</i>
C	+	Rod	-	-	-	-	-	-	-	-	-	-	-	-	-	<i>Escherichia coli</i>
D	+	Rod	+	+	+	-	+	-	-	+	-	+	-	-	-	<i>Bacillus cereus</i>
E	-	Rod	-	+	+	+	+	-	-	-	-	-	-	-	-	<i>Pseudomonas aeruginosa</i>
F	+	Coccus	-	-	+	-	-	-	+	+	+	+	+	+	+	<i>Staphylococcus aureus</i>
G	-	Rod	-	-	+	-	+	-	-	+	-	+	+	-	+	<i>Serratia marcescens</i>
H	+	Rod	+	-	+	-	+	-	-	+	-	-	-	-	+	<i>Bacillus subtilis</i>

KEY: + = Positive, - = Negative

Table 4. Morphological and Biochemical Characteristics of Hydrocarbon-Utilizing Bacteria Isolated from the Crude Oil-Polluted Soil

Isolates	Gram stain	Form	Spore stain	Motility	Catalase	Oxidase	Citrate	Indole	Methyl red	Voges Proskauer	Coagulase	Glucose	Sucrose	Lactose	Mannitol	Identity
A	+	Rod	+	+	+	-	+	-	-	+	-	+	-	-	-	<i>Bacillus cereus</i>
B	-	Rod	-	+	+	+	+	-	-	-	-	-	-	-	-	<i>Pseudomonas aeruginosa</i>
C	-	Rod	-	-	+	-	+	-	-	+	-	+	+	-	+	<i>Serratia marcescens</i>
D	+	Rod	+	-	+	-	+	-	-	+	-	-	-	-	+	<i>Bacillus subtilis</i>
E	+	Coccus	-	-	+	+	-	-	-	-	-	-	-	-	-	<i>Micrococcus luteus</i>
F	-	Rod	-	+	+	-	-	+	+	-	-	+	+	-	-	<i>Proteus vulgaris</i>
G	-	Rod	-	-	+	-	-	+	+	+	-	+	+	+	+	<i>Klebsiella aerogenes</i>
H	-	Rod	-	-	+	-	+	+	+	-	-	-	-	-	-	<i>Arthrobacter oxydans</i>

KEY: + = Positive, - = Negative

Table 5. Colonial and microscopic features of the heterotrophic fungi isolated from the unpolluted soil

Isolate	Colonial Characteristics	Microscopic Characteristics	Identity
1	Colonies were initially white but quickly formed black conidia. The reverse was yellow-grey.	Hyphae were septate. Conidiophores were long, smooth-walled, hyaline and contained globose vesicles each covered completely with biserial phialides. Conidia were globose, dark and rough-walled. Conidia heads were large, globose, dark brown and biserial.	<i>Aspergillus niger</i>
2	Colonies are cream-coloured, oval, smooth, glabrous, dull, soft and yeast-like in appearance with a yeasty odour.	Sub-spherical budding yeast-like cells called blastospores are present, and terminal chlamydoconidia are absent. Ferment glucose and maltose. Does not ferment lactose.	<i>Candida utilis</i>
3	Dull green mycelia with whitish lining were seen.	Non-ceonocytic (septate) hyphae were seen.	<i>Penicillium expansum</i>
4	Whitish to pink cottony mycelium was seen. The reverse was dark purple.	Conidiophores were short with single lateral momophialides, Microconidia were in fusiforms and pointed at the tip.	<i>Fusarium oxysporium</i>
5	Colonies were slow-growing, dark green with black edges and powdery. The reverse was greenish brown	Conidiophores were erect, unbranched and dark, shield smooth and occurred in long branching chains.	<i>Trichoderma herbarum</i>
6	Colonies were slow-growing, dark green with black edges and powdery. The reverse was greenish brown	Conidiophores were erect, unbranched and dark, shield smooth and occurred in long branching chains.	<i>Cladosporium resinae</i>

Table 6. Colonial and microscopic features of the hydrocarbon utilizing fungi isolated from the crude oil-polluted soil

Isolate	Colonial Characteristics	Microscopic Characteristics	Identity
1	Colonies were initially white but quickly formed black conidia. The reverse was yellow-grey.	Hyphae were septate. Conidiophores were long, smooth-walled, hyaline and contained globose vesicle each covered completely with biserial phialides. Conidia were globose, dark and rough-walled. Conidia heads were large, globose, dark brown and biserial.	<i>Aspergillus niger</i>
2	Dull green mycelia with whitish lining, were seen.	Non-ceonocytic (septate) hyphae were seen.	<i>Penicillium expansum</i>
3	Colonies are tannish white and butyrous to hyphal, ascospores formed are released by ascus, cells are short, ellipsoidal or elongated and may form pseudohyphae and true hyphae.	Yeast like conidia borne laterally and terminally on short denticles along the parental, septate true hyphae.	<i>Hyphopichiaburtonii</i>
4	Colonies are cream-coloured, oval, smooth, glabrous, dull, soft and yeast-like in appearance with a yeasty odour.	Sub-spherical budding yeast-like cells called blastospores are present, and terminal chlamydoconidia are absent. Ferment glucose and maltose. Does not ferment lactose.	<i>Candida utilis</i>
5	Whitish to pink cottony mycelium was seen. The reverse was dark purple.	Conidiophores were short with single lateral monophialides. Microconidia were fusiform and pointed at the tip.	<i>Fusarium oxysporium</i>
6	Colonies were slow-growing, dark green with black edges and powdery. The reverse was greenish brown.	Conidiophores were erect, unbranched and dark, shield smooth and occurred in long branching chains.	<i>Cladosporium resinae</i>
7	Colonies are cream-coloured, oval, smooth, glabrous, dull, soft and yeast-like in appearance with a yeasty odour.	Sub-spherical budding yeast-like cells called blastospores present at both terminal chlamydoconidia are absent. Ferment glucose and maltose, does not ferment sucrose.	<i>Candida tropicalis</i>

Table 7. Screening characteristics for hydrocarbon utilization by the microbial isolates from the crude oil polluted soil

Microorganism	Hydrocarbon utilization
<i>Bacillus cereus</i>	+++
<i>Candida utilis</i>	+
<i>Hyphopichiaburtonii</i>	+++
<i>Penicillium expansium</i>	+
<i>Aspergillus niger</i>	+++
<i>Bacillus subtilis</i>	+++
<i>Candida tropicalis</i>	++

+ Minimal turbidity ++ Moderate turbidity +++ Heavy turbidity

4. DISCUSSION

This estimated and characterized the microbial population in crude oil polluted and unpolluted soil. The total viable count of bacteria from contaminated soil with used hydrocarbon was impressive showing the presence of some bacteria and fungi that utilize hydrocarbon which makes bioremediation effective. Crude oil impaction caused decreased values of all the bacterial and fungal populations as seen in Table 2. The total heterotrophic bacterial and fungal count in the unpolluted soil were 6.3×10^4 CFU/g and 3.5×10^4 CFU/g (Table 1), while that of hydrocarbon-utilizing bacteria and fungi were 4.6×10^4 CFU/g and 2.8×10^4 CFU/g respectively. Morphological and biochemical characteristics of the heterotrophic bacteria isolated from the unpolluted soil are shown in Table 3, the organisms isolated were *Bacillus cereus*, *Enterococcus faecalis*, *Micrococcus luteus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Serratia marcescens* and *Bacillus subtilis* this agrees with the findings of Udebuani et al. [22].

Table 4 shows the morphological and biochemical characteristics of hydrocarbon utilizing bacteria isolated from the crude oil-polluted soil. The organisms isolated were *Bacillus cereus*, *Bacillus subtilis*, *Proteus vulgaris*, *Klebsiella aerogens*, *Athrobacter oxydans*, *Pseudomonas aeruginosa*, *Serratia marcescens* and *Micrococcus luteus*. The hydrocarbon utilizing bacteria isolated conforms to the finding of Chikere et al. [23], in their assessment of bioreactor-based bioremediation of hydrocarbon-polluted Niger Delta marine sediment, Nigeria. This finding again revealed that there is a high population of active indigenous hydrocarbon-utilizing bacteria which can be monitored and enhanced to bring about bioremediation. This agrees with the work of Chikere and Ekwuabu [12] in their study of the

culture-dependent characterization of hydrocarbon utilizing bacteria in the selected crude oil-impacted sites in Bodo, Ogoni land, Nigeria. Again Omotayo et al. [24], isolated similar organisms in their work on the degradation of aviation fuel by microorganisms isolated from tropical polluted soils. Also, Yakubu and Bello [25] isolated similar organisms in their work on the biodegradation of Lagoma crude oil using pig dung.

Table 5 shows the colonial and microscopic features of hydrocarbon-utilizing moulds isolated from the pristine soil. The isolates were *Aspergillus niger*, *Cladosporium resineae*, *Fusarium oxysporium* and *Penicillium expansium*. Chikere and Azubike [12] in their work of characterization of hydrocarbon utilizing fungi from hydrocarbon-polluted sediments and water, isolated similar organisms.

Table 6 shows the colonial and microscopic features of the hydrocarbon utilizing fungi isolated from crude oil-polluted soil. The isolates were *Aspergillus niger*, *Hyphopichia burtonii*, *penicillium expansium*, *Cladosporium resineae*, *Fusarium oxysporium*, *Candida tropicalis* and *Candida utilis*,. Omotayo et al. [24], in their work on the degradation of aviation fuel by microorganisms isolated from tropical polluted soils, isolated *Aspergillus*, *Penicillium* and *Candida*, while Stephen and Panneerselvam [26] in their work on the study of the prevailing fungi on the hydrocarbon polluted soil, isolated similar organism, however, the fungal isolates in the crude oil-polluted soil differ with the result of this study. This could be a result of environmental conditions like pH and the percentage of mineral nutrients in the soil.

Table 7 shows the result of the screening test for hydrocarbon utilization by the microbial isolates from the crude oil-polluted soil before amendment with Livestock manure. *Bacillus*

cereus and *Hypopichia burtonii* produced heavy turbidity (+++). *Aspergillus niger* had moderate turbidity (++) while *Penicillium expansum* and *Candida utilis* had minimal growth (+). This result corroborates the result of Chikere and Ekwuabu [12] in their study on molecular characterization of autochthonous hydrocarbon utilizing bacteria in oil-polluted sites at Bodo Community, Ogoni land, Niger Delta, Nigeria. The authors reported that *Aspergillus* and *Penicillium sp* produced heavy turbidity. Also, Yakubu and Bello [25], who worked on the biodegradation of Lagoma crude oil using pig dung, reported that *Bacillus sp* produced heavy growth, *Micrococcus* moderate growth while *Proteus* had minimal growth. .

5. CONCLUSION

The result obtained in this study has revealed that petroleum hydrocarbon-degrading bacteria and fungi isolated from the work support other people's findings. The isolates such as *Bacillus cereus*, *Bacillus subtilis*, and *Hypopichia butornii*, which are excellent hydrocarbon utilizers, can be explored for use as hydrocarbon degraders in bioremediation.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Agarry SE. Evaluation of the effects of inorganic and organic fertilizers and activated carbon on bioremediation of soil contaminated with weathered crude oil. *Journal of Applied Sciences and Environmental Management*, 2018;22(4): 587-595.
2. Schoonover Jon E, Jackie F. Crim. An Introduction to Soil Concepts and the Role of Soils in Watershed Management. *Journal of Contemporary Water Research & Education*. 2015;154(1):21-47.
3. Agarry SE, Oghenejoboh KM. Enhanced Aerobic Biodegradation of Naphthalene in Soil: Kinetic Modelling and Half-Life Study. *International Journal Environmental Bioremediation. and Biodegradation*. 2015; 3(2):48 – 53.
4. Kalev SD, Gurpal ST. "The Composition of Soils and Sediments." *Green Chemistry: An Inclusive Approach*. (November) 2018: 339-57.
5. Akarator SE, Molindo WA. Bioremediation Potential of Piggery Manure in a Soil Contaminated with Crude Oil in Benin City, Nigeria. *Journal of Agricultural Science and Technology*. 2016;5(1):33-41.
6. Bento FM, Camargo FAO, Okeke, BC, Frankenberger WT. Comparative bioremediation of soils contaminated with diesel oil by natural attenuation, biostimulation and bioaugmentation. *Bioresource Technology*. 2005;96:1049-1055.
7. Agarry SE, Olu-Arotiowa OA, Ajani AO, Aremu MO. Bioremediation of atrazine herbicide-contaminated soil using different bioremediation strategies. *Journal of Applied Sciences and Environmental Management*. 2019;23(1):99-109.
8. Chemlal R, Abdi N, Lounici H, Pauss A, Mameri N. Modelling and qualitative study of biodegradation using biopile process in sandy soil. *International Biodeterioration and Biodegradation*. 2013;78:43 – 48.
9. Agarry SE, Aremu MO, Aworanti OA Kinetic modelling and Half study on enhanced soil Bioremediation of Bonny light crude oil Amended with crop and animal-derived organic wastes. *Journals of petroleum environmental biotechnology*, 2013;4:137.
10. Ekpo FE, Nya EJ. Effect of poultry manure amendments on diesel oil polluted soil on germination and growth performance of some forest tree species. *Journal of Research in Environmental Science and Toxicology*. 2012;1(7):195-200.
11. Thouand G, Bauda P, Oudot J, Kirsch G, Sutton C, Vidallie JF. Laboratory evaluation of crude oil biodegradation with commercial or natural microbial inocula. *Canadian Journal of Microbiology*. 2016; 45(2):106-115.
12. Chikere CB, Ekwuabu CB. Molecular characterization of autochthonous hydrocarbon utilizing bacteria in oil-polluted sites at Bodo Community, Ogoni land, Niger Delta, Nigeria. *Nigerian Journal of Biotechnology*; 2014;27(1):23 - 45.
13. Onuorah S, Soludo OC, Odibo F. Impact of pig manure on the chemical characteristics of and Microbial population of crude oil - polluted soil in Awka, Nigeria. *American Journal of Life Science Researches*. 2018; 6(1):47-59
14. Onuorah S, Soludo OC, Odibo F. Effect of pig manure on the microbial remediation of

- crude oil polluted soil. American Journal of Life Science Researches. 2018;6(2):76-90.
15. Olukunle OF. Characterization of indigenous microorganisms associated with crude oil-polluted soils and water using traditional techniques. Microbiology Journal. 2013;3:1-11.
 16. Franco I, Contin M, Bragato G, De-Nobili, M. Microbiological resilience of soils contaminated with crude oil. Journal of Biotechnology. 2004;1(2):17-30.
 17. Chessbrough M. Laboratory manuals. District laboratory practice in Tropical Countries. Cambridge University Press, United Kingdom. 2006;145 – 157.
 18. Oyeleke SB, Manga SB. Essentials of laboratory Practical in Microbiology. First edition, Tobest Publishers, Minna, Nigeria. 2008;40-43.
 19. Ramaraj B. Isolation and characterization of a phenol-degrading sulphate-reducing bacterium from swine manure. Bioresources Technology. 1995;54: 29-33.
 20. Thijsee GJE, van der Linden AC. Iso-alkane oxidation by *Pseudomonas*. Antonie Van Leeuwenhoek. 2017;27: 171-179.
 21. Taiwo LB, Oso BA. Influence of composting techniques on microbial succession, temperature and pH in a composting municipal solid waste. African Journal of Biotechnology. 2004;3(4):239-243.
 22. Udebuani AC, Okoli CI, Nwigwe S, Harriet C, Ozoh PTE. The value of animal manure in the enhancement of bioremediation processes in petroleum hydrocarbon contaminated agricultural soils. Journal of Agricultural Technology, 2012;8(6):1935-1952.
 23. Chikere CB, Chikere BO, Okpokwasili GC. Bioreactor-based bioremediation of hydrocarbon-polluted Niger Delta marine sediment, Nigeria. Journal of Biotechnology. 2011;2(1):53 – 66.
 24. Omotayo AE, Efetie AO, Oyetibo G, Ilori MO, Amund OO. Degradation of aviation fuel by microorganisms isolated from tropical polluted soils. International Journal of Biological and Chemical Sciences, 2011;5(2):698-708.
 25. Yakubu M, Bello C. Biodegradation of Lagoma crude oil using pig dung. African Journal of Biotechnology. 2007;6(2):821-825.
 26. Stephen M, Panneerselvam A. Study of the prevailing fungi on the hydrocarbon-polluted soil. World Journal of Pharmaceutical Research, 2014;4(2): 1102-1108.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/107825>