



Effect of Land Configurations and Varieties on Yield and Economics of Summer Groundnut

S.S. Bharade ^{a++}, B.V. Asewar ^{b#} and B.M. Satale ^{a†*}

^a Department of Agronomy, VNMKV, Parbhani (M.S), India.

^b College of Agriculture, Golegaon, VNMKV, Parbhani (M.S), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i74787>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/118319>

Original Research Article

Received: 13/04/2024

Accepted: 15/06/2024

Published: 27/06/2024

ABSTRACT

The field investigation was conducted at experimental farm, Department of Agronomy, College of Agriculture, VNMKV, Parbhani, during summer season to find out results of yield and economic parameters. The experimental field was levelled and well drain. The soil was clay in texture, low in nitrogen, medium in phosphorous, medium in potash and alkaline in reaction. The experiment was laid out in a Split plot design with 12 treatment combinations, which comprised of four land configurations and three varieties. The main plots treatments were L₁- Ridges and furrows, L₂-BBF, L₃- Sara method and L₄- Flat beds and varieties were LGN-1, TAG-24, and SB-XI. Each experimental unit was replicated three times. Amongst land configuration BBF recorded significantly higher yield parameters viz., Dry pod yield plant⁻¹ (g), 100 kernel weight (g), kernel yield (kg ha⁻¹), Dry pod yield (kg ha⁻¹), haulm yield (kg ha⁻¹), biological yield (kg ha⁻¹), harvest index

⁺⁺ MSc. Agronomy;

[#] Associate Dean and Principal;

[†] Ph.D. Scholar;

^{*}Corresponding author: E-mail: satalebm92@gmail.com;

Cite as: Bharade, S.S., B.V. Asewar, and B.M. Satale. 2024. "Effect of Land Configurations and Varieties on Yield and Economics of Summer Groundnut". *International Journal of Plant & Soil Science* 36 (7):740-45. <https://doi.org/10.9734/ijpss/2024/v36i74787>.

(%) and in Economic parameter gross monetary returns (Rs ha⁻¹), net monetary returns (Rs ha⁻¹) and Benefit cost ratio (%) followed by Ridges and furrows. Among the varieties TAG-24 recorded highest values of yield and economic parameters as compared to the varieties SB-XI and LGN-1.

Keywords: Summer groundnut; commercial cultivation; nutrients; harvest index.

1. INTRODUCTION

“Groundnut, also called pea nut, earth nut, monkey nut, goober nut, manila nut, pinder and panda nut is a native of South American leguminous oil seed” [1]. “It was first found in Brazil or Peru as early as 950 B.C” [2]. According to Weiss [3], “a peanut was probably brought to West Africa from Brazil in the 16th century and then to the African East coast and to India”. “In India the plant was introduced by the Portuguese in 16th century as an oil seed crop for commercial cultivation. In the developed world, major pea nut produce is processed in to a variety of food products such as pea nut butter, salted pea nut, candies, salads, cheese and yoghurt like products, protein concentrate and pea nut meals” [4]. “Pea nut is fairly good source of some dietary minerals. The total minerals contains in pea nut ranges from 1.08 to 1.3 %. It contains (Mg per 100 g.) primary element such as Ca 50 to 90 mg, Mg 200 to 250 mg, P 340 to 430 mg, S 240 to 300 mg, K 500 to 730 mg and certain trace elements such as Mn 1.3 to 2 mg, Zn 2.3 to 4 mg, Boron 0.9 to 1.8 mg, Iron 2.3 to 3.2 mg. As groundnut helps to maintain blood cholesterol levels they have been recognized as heart friendly. Ground nuts provides over 30 essential nutrients are considered a rich source of fiber, vitamin (niacin, folate and vit. E) and minerals (Mg, Mn and P) and free from Na” [5]. The vitamins contents in pea nut have been studied by several authors [6] and reported that “pea nut have little or no vitamin A, D, K and B₁₂. Groundnut is a good source of all vitamins B except B₁₂”. “This is a rich source of thiamin, riboflavin, nicotinic acid and Vit E. The choline, inositol and pentathenic acid were present relatively higher amounts. India share 23 per cent of the world’s groundnut area and production. In India, it is grown in area of 4.77 M ha with total production 4.75 MT. Above 80 % of the area under groundnut is concentrated in five states viz. Gujarat, Andhra Pradesh, Tamil Nadu, Maharashtra and Karnataka. Gujarat occupies the first place in regard to area and production. The average yield of groundnut in India is 999 kg ha⁻¹, which is less than the average yield of groundnut in world has 1600 kg ha⁻¹. Maharashtra is one of the important groundnuts

growing states in India. The total area under groundnut in Maharashtra is 3.57 lacks ha out of which 2.57 in kharif and 0.87 summer with total production of 4.44 lack tonnes with average yield of 1150 kg ha⁻¹ (www.indiastat.com). “The botanical name of groundnut, *Arachis hypogaea* L. is derived from Greek word *Arachis* meaning a legume and *hypogaea* meaning below ground, referring to the formation of pods in the soil. Ground nut belongs to the genus *Arachis* of sub-family papilionaceae of the family Leguminaceae *Arachis hypogaea* L. ssp. procumbent are always trailing, runner or spreading type with a central axis which does not bear inflorescence and has produce lateral branches, plant is late maturing and prostrate habit” [7-9]. The plants of *Arachis hypogaea* L. sp. fastigiata are always bushy upright erect or bunchy type with inflorescence in the central axis and without a regular pattern in the sequence of reproductive and vegetative branches. Fruits are concentrated around the central axis and plants are early maturing.

2. MATERIALS AND METHODS

The field investigation was conducted at experimental farm, Department of Agronomy, College of Agriculture, VNMKV, Parbhani, during summer season to find out results of growth and quality parameters. The topography of experimental plot was fairly levelled. The soil was medium black in colour, and fairly well drained. The soil was clay in texture, low in nitrogen, medium in phosphorous, medium in potash and alkaline in reaction. “The experiment was laid out in a Split plot design with 12 treatment combinations, which comprised of four land configurations and three varieties. The experiment was laid out in split plot design where in the main plot were assigned to four land configuration (Flatbed, Ridges and furrow, BBF and Sara method) and subplots to three varieties of groundnut (LGN-1, TAG-24 and SB-XI) and the treatment combinations were randomly replicated thrice. The treatments were allotted randomly to each replication. The gross plot size was 5.4 m x 4.2 m and net plot size was 4.8 m x 3.8 m. The recommended dose of fertilizer was 25: 50: 00 kg NPK ha⁻¹ which is applied through Urea and D.A.P” [5].

3. RESULTS AND DISCUSSION

The mean dry pod yield per plant in (g) was significantly influenced by the various treatments. The land configuration BBF recorded significantly higher dry pod yield per plant (7.38 g) followed by the land configuration ridges and furrows (7.22 g). Shown in Table 1. Same result was reported by Hadvani et al. [10]. The data on dry pod yield per plant (g) revealed that the variety TAG-24 recorded significantly higher mean dry pod yield per plant (7.48 g) as compared to the other varieties SB-XI (6.96.g) and LGN-1 (6.74 g).

The effect of different land configuration on mean 100 kernel weight was found to be non-significant. But the highest 100 kernel weight was observed by the land configuration BBF (45.90 g). This might be due to higher soil moisture availability showed in Table 1. The data presented in Table 1 100 kernel weight (g) revealed that the variety TAG-24 recorded significantly higher mean 100 kernel weight (46.43 g) as compared to other varieties SB-XI (42.25g) and LGN-1 40.00 g).

The effect of different land configuration on mean kernel yield (kg ha^{-1}) was found to be significant. The land configuration BBF recorded significantly higher mean kernel yield (1341 kg ha^{-1}) followed by the land configuration ridges and furrows (1246 kg ha^{-1}). This might be due to higher soil moisture and also more surface area for peg penetration. showed in Table 1. The TAG-24 recorded significantly higher mean kernel yield (1375 kg ha^{-1}) as compared to the other varieties SB-XI and LGN-1. The varietal variation in yields might be due to differential genetic make-up. Similar result was reported by Ratankumar et al. [11] and Patra et al. [12].

Data presented in Table 1 on mean dry pod yield kg ha^{-1} as influenced by different land configuration was found to be significant. The land configuration BBF recorded significantly highest mean dry pod yield (1939 kg ha^{-1}) followed by the ridges and furrows (1835 kg ha^{-1}) land configuration with BBF 4% increased dry pod yield kg ha^{-1} over flat bed. This may be due higher dry pod weight, highest kernel easy harvesting and less pod retainment in soil. Same result was reported by Patel et al. [13], Patra et al. [12], Tirakannavar [14] and Bheemappa [15]. The land configuration BBF recorded significantly higher mean haulm yield (2756 kg ha^{-1}) followed by the ridges and furrows (2688 kg ha^{-1}) as

compared to all other land configurations. This may be due to all growth parameters are highest with BBF. Same result was reported by Baskaran et al. [16], Patil et al. [17]. The land configuration BBF recorded significantly higher mean biological yield (4729 kg ha^{-1}) followed by the land configuration ridges and furrows (4381 kg ha^{-1}) as compared to the all other land configurations shown in Table 1. Same result was reported by Baskaran et al. [16].

The data in Table 1 on dry pod yield kg ha^{-1} , haulm yield kg ha^{-1} , biological yield kg ha^{-1} and kernel yield kg ha^{-1} revealed that the genotype TAG-24 recorded significantly higher mean dry pod yield (1999 kg ha^{-1}), haulm yield (2820 kg ha^{-1}), biological yield (4820 kg ha^{-1}) and kernel yield (1375 kg ha^{-1}) as compared to the other varieties SB-XI and LGN-1. The varietal variation in yields might be due to differential genetic make-up. Similar result was reported by Ratankumar et al. [11] and Patra et al [12].

Data presented in Table 1 on harvest index showed that there was no any significant effect by the different land configurations on harvest index. But the highest harvest index was observed (41.48 %) by the land configuration ridges and furrows as compared to other land configurations shows in Table 1. The harvest index it was revealed that the response of genotypes on harvest index was found to be non significant. But the highest harvest index was observed in the variety TAG-24 (41.43%) as compared to the other varieties SB-XI (39.81%) and LGN-1 (39.11%).

The presented in Table 2 data on gross monetary returns revealed that the land configuration with BBF recorded the maximum gross returns ha^{-1} (84469 Rs ha^{-1}) followed by ridges and furrows (80125 Rs ha^{-1}) as compared to the other land configurations. This may be due to higher pod yield and haulm with BBF. Similar result was recorded by Tarde [18] and Baskaran et al. [16]. The net monetary returns per hectore revealed that the land configuration BBF recorded highest net monetary returns per hectore (59188 Rs ha^{-1}) followed by the land configuration ridges and furrows (54145 Rs ha^{-1}). Similar result was recorded by Tarde [18] and Bheemappa et al. [15]. The data presented in Table 2 on benefit:cost ratio it was seen that the land configuration BBF recorded highest benefit:cost ratio (3.34) followed by the land configuration ridges and furrows (3.09) and land configuration Sara (3.03). The land configuration

flat beds recorded the lowest benefit:cost ratio (3.00).

The gross monetary returns Rs ha⁻¹ and net monetary returns Rs ha⁻¹ revealed that the variety TAG-24 recorded significantly higher gross monetary returns (87047 Rs ha⁻¹) and net monetary returns (62325 Rs ha⁻¹) as compared

to the other varieties SB-XI and LGN-1. This may be due to higher dry pod yield and haulm yield. Similar result was reported by Chandapur [19] and Meerasab et al. [20]. The benefit: cost ratio it was revealed that the variety TAG-24 recorded higher benefit: cost ratio (3.51) as compared to the varieties SB-XI (3.12) and LGN-1 (2.70) [21].

Table 1. Dry pod yield plant⁻¹ (g), 100 kernel weight (g), and kernel yield (kg ha⁻¹), Dry pod yield (kg ha⁻¹), haulm yield (kg ha⁻¹), biological yield (kg ha⁻¹) and harvest index (%) of groundnut as influenced by various treatments

Treatments	Dry pod Yield plant ¹ (g)	100 Kernel Weight (g)	Kernel yield (Kg ha ⁻¹)	Dry pod Yield (Kg ha ⁻¹)	Haulm yield (Kg ha ⁻¹)	Biological yield (Kg ha ⁻¹)	Harvest Index (%)
L) Land Configuration (L)							
L ₁ - Ridges and Furrows	7.22	43.79	1246	1835	2688	4523	40.53
L ₂ – BBF	7.38	45.90	1341	1939	2756	4729	41.48
L ₃ - Sara method	6.84	42.78	1015	1642	2480	4089	40.0
L ₄ - Flat bed	6.70	41.60	947	1553	2396	3950	39.11
SE ±	0.047	0.22	12.26	26.16	23.02	39.62	-
CD at 5 %	0.16	NS	42.44	90.55	79.68	137.12	-
V) Varieties (V)							
V ₁ - TAG-24	7.40	46.43	1375	1999	2820	4820	41.43
V ₂ - SB-XI	6.96	42.25	1078	1678	2516	4195	39.81
V ₃ - LGN-1	6.74	40.91	958	1549	2405	3954	39.11
SE ±	0.045	0.18	13.29	13.75	19.82	40.93	-
CD at 5 %	0.13	0.54	39.84	41.23	59.44	122.71	-
Interaction (L x V)							
SE ±	0.18	0.73	53.16	55.94	79.13	163.71	-
CD at 5%	NS	NS	NS	NS	NS	NS	-
General Mean	7.03	43.2	1137	1742	2580	4323	-

Table 2. Economics of the groundnut cultivation as influenced by various treatments

Treatments	Gross Monetary Returns (Rs ha ⁻¹)	Net monetary Returns (Rs ha ⁻¹)	Benefit cost Ratio
L) Land Configuration (L)			
L ₁ - Ridges and Furrows	80125	25480	54645
L ₂ – BBF	84469	25280	59188
L ₃ - Sara method	71885	23780	48105
L ₄ - Flat bed	68134	22780	45464
SE ±	1065	-	788
CD at 5 %	3686	-	2729
V) Varieties (V)			
V ₁ - TAG-24	87047	24597	62450
V ₂ - SB-XI	73437	23397	50040
V ₃ - LGN-1	67976	24997	43062
SE ±	602	-	466
CD at 5 %	805	-	1397
Interaction (L x V)			
SE ±	2409	-	1865
CD at 5%	NS	-	NS
General Mean	76153	24330	51851

The data presented in Table 2. on dry pod yield per plant (g), dry pod yield kg ha⁻¹, 100 kernel weight (g), haulm yield kg ha⁻¹, biological yield kg ha⁻¹, kernel yield kg ha⁻¹ and harvest index revealed that the interaction effect of land configurations and genotypes on above yield and yield attributes was found to be non-significant. The data presented in Table 2 on gross monetary returns Rs ha⁻¹ and net monetary returns Rs ha⁻¹ revealed that the interaction effect of land configurations and varieties on gross monetary returns and net monetary returns was found to be non-significant [22,23].

4. CONCLUSION

Based on year experimentation, it is concluded that, Land configuration with BBF produced superior growth and dry pod yield per plant (g), dry pod yield kg ha⁻¹, 100 kernel weight (g), haulm yield kg ha⁻¹, biological yield kg ha⁻¹, kernel yield kg ha⁻¹ and harvest index and it was comparable with the ridges and furrows. The variety TAG-24 was found superior in growth and development and dry pod yield per plant (g), dry pod yield kg ha⁻¹, 100 kernel weight (g), haulm yield kg ha⁻¹, biological yield kg ha⁻¹, kernel yield kg ha⁻¹ and harvest index as compared to the variety SB-XI and LGN-1. The variety TAG-24 was found to be productive and remunerative. Highest GMR and NMR were recorded with BBF and it was comparable with other land configurations and among the varieties TAG-24 recorded highest GMR and NMR.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Hammons RO. Origin and early history of the pea nut. In pea nut science and Technology, ed. Pattee HE, young CT. 1982;1-20.
2. Higgins BB. Origin and early history of the pea nut. The pea nut the Unpredictable Irgume. Washington:National Fertilizer Association; 1951.
3. Weiss EA. Oilseed crops, Tropical agriculture series, London: Longman; 1983.
4. Chavan UD, JK Chavan, MB Pinjari, SS Kadam. Nutritional composition and oil quality of promising groundnut cultivars. *J. Mah. Agril. Univ.* 1990;16:48-50.
5. Bharade SS, Asewar BV, Bobade BR, Mirza IA. Effect of land configurations and varieties on growth and quality of summer groundnut. *Int. J. Curr. Microbiol. Appl. Sci.* 2019;8:1392-398.
6. Rao PV, Rao SP. Chemical composition of fatty acid profile of high yielding varieties of oil seeds. *Indian J. of Agric. Science.* 1981; 51:703-707.
7. Malviya Shreya, Harish Chandra Patel. Influence of land configuration, seed rates, and variety on physiology and yield of rainfed soybean in semi-arid tropics of Central India. *Journal of Experimental Agriculture International.* 2023;45(9):26-33. Available:<https://doi.org/10.9734/jeai/2023/v45i92172>.
8. Bhatt Kanika, Nilutpal Saikia, Jitendra Singh Bohra, Kadagonda Nithinkumar, Shreyas Bagrecha, Tejbal Singh, Harsita Nayak, Aakash. Integrated nutrient management (INM) and varietal impact on toria (*Brassica campestris* L. Var. Toria) growth, yield, and economics. *Asian Journal of Soil Science and Plant Nutrition.* 2024;10(1):23-29. Available:<https://doi.org/10.9734/ajsspn/2024/v10i1207>.
9. Das A, Layek J, Ramkrushna GI, Patel DP, Choudhury BU, Chowdhury S, Ngachan SV. Raised and sunken bed land configuration for crop diversification and crop and water productivity enhancement in rice paddies of the north eastern region of India. *Paddy and Water Environment.* 2015, Oct;13:571-80.
10. Hadvani RG, Ahlawat IPS, Trivedy J. Effect of method of sowing and levels of sulphur on growth and yield of groundnut. *Indian Journal of Agronomy.* 1993;38(2): 325-327.
11. Ratankumar AU, Hariprasanna K, Lalwni HB. Genetic improvement in Spanish type groundnut, *Arachis hypogaea* L.varieties in India over the years. *Journal of Oilseed Research.* 2010;27(1):1-7.
12. Patra AK, Samui RC, Tripathy SK. Response of summer groundnut varieties to potassium and planting methods.

- Journal of Oilseed Research. 1996;13(1): 26-31.
13. Patil SH, Kale DM, Deshmukh SN, Fulzele GR, Weginwar BG. Semi dwarf, early maturing and high yielding new groundnut variety variety, TAG-24. Journal of Oilseed Research. 1995;12(2): 254-257.
 14. Tirakannavar AV. Response of soybean to planting methods plant density and no. of seeds per hill under rainfed condition in Vertisol. M.Sc. Agri. Thesis, University of Agriculture Science, Dharwad; 1999.
 15. Bheemappa A, Meti SK, hanchinal SN. Effectiveness of broad bed and furrow method of groundnut cultivation as perceived by farmers. Karnataka J. Agric. sci. 1994;7(2):205-210.
 16. Baskaran R, Solaimalai A, Subbaramu K. Effect of water harvesting techniques and IPM practices on productivity of a rainfed groundnut. Crop Research. 2003;26(3): 424-428.
 17. Patil HM, Kolekar PT, Shete BT. Effect of layouts and spacing on yield and quality of bold seeded summer groundnut. International Journal of Agric. Sci. 2007; 3(2):210-213.
 18. Tarde NR. Studies on growth, yield and quality performance of bunch type groundnut (*Arachis hypogea*) variety FSB-7-2) as influenced by planting layout between row spacing and plant population under summer condition. M.Sc. Agri. Thesis, MPKV, Rahuri; 1984.
 19. Chandapur VB. Studies on growth and yield of groundnut varieties under different land configuration. M.Sc. (Agri.) Thesis, VNMKV, Parbhani; 2012.
 20. Meerasab HJ, Guled MB. Effect of sara method of in situ moisture conservation practices and intercropping system on yield of rabi sorghum and chickpea. Karnataka J. Agric. Sci. 2013;26(3):356-359.
 21. Desai MM. Effect of planting layout and gypsum level on the growth, yield and quality of groundnut variety Phule Pragati during Kharif. M.Sc. Agri. Thesis, MPKV, Rahuri; 1989.
 22. Jadhav AP. Response of gram Phule-5 to irrigation applied at physiological growth stages, layouts and plant densities. M.Sc. Agri. Thesis. MPKV, Rahuri; 1983.
 23. Rasve SD, Bharambe PR, Ghonsikar GP. Effect irrigation frequency and method of cultivation on yield and quality of summer groundnut. J. Maharashtra Agric. Univ. 1983;8(1):57-59.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/118319>