



Effect of Integrated Use of Manures and Fertilizers on Physico-chemical Properties of Soil and Yield of Coconut (*Cocos nucifera* L.)

S. S. Pawar ^{a++*}, R.T. Bankar ^{a++} and N.H. Khobragade ^{a#}

^a Department of Soil Science and Agril. Chemistry, Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli- 415 712, Dist. Ratnagiri (M.S.), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i74803>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/118962>

Original Research Article

Received: 25/04/2024
Accepted: 29/06/2024
Published: 02/07/2024

ABSTRACT

A field experiment was conducted during 2014-15 and 2015-16 in a thirty years old coconut garden of cultivar *Pratap* to study the effect of Integrated Use of Manures and Fertilizers on Physico-chemical Properties of Soil and Yield of Coconut (*Cocos nucifera* L.). Accordingly, treatments were applied to coconut orchard in three splits (*viz.*, Stage I- June, Stage II- October and Stage III- February) in a year. Soil samples were collected periodically before fertilizers application and analysed for different physico-chemical properties (*viz.*, pH, EC and organic carbon). The yield of each palm was recorded throughout the year under experimental plot. The application of RDN

⁺⁺ Ph.D. Scholar;

[#] Assistant Professor;

^{*}Corresponding author: E-mail: pawar_667@rediffmail.com;

Cite as: Pawar, S. S., R.T. Bankar, and N.H. Khobragade. 2024. "Effect of Integrated Use of Manures and Fertilizers on Physico-Chemical Properties of Soil and Yield of Coconut (*Cocos Nucifera* L.)". *International Journal of Plant & Soil Science* 36 (7):908-19. <https://doi.org/10.9734/ijpss/2024/v36i74803>.

through FYM at 25 kg/palm/year + Neem Cake at 15 kg/palm/year + Vermicompost at 6 kg/palm/year (T₁₀) recorded significant increase in pH, EC and organic carbon content in soil of coconut orchard. Also the physico-chemical properties of soil found to be improved during second year over the first year of experimentation. The application of RDF + Azadirachtin + Micronutrients through briquettes (T₅) received highest yield of coconut. Therefore, it is concluded that application of RDF along with Azadirachtin and Micronutrients through Konkani Annapurna Briquettes in three splits (i.e. June, October and February) is beneficial for increasing the yield of coconut.

Keywords: INM; coconut; soils; physico-chemical properties; yield; Maharashtra.

1. INTRODUCTION

The coconut (*Cocos nucifera* L.) palm eulogized as “Kalpavriksha” and “Tree of Heaven”, provides not only edible parts but also fuel, shelter, medicine and employment to the millions of people in tropics. In India, coconut is grown in 19 states and 3 union territories, mostly along the coastal region of the country. The area under coconut in Ratnagiri district (M.S.) is 4,549 hectares with an annual production of 409.41 lakh nuts and average productivity of 9,001 nuts ha⁻¹ [1]. The coconut palm requires balanced fertilization of NPK; the response of N and K was common while response of P was under certain conditions. Palms with larger yield potential need higher dose of NPK. Worldwide people realized that pure chemical farming undermines the natural mechanism operating in the ecosystem and often leads to soil degradation, pollution of ground water and eutrophication of water bodies with nitrates, phosphates and pesticides Upadhyay et al. [2], [3-6]. Organic manures are important in sustaining soil productivity especially for a perennial crop like coconut, which requires continuous supply of nutrients.

Therefore, integrated nutrient management been always considered as the best option to increase the crop yield. In this regard, present study was conducted on Effect of Integrated Use of Manures and Fertilizers on Physico-chemical Properties of Soil and Yield of Coconut.

2. MATERIALS AND METHODS

The experiment was conducted during 2014-15 and 2015-16 in a thirty years old coconut garden of cultivar Pratap at the farm of Asond block,

Central Experiment Station, Wakawali, Dr. Balasaheb Sawant Konkani Krishi Vidyapeeth, Dapoli, Tal.- Dapoli, Dist.- Ratnagiri (M.S.) which is located at 17.68337° N latitude and 73.27761° E longitudes with an altitude of 250 m MSL (Fig. 1). The research station receives very high annual average rainfall of 3500 mm in rainy days. The soils of experimental site are well drained, sandy clay loam in texture, slightly acidic in reaction with low electrical conductivity and high in organic carbon content. The soil is medium in available nitrogen, low in available phosphorus and high in available potassium content. Kaolinite is the dominant clay mineral in this soil.

The experiment was laid out in Randomized Block Design comprising ten treatments (*viz.*, T₁- Absolute Control (No manure, no fertilizer), T₂- Recommended Dose of Fertilizers (RDF) only, T₃- Application of RDF through briquettes, T₄- Application of RDF + Azadirachtin through briquettes, T₅- Application of RDF + Azadirachtin + Micronutrients through briquettes, T₆- Application of RDF + Neem oil through briquettes, T₇- Application of RDF through briquettes and neem cake at 15 kg/palm, T₈- Application of RDF + Root feeding with Azadirachtin 5 % @ 7.5 mL + 7.5 mL water, T₉- Application of RDF + Drenching with Eriophyid smash 250 mL/ 20 L of water and T₁₀- RDN through FYM at 25 kg/palm/year + Neem cake at 15 kg/palm/year + Vermicompost at 6 kg/palm/year) replicated thrice. Two palms for each treatment were selected at Farm of Asond Block, C.E.S. Wakawali, Dr. B. S. Konkani Krishi Vidyapeeth, Dapoli, Dist.- Ratnagiri during 2014-15 and 2015-16. The treatments were applied in three splits (*viz.*, Stage I- June, Stage II- October and Stage III- February) in a year.





Fig. 1. Location map of study area



Picture 1. Formulation of briquettes using different ingredients for different treatments



Picture 2. Application of manures and fertilizers in the fertilizer ring



Picture 3. Root feeding with Azadirachtin 5% @ 7.5 mL +7.5 mL water



Picture 4. Harvesting of coconut

Soil samples were collected periodically before fertilizers application and analyzed for different physico-chemical properties (*viz.*, pH, EC and organic carbon). The pH and EC in soil was determined by potentiometrically and conductometrically as per the procedure given by Jackson [7] while, organic carbon was determined by Walkley and Black's wet digestion method as described by Black [8]. The yield of each palm was recorded throughout the year under experimental plot. The year wise yield was worked out and analyzed statistically for Randomized Block Design by following the standard procedures given by Panse and Sukhatme [9].

3. RESULTS AND DISCUSSION

The results of study presented in Table 1, it is clearly observed that the physico-chemical

properties of soil in coconut orchard were improved due to application of manures and fertilizers during both the years of experimentation. The treatments receiving application of RDN through FYM at 25 kg/palm/year + Neem Cake at 15 kg/palm/year + Vermicompost at 6 kg/palm/year (T_{10}) and application of RDF through briquettes and neem cake at 15 kg/palm (T_7) showed the maximum improvement in pH of soil. An overall the mean soil pH during first year and second year ranged between 5.46 to 5.89 and 5.43 to 6.04, respectively. In comparison of both years an increasing trend of soil pH was noticed from first year to second year due to application of organic manures in treatments T_{10} and T_7 . In general, due to application of organic manure like FYM, neem cake and vermicompost soil pH was found to be increased over rest of the treatments including recommended dose of fertilizers (T_2)

during both years of the experiment. The soil samples in all treatments were 'moderately to slightly acidic' in reaction as per the ratings given by Bangar and Zende [10]. The increase in pH of acid soil due to addition of organic manures is attributed to the deactivation of Fe^{3+} and concomitant release of basic cations during their decomposition [11]. Similar results were also recorded by Temgire [12] and Maheswarappa et al. [13] in lateritic soils of coconut orchard due to application of manures and fertilizers.

In case of electrical conductivity (Table 2), it did not much more influenced due to application of manure and fertilizers through different treatments applied during both the years of experiment. The electrical conductivity of soil during first year and second year varied between 0.062 to 0.102 dS m^{-1} and 0.073 to 0.118 dS m^{-1} , respectively. There is slight increase in electrical conductivity of soil due to application of all treatments of manures and fertilizers over absolute control. Comparatively, the electrical conductivity of soil during second year was increased over first year. The electrical conductivity of soil in all treatments was low and had no any potential threat to the productivity of the soils to crop growth. It was also observed that the electrical conductivity of soil was significantly increased due application of manures and fertilizers over absolute control during both years. In general, the soil samples under all treatments were found under 'normal' (i.e. $< 1.0 \text{ dS m}^{-1}$) class of electrical conductivity based on the ratings given by Seth [14] indicating low salt concentration in these soils. Increase in electrical conductivity with application of organic manures and fertilizers were also reported by Diwale [15] and Bhosale [16] for lateritic soils of Konkan.

The data given in Table 3, clearly explain that application of manures and fertilizers significantly increased organic carbon content of soil in all stages of both the years of experimentation. The organic carbon content in soil during first year and second year was ranged between 1.57 to 2.36 per cent and 1.47 to 2.73 per cent, respectively. The maximum built up of soil organic carbon in soil was noticed due to application of treatments receiving application of RDN through FYM at 25 kg/palm/year + Neem Cake at 15 kg/palm/year + Vermicompost at 6 kg/palm/year (T_{10}) and application of RDF through briquettes and neem cake at 15 kg/palm (T_7) and during both the years of trial. In

comparison of both years, an increasing trend of soil organic carbon was noticed from first year to second year due to application of organic manures in treatments T_{10} and T_7 .

The soil samples in all treatments were categorized under 'very high' class due to the presence of excess amount of organic carbon content in the soils of coconut orchards Bangar and Zende [10].

The improvement in organic carbon content of soil in organic manure treated plots might be ascribed to direct addition of organic matter through organic manures and due to addition of considerable amount of leaf litter of crops. The High amount of organic carbon content in these soils might be attributed to luxurious growth of grasses and vegetation due to high rainfall and thus addition of organic matter through litter, residues and cover crops and thereby subsequent increased humification Preethi et al. [17]. An increase in organic carbon content of soil due to the application of various organic manures were noted by Hapse et al. [18], Kadam [19], Yaduvanshi [20], Gedam et al. [21], Gadade [22], Yadav et al. [23] and Diwale [15] for different soils.

Effect on Yield: The data given in Table 4 and depicted in Fig. 2, clearly explain that the yield of coconut significantly increased due to application of manures and fertilizers in both the years of trial i.e. 2014-15 and 2015-16. The application of RDF + Azadirachtin + Micronutrient through briquettes (T_5) recorded significantly higher nut yield (125.33 and 129.50 nuts palm⁻¹ year⁻¹) over rest of the treatments during both years of experimentation, respectively except treatments T_8 and T_9 . Treatment T_5 was at par with treatments T_8 and T_9 . The lowest yield was recorded by the treatment T_1 (Absolute control).

An overall the nut yield of coconut during first year and second year ranged between 85.33 to 125.33 nuts palm⁻¹ year⁻¹ and 84.67 to 129.50 nuts palm⁻¹ year⁻¹, respectively. In comparison of both years an increasing trend of nut yield from first year to second year was noticed due to application of manures and fertilizers. The increase in nut yield during second year might be due to combined effect of manures and fertilizers applied during second year as well as the residual impact of manures and fertilizers applied during first year.

Table 1. Effect of integrated use of manures and fertilizers on pH of soils in coconut orchard

Treat. No.	Treatments	2014-15				2015-16			
		Stage I	Stage II	Stage III	Mean	Stage I	Stage II	Stage III	Mean
T ₁	Absolute Control (No manure, no fertilizer)	5.58	5.50	5.44	5.51	5.44	5.50	5.53	5.49
T ₂	Recommended Dose of Fertilizers (RDF) only	5.52	5.44	5.43	5.46	5.47	5.40	5.44	5.44
T ₃	Application of RDF through briquettes	5.79	5.70	5.49	5.66	5.47	5.41	5.42	5.43
T ₄	Application of RDF + Azadirachtin through briquettes	5.59	5.59	5.51	5.56	5.48	5.54	5.53	5.52
T ₅	Application of RDF + Azadirachtin + Micronutrient (B) through briquettes	5.56	5.54	5.36	5.49	5.54	5.74	5.46	5.58
T ₆	Application of RDF + Neem oil through briquettes	5.50	5.60	5.46	5.52	5.37	5.47	5.45	5.43
T ₇	Application of RDF through briquettes and neem cake at 15 kg/palm	5.64	5.84	6.00	5.83	6.02	5.76	5.85	5.88
T ₈	Application of RDF + Root feeding with Azadirachtin 5% @ 7.5 mL + 7.5 mL water	5.49	5.69	5.50	5.56	5.48	5.54	5.50	5.51
T ₉	Application of RDF + Drenching with Eriophyid smash 250 mL/20 L of water	5.73	5.60	5.49	5.61	5.49	5.52	5.53	5.51
T ₁₀	RDN through FYM at 25 kg/palm/year + Neem cake at 15 kg/palm/year + Vermicompost at 6 kg/palm/year	5.54	6.05	6.07	5.89	6.12	5.98	6.01	6.04
	SE ±	0.11	0.05	0.06		0.05	0.06	0.02	
	CD @ 5 %	N.S.	0.15	0.19		0.14	0.18	0.07	

Note: Stage-I of first year is the initial stage and the values given under Stage-I of first year are irrespective of treatments

Table 2. Effect of integrated use of manures and fertilizers on electrical conductivity (dS m⁻¹) of soil in coconut orchard

Treat. No.	Treatments	2014-15				2015-16			
		Stage I	Stage II	Stage III	Mean	Stage I	Stage II	Stage III	Mean
T ₁	Absolute Control (No manure, no fertilizer)	0.065	0.060	0.061	0.062	0.078	0.072	0.068	0.073
T ₂	Recommended Dose of Fertilizers (RDF) only	0.060	0.059	0.079	0.066	0.088	0.079	0.070	0.079
T ₃	Application of RDF through briquettes	0.088	0.071	0.106	0.088	0.124	0.088	0.076	0.096
T ₄	Application of RDF + Azadirachtin through briquettes	0.078	0.066	0.126	0.090	0.140	0.103	0.080	0.108
T ₅	Application of RDF + Azadirachtin + Micronutrient (B) through briquettes	0.082	0.092	0.099	0.091	0.088	0.100	0.111	0.100
T ₆	Application of RDF + Neem oil through briquettes	0.082	0.086	0.100	0.089	0.111	0.117	0.086	0.105
T ₇	Application of RDF through briquettes and neem cake at 15 kg/palm	0.081	0.086	0.096	0.088	0.139	0.118	0.084	0.114
T ₈	Application of RDF + Root feeding with Azadirachtin 5% @ 7.5 mL + 7.5 mL water	0.096	0.096	0.115	0.102	0.109	0.117	0.095	0.107
T ₉	Application of RDF + Drenching with Eriophyid smash 250 mL/20 L of water	0.091	0.093	0.110	0.098	0.135	0.112	0.107	0.118
T ₁₀	RDN through FYM at 25 kg/palm/year + Neem cake at 15 kg/palm/year + Vermicompost at 6 kg/palm/year	0.090	0.081	0.089	0.087	0.116	0.108	0.121	0.115
	SE ±	0.007	0.004	0.007		0.009	0.007	0.005	
	CD @ 5 %	N.S.	0.011	0.020		0.027	0.021	0.016	

Note: Stage-I of first year is the initial stage and the values given under Stage-I of first year are irrespective of treatments

Table 3. Effect of integrated use of manures and fertilizers on organic carbon (%) of soil in coconut orchard

Treat. No.	Treatments	2014-15				2015-16			
		Stage I	Stage II	Stage III	Mean	Stage I	Stage II	Stage III	Mean
T ₁	Absolute Control (No manure, no fertilizer)	1.51	1.49	1.70	1.57	1.51	1.43	1.46	1.47
T ₂	Recommended Dose of Fertilizers (RDF) only	1.55	1.50	1.66	1.57	1.74	1.57	1.34	1.55
T ₃	Application of RDF through briquettes	1.64	1.94	1.63	1.74	1.78	1.66	1.68	1.71
T ₄	Application of RDF + Azadirachtin through briquettes	1.55	1.77	1.74	1.69	1.72	1.65	1.62	1.66
T ₅	Application of RDF + Azadirachtin + Micronutrient (B) through briquettes	1.63	1.77	1.61	1.67	1.25	1.83	1.75	1.61
T ₆	Application of RDF + Neem oil through briquettes	1.51	1.78	1.49	1.59	1.48	1.51	1.57	1.52
T ₇	Application of RDF through briquettes and neem cake at 15 kg/palm	1.69	2.15	2.07	1.97	2.12	2.22	2.29	2.21
T ₈	Application of RDF + Root feeding with Azadirachtin 5% @ 7.5 mL + 7.5 mL water	1.59	1.52	1.81	1.64	1.55	1.72	1.54	1.60
T ₉	Application of RDF + Drenching with Eriophyid smash 250 mL/20 L of water	1.77	1.40	1.76	1.64	1.62	1.75	1.59	1.65
T ₁₀	RDN through FYM at 25 kg/palm/year + Neem cake at 15 kg/palm/year + Vermicompost at 6 kg/palm/year	1.85	2.54	2.68	2.36	2.59	2.70	2.89	2.73
	SE ±	0.09	0.09	0.08		0.12	0.09	0.06	
	CD @ 5 %	N.S.	0.26	0.24		0.36	0.26	0.18	

Note: Stage-I of first year is the initial stage and the values given under Stage-I of first year are irrespective of treatments

Table 4. Effect of integrated use of manures and fertilizers on yield (nuts palm⁻¹ year⁻¹) of coconut

Treat. No.	Treatments	2014-15	2015-16
T ₁	Absolute Control (No manure, no fertilizer)	85.33	84.67
T ₂	Recommended Dose of Fertilizers (RDF) only	98.67	104.50
T ₃	Application of RDF through briquettes	102.17	114.17
T ₄	Application of RDF + Azadirachtin through briquettes	112.50	117.67
T ₅	Application of RDF + Azadirachtin + Micronutrient (B) through briquettes	125.33	129.50
T ₆	Application of RDF + Neem oil through briquettes	103.33	113.00
T ₇	Application of RDF through briquettes and neem cake at 15 kg/palm	114.83	116.67
T ₈	Application of RDF + Root feeding with Azadirachtin 5 % @ 7.5 mL + 7.5 mL water	122.67	126.17
T ₉	Application of RDF + Drenching with Eriophyid smash 250 mL/ 20 L of water	123.83	127.17
T ₁₀	RDN through FYM at 25 kg/palm/year + Neem cake at 15 kg/palm/year + Vermicompost at 6 kg/palm/year	104.00	101.17
	SE ±	3.43	3.31
	CD @ 5 %	10.19	9.84




Fig. 2. Effect of Integrated use of manures and Fertilizers on nut yield of coconut orchard during 2014-15 and 2015-16

In general, due to application of manure and fertilizers through different treatments the nut yield in coconut orchard was found to be increased over absolute control during both years of the experiment. The increase in nut yield with integration of organics with fertilizers was attributed to increased female flowers and nut setting per cent due to improved availability of nutrients to coconut.

The beneficial response of manures and fertilizers over absolute control to yield might be attributed to the availability of sufficient amount of plant nutrients throughout the year to the crop, improvement of soil environment resulting in higher root proliferation leading to better absorption of moisture and nutrients, plant vigour and ultimately higher yield. After proper decomposition and mineralization, the manures supplied available nutrients directly to the plants and also had solubilising effect on fixed forms of nutrients in soil Sinha et al. [24].

Similar results of higher nut yield with integrated use of manures and fertilizers were also reported by Temgire [12] and Talashilkar et al. [25].

4. CONCLUSION

Based on data obtained from the present investigation, it could be concluded that the application of RDN through FYM at 25 kg/palm/year + Neem Cake at 15 kg/palm/year + Vermicompost at 6 kg/palm/year in three splits (i.e. June, October and February) is beneficial for improving physico-chemical properties of soil of

coconut orchard. The application of RDF + Azadirachtin + Micronutrients through briquettes (T₅) receiving highest yield. Therefore, it is concluded that the application of RDF along with Azadirachtin and Micronutrients through Konkan Annapurna Briquettes in three splits (i.e. June, October and February) is beneficial for increasing the yield of coconut in lateritic soils of Konkan.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (Chat GPT, COPILOT, etc.) and text-to-image generators have been used during writing or editing of manuscripts.

ACKNOWLEDGEMENT

The authors express his heartfelt gratitude to Dr. S. S. Prabhudesai, Head, Department of Soil science and Agril. Chemistry, Dr. V. G. Salvi, Associate professor Department of Soil science and Agril. Chemistry, Dr. N.H. Khobragade, Assistant Professor, Department of Soil science and Agril. Chemistry, Dr. J. S. Dhekale, Associate professor, Department of Agril. Economics, Dr. V. S. Desai, Manager, Agricultural Technology Information Centre and Dr. P. C. Haldavnekar, Associate Director of Research, RFRS, Vengurla, Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth Dapoli for their excellent guidance, suggestions and regular encouragement during the course of investigation.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Anonymous; 2015.
DOI:www.coconutboard.gov.in/presentation/statistics/statistics.aspx..
2. Upadhyay AK, Srinivasa Reddy DV, Biddappa CC. Organic farming technology for coconut, Indian Coconut Journal. 1998; 24:74-78.
3. Diwedi, Namrata, Sudhakar Singh, Deepak Pandey, Singh PK, Chanda SS, Tiwari HN, Singh DK, Gajendra Singh. The effect of integrated nutrient management on growth, yield attributes and yield of transplanted rice under irrigated condition (*Oryza sativa* L.). Journal of Experimental Agriculture International. 2024;46(1):27-36.
Available:https://doi.org/10.9734/jeai/2024/v46i12288.
4. Kumar Amit, Singh SK, Rohtas Kumar, Sandip Kumar Gautam, Amit Sharma. Evaluating the effects of organic manures and fertilizer N on soil properties in rice cultivation. International Journal of Environment and Climate Change. 2022;12(12):1140-48.
Available:https://doi.org/10.9734/ijecc/2022/v12i121552.
5. Kumar D, Rakshit R, Rani R, Bharti P, Das A, Kundu M. Integrated nutrient management in coconut (*Cocos nucifera* L.): An assessment of soil chemical and biological parameters under subtropical humid climate. Journal of Soil Science and Plant Nutrition. 2022;22(2):2695-2706.
6. Andoh-Mensah E, Anthonio CK, Sossah FL, Fianko DA, Yankey EN. Integrated soil fertility management using cocoa bean shells improves soil chemical properties, coconut yield and mitigates environmental pollution. Journal of Cleaner Production. 2023;428:139418.
7. Jackson ML. Soil Chemical Analysis, Prentice Hall of India Pvt. Ltd., New Delhi. 1973;134-182.
8. Black CA. Methods of Soil Analysis, Part-I, American Society for Agronomy, Inc., Madison, Wisconsin, USA; 1965.
9. Panse VG, Sukhatme PV. Statistical methods for Agricultural Workers, Indian Council of Agricultural Research Publication, New Delhi; 1967.
10. Bangar AR, Zende GK. Soil testing: A new basis for efficient fertilizer use. Journal of Maharashtra Agriculture University. 1978; 3(2):81-84.
11. Lal S, Mathur BS. Effect of long term manuring fertilization and liming on crop yield and some physical properties of acid soil. Journal of the Indian Society of Soil Science. 1988;36:113-119.
12. Temgire AM. Effect of integrated use of manures and fertilizers on yield, nutrient uptake and quality of coconut (*Cocos nucifera* L.) and soil properties, M.Sc. (Agri.) Thesis submitted to Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli, Dist. Ratnagiri, Maharashtra; 2007.
13. Maheswarappa HP, George VT, Gupta A, Bhat R and Palaniswami C. Productivity and nutrient status of coconut (*Cocos nucifera*) as influenced by integrated nutrient management with vermicomposted coconut leaves. Indian Journal of Agronomy. 2014;59(3):455-459.
14. Seth SP. Indices for diagnosis and salinity in soils of Rajasthan canal area; 1967.
15. Diwale SR. Effect of Manures on soil properties and crop response under cowpea-green gram- rice cropping sequence in lateritic soil. Ph. D Thesis submitted to Dr. Balasaheb Sawant Konkan Krushi Vidyapeeth, Dapoli, Dist. Ratnagiri, (M.S); 2012.
16. Bhosale AR. Effect of graded levels of nitrogen and potassium on yield and quality of watermelon in lateritic soils of Konkan, M.Sc. (Ag.), Thesis submitted to Dr. Balasaheb Sawant Konkan Krushi Vidyapeeth, Dapoli, Dist. Ratnagiri, (M.S); 2016.
17. Preethi P, Raja P, Sehgal J and Gajbhiye KS. Study on mineralogy of some selected soils from hot humid to per humid ecosystem of Kozhikode, Palaghat and Emalkulam areas in Kerala. Journal of Indian Society of Soil Science.1998;46:430-435.
18. Hapse DG, Murkute SB, Zende NA. Effect of vermicompost on sugarcane yield and sugar recovery. 10th annual State Level Sugarcane Development Workshop on low cost Technology for cane and sugar production organized by U.S. I. Manjari (BK) on 9-10 the July; 1993.
19. Kadam RG. Effect of vermicompost with and without inorganic fertilizers on yield, quality and mineral nutrition of cowpea-cowpea cropping sequence. M.Sc. (Agri.)

- thesis, submitted to Konkan Krishi Vidyapeeth, Dapoli Dist. Ratnagiri (MS); 2000.
20. Yaduvanshi NPS. Substitution of inorganic fertilizers by organic manures and their effect on soil fertility in rice-wheat rotation on reclaimed Sodic Soil in India. *Journal of Agricultural Sciences*. 2003;14(2):161-168.
 21. Gedam VB, Rametke JR, Rudragouda, Powar MS. Influence of organic manures on yield, nutrient uptake of groundnut and change in physico-chemical properties of soil after harvest of groundnut. *Crop Research*. 2008;36(1, 2 & 3):111-114.
 22. Gadade OB. Effect of integrated nutrient management on yield, quality and nutrient uptake by cowpea (*Vigna unguiculata* (L) Walp) in lateritic soil. M.Sc. (Agri) thesis, submitted to Konkan Krishi Vidyapeeth, Dapoli; 2007.
 23. Yadav DS, Kumar V, Yadav V. Effect of organic farming on productivity, soil health and economics of rice (*Oryza sativa*)-wheat (*Triticum aestivum*) system. *Indian Journal of Agronomy*. 2009;54(3):267-271.
 24. Sinha NP, Prasad B, Ghosh AB. Effect of continuous use of fertilizers on yield and nutrient uptake in wheat-soybean-potato cropping system. *Journal of the Indian Society of Soil Science*. 1981;29:537-542.
 25. Talashilkar SC, Nagwekar DD, Akhave SR, Dabke DJ. Effect of integrated use of manures and fertilizers on coconut yield and changes in available nutrient content and N, P and K fractions of Inceptisols of Konkan region. *Journal of Plantation Crops*. 2008;36(2):112-116.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/118962>