



Antinutrient Contents of Fermented and Extruded Unripe Plantain and Pigeon Pea Blends

Orekoya, Elias Similoluwa ^{a*}, Ojokoh Anthony Okhonlaye ^b,
Arogunjo Ayodele Openla ^a, Akinwunmi Iseoluwa Moses ^a,
Ajayi-Moses Oluwatayo Benjamin ^a, Gabriel Paul Oladimeji ^a
and Aribisala Jamiu Olaseni ^a

^a Department of Microbiology, Federal University of Technology, Akure, Nigeria.
^b Department of Biotechnology, Federal University of Technology, Akure, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. The etiquette and first draft of the manuscript was drafted by authors OES, OAO designed the study. Second draft and corrections were performed by authors AIM managed the literature searches. Authors AMOB performed the statistical analyses. Authors GPO and AAO, AJO managed the analyses of the study. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/SAJRM/2022/v14i1260

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/69401>

Original Research Article

Received 08/04/2021
Accepted 15/06/2021
Published 09/12/2022

ABSTRACT

The study investigated effects of fermentation and extrusion on the antinutrient composition of unripe plantain and pigeon pea blends. The blended samples were prepared in three combinations (A=100g unripe plantain; B= 70g unripe plantain: 30g pigeon pea; C= 50g unripe plantain: 50g pigeon pea) and separated into four batches (i.e. first batch = preconditioned and fermented; second batch = extruded; third batch = fermented and extruded; and fourth batch =

*Corresponding author: Email: orekoyasimiloluwa@yahoo.com;

unfermented/unextruded). The blended samples were fermented semi-solid state fermentation. The anti-nutrient content of fermented and extruded blends decreased significantly ($P<0.05$) when compared to the raw blends. Hence, it can be concluded based from the available information from this study that fermentation and extrusion decreased the antinutrient composition of unripe plantain and pigeon pea blends.

Keywords: Unripe plantain; pigeon pea; fermentation; extrusion.

1. INTRODUCTION

Fermentation and extrusion improve the nutritional value of foods by reducing the water-binding capacity of cereal flour. This allows the fortified to have a free-flowing consistency even with high proportion of flour. Extrusion has been reported as an effective processing treatment to improve the nutritional quality of cereals [1]. In the developing world, fermentation is one of the oldest technologies used for food processing and preservation. Fermentation reduces antinutrient properties of foods. It can be described as a desirable process of biochemical modification of primary food products brought about by microorganisms and their enzymes [2] Extrusion cooking technology has been described as a process in which raw materials are heated and worked upon mechanically while passing through compression screws [3].

Plantain (*Musa paradisiaca*) is a giant perennial crop, cultivated in many tropics and subtropical countries of the world [4]. Plantains are staple food that provides 60 million people with 25% calories [5]. Plantain is used as a source of starchy staple food for millions of people in Nigeria. Mature plantain pulp is rich in iron, potassium and vitamin A but low in protein and fat [6]. Unripe plantain meal is usually consumed by diabetics to reduce postprandial glucose level. This is because the propensity of individual to develop diabetes and obesity is due to the increased consumption of carbohydrate-rich foods with a high glycemic index [7].

Pigeon peas are leguminous shrubby herb, with trifoliolate leaves, yellow flowers and flattened pods that is much cultivated especially in the tropics [8]. Pigeon pea is well adapted to the tropical regimes. One of the best solutions to protein energy malnutrition in developing countries is supplementing cereals with protein rich legumes. Pigeon pea flour has been tested and found to be suitable as a protein source for supplementing cereal food products due to its high level of protein, iron and phosphorus [9].

The main objective of this study is to evaluate the fermentation and extrusion effects on the antinutrient compositions of unripe plantain and pigeon pea flour blends for human consumption.

2. MATERIALS AND METHODS

2.1 Collection of Samples

Green matured unripe plantain and pigeon pea seeds used for this study were obtained from Oja Oba, Akure metropolis in Ondo state, Nigeria.

2.2 Processing of Unripe Plantain Flour

The unripe plantain was sorted for maturity and cleaned by washing with water. The clean unripe plantains were peeled and sliced thinly into 2 mm diameter and sun dried for 72 hours. The dried unripe plantain was then fed into a Bental attrition mill (Model 200L090). The milled flour was sieved with 0.25 mm mesh sieve into fine flour and kept in an air tight container.

2.3 Processing of Pigeon pea Flour

Pigeon pea seeds were cleaned by sorting out dirt and stones. The cleaned pigeon pea seeds were coarsely milled to separate the coat from the cotyledon. The husk was separated from the seed by blowing air into it. The dehulled pigeon pea seeds were milled into fine flour using an attrition mill after which it was sieved through 0.25 mm mesh. The pigeon pea flour was kept in an airtight container.

2.4 Formation of Pigeon pea-plantain Blends

The unripe plantain and pigeon pea flours were formulated in the ratio of (unripe plantain: pigeon pea) 100:0; 70:30; and 50:50.

Sample A (100:0) = 100% unripe plantain flour
Sample B (70:30) = 70% unripe plantain flour
and 30% pigeon pea flour
Sample C (50:50) = 50% unripe plantain flour
and 50% pigeon pea flour

2.5 Fermentation and Extrusion of Flour Blends

A batch of the flour blend was fermented using semi- solid state fermentation for 96 hours. 70 ml of sterilized water was added to 100 g of each sample in cleaned containers and properly sealed. The fermentation process was terminated by oven drying at 60°C for 24 hours. Two batches of samples were subjected to extrusion cooking. The first batch consists of the unfermented blends. The blends were hydrated and preconditioned by adding 50 ml of water to 1000 g of the sample and manually mixed in a sterile bowl to ensure even distribution of water. The samples were extruded using a Brabender 20DN single screw laboratory extruder (Brabender OHG, Duisburg, Germany). The second batch of the samples consists of the fermented samples. The fermented samples were also extruded using a Brabender 20DN single screw laboratory extruder (Brabender OHG, Duisburg, Germany). The samples were extruded at 100°C, 20 revolution per minute and feeding rate of 30 kg/h. All the extrudates were air dried for 12 hours after which they were stored at 32°C in sterile polyethylene bags and kept in properly labeled air tight containers. The control which consists of the raw blends which were neither fermented nor extruded was kept in air tight containers.

2.6 Determination of Antinutrients

2.6.1 Determination of phytate

The phytate content in each sample was determined using the modified method of Chen (2004). Phytate was extracted by adding 0.1g of the sample into 100 ml 0.2M HCl and shaken for 1 hour before centrifuging at 5000 rpm for 15 minutes. A 0.5 ml of the supernatant was pipetted into a test tube fitted with ground glass stopper before adding 1ml acidic ammonium iron (3) sulphate dodecahydrate (0.2g $\text{NH}_4\text{Fe}(\text{SO}_4)_2$)

2.6.2 Determination of alkaloid content

The sample (100g) was grounded and then extracted with methanol for 24 hours in a continuous extraction (soxhlet) apparatus. The extract was filtered and methanol was evaporated on a rotary evaporator under vacuum at a temperature of 45°C to dryness. A part of this residue was dissolved in 2 N HCl and then filtered. One ml of this solution was transferred to a separatory funnel and washed with 10 ml

chloroform (3 times). The pH of this solution was adjusted to neutral with 0.1 N NaOH. Then 5 ml of BCG solution and 5 ml of phosphate buffer were added to this solution. The mixture was shaken and the complex formed was extracted with 1, 2, 3, and 4 ml chloroform by vigorous shaking. The extracts were collected in a 10ml volumetric flask and diluted to volume with chloroform. Presence of alkaloid was confirmed by Dragen droff's method. A part of extract was dissolved in dilute HCL and 2 drops of Dragon drop's was added, a crystalline precipitate indicates presence of alkaloid [10].

2.6.3 Determination of saponin

The sample was grounded and 20g of each were put into a conical flask and 100 cm³ of 20 % aqueous ethanol was added. The samples were place over a hot water bath for 4 hours with continuous stirring at 55°C. The mixture was filtered and the residue re extracted with another 200 ml of 20 % ethanol. The combined extracts were reduced to 40ml over water bath at 90°C. The concentrate was transferred into 250 ml separating funnel and 20 ml of diethyl ether was added and shaken vigorously. The aqueous layer was recovered while the ether was later discarded. The purification process was repeated and 60 ml of n- butanol was added. The combined n-butanol extract was washed twice with 10 ml of 5 % aqueous sodium chloride. The remaining solution was heated in a water bath. After evaporation, the sample was dried in the oven to a constant weight; the saponin content was calculated as percentage [11].

$\% \text{ Saponin} = (\text{Initial weight} - \text{final weight of the sample}) / \text{Initial weight} \times 100$

2.6.4 Determination of phenol

The concentration of phenolics in the sample was determined using spectrophotometric method [12]. Methanolic solution of the sample in the concentration of 1 mg/ml was used in the analysis. The reaction mixture was prepared by mixing 0.5 ml of methanolic solution of the sample, 2.5 ml of 10% Folin-Ciocalteu's reagent dissolved in water and 2.5 ml of 7.5 % NaHCO_3 . Blank was concomitantly prepared, containing 0.5 ml methanol, 2.5 ml 10 % Folin-Ciocalteu's reagent dissolved in water and 2.5 ml of 7.5 % of NaHCO_3 . The samples were thereafter incubated in a thermostat at 45°C for 45 min. The absorbance was determined using spectrophotometer at $\lambda_{\text{max}} = 765 \text{ nm}$. The

samples were prepared in triplicate for each analysis and the mean value of absorbance was obtained. The same procedure was repeated for the standard solution of gallic acid and the calibration line was constructed. Based on the measured absorbance, the concentration of phenolics was read in (mg/ml) from the calibration line; then the content of phenolics in extracts was expressed in terms of gallic acid equivalent (mg of GA/g of extract) [12,13].

2.6.5 Determination of tannin content

A 200mg of finely ground sample was weighed into a 50 ml sample bottle. Ten (10) ml of 70 % aqueous acetone was added and properly covered. The bottles were put in an ice bath shaker for 2 hours at 30°C. Each solution was then centrifuged and the supernatant stored in ice. A 0.2 ml of each solution was pipetted into test tubes and 0.8 ml of distilled water was added. Standard tannic acid solutions were prepared from a 0.5 mg/ml stock and the solution made up to 1 ml with distilled water. A 0.5 ml of folin reagent was added to both sample and standard followed by 2.5 ml of 20% Na₂CO₃. The solutions were then vortexed and allowed to incubate for 40 minutes at room temperature after which absorbance was read at 725 nm against a reagent blank concentration of the samples from a standard tannic acid curve [14].

2.7 Statistical Analysis

Statistical analyses of the Data were obtained using SPSS statistical software (SPSS for window version 20). Data obtained as mean standard deviations were analyzed by Analysis of Variance (ANOVA), followed by Duncan's New Multiple Range Test ($P < 0.05$) to determine the significant differences between the mean values.

3. RESULTS

3.1 Anti-Nutritional Composition of Unripe Plantain and Pigeon Pea Flour Blends

3.1.1 Changes in alkaloid content

The alkaloid content of unripe plantain and pigeon pea blends are shown in Table 1. The result obtained from the anti-nutrient composition of the samples indicated that the raw samples contained the highest alkaloid content ranged from 1.47±0.03 to 3.83±0.02. There was no significant difference ($P \leq 0.05$) in the fermented-

unextruded blends (0.75±0.02 to 1.13±0.04) and unfermented-extruded blends (0.88±0.03 to 1.46±0.01). The fermented-extruded blends recorded significantly low alkaloid content ranging from 0.55±0.01 to 1.03±0.05.

3.1.2 Changes in phenol content

The phenol content of unripe plantain and pigeon pea blends are shown in Table 1. There was significant difference between the phenol content of all the blends. Raw blends had the highest phenol content with values ranged from 11.17±0.01 to 17.03±0.00. Fermented blends had phenol content ranging from 2.35±0.03 to 15.29±0.01. Extruded unfermented blends had phenol content with values ranging from 1.13±0.0 to 9.15±0.03. Extruded fermented blends had the lowest phenol content ranging from 1.03±0.01 to 7.52±0.01.

3.1.3 Changes in tannin content

Tannin content of the blends is shown in Table 1. There was significant difference ($P \leq 0.05$) between sample A, B and C of the raw blends (0.85±0.01, 1.63±0.00 and 1.28±0.00 respectively). There was no significant difference ($P \leq 0.05$) for samples A and B for the fermented blends (0.47±0.00 and 0.39±0.00 respectively), but sample C recorded significant difference ($P \leq 0.05$). Extruded unfermented blends had no significant difference for sample B and C (1.07±0.01 and 1.03±0.01 respectively) but sample a recorded significant difference ($P \leq 0.05$). The extruded fermented blends had values ranging from 0.25±0.01 to 1.13±0.01.

3.1.4 Changes in saponin content

Saponin content of the blends is shown in Table 1. The raw blends had the highest saponin content (1.32±0.08 to 1.99±0.07). Saponin content reduced significantly in the fermented unextruded blends having values that ranged from (0.31±0.04 to 0.78±0.06). There was slight decrease in the saponin content of unfermented extruded (1.25±0.24 to 1.52±0.13) and fermented extruded blends (0.57±0.13 to 1.07±0.23).

3.1.5 Changes in phytate content

Phytate content of the blends are shown in Table 1. There was significant difference ($P \leq 0.05$) in the phytate content of the raw blends, fermented blends and fermented-extruded blends. The extruded-unfermented blends of

Table 1. Anti-nutrient composition of unripe plantain and pigeon pea blends (mg/g)

Sample codes	Alkaloid (mg/g)	Phenol (mg/g)	Tannin (mg/g)	Saponin (mg/g)	Phytate (mg/g)
ARF	1.47±0.01 ^d	11.17±0.00 ^g	0.85±0.01 ^{ab}	1.32±0.00 ^c	23.17±0.01 ^g
BRF	2.25±0.01 ^e	17.03±0.01 ^j	1.63±0.02 ^d	1.79±0.01 ^e	21.35±0.02 ⁱ
CRF	3.83±0.00 ^f	16.25±0.00 ^j	1.28±0.00 ^c	1.99±0.00 ^f	22.69±0.01 ⁱ
AFU	1.13±0.01 ^c	2.35±0.00 ^c	0.47±0.00 ^a	0.78±0.00 ^{ab}	13.37±0.00 ^b
BFU	1.07±0.00 ^c	15.29±0.00 ⁱ	0.39±0.02 ^a	0.69±0.01 ^a	11.53±0.00 ^a
CFU	0.75±0.01 ^b	13.48±0.01 ^h	1.15±0.00 ^b	0.31±0.01 ^a	15.25±0.00 ^d
AUE	1.46±0.00 ^d	1.13±0.01 ^a	0.75±0.00 ^{ab}	1.25±0.00 ^c	17.43±0.01 ^d
BUE	1.37±0.00 ^d	4.27±0.00 ^b	1.07±0.00 ^b	1.30±0.00 ^c	18.15±0.00 ^e
CUE	0.88±0.00 ^b	9.15±0.01 ^f	1.03±0.00 ^b	1.52±0.00 ^d	18.03±0.01 ^e
AFE	1.03±0.01 ^c	1.03±0.01 ^a	0.25±0.00 ^a	0.57±0.00 ^a	15.31±0.00 ^c
BFE	0.71±0.02 ^b	7.52±0.00 ^e	1.13±0.01 ^b	0.63±0.00 ^a	12.03±0.01 ^a
CFE	0.55±0.00 ^a	5.97±0.01 ^d	1.01±0.00 ^b	1.07±0.02 ^b	13.77±0.00 ^b

Values are means of triplicate determinations ± SD. Means in the same column with different superscripts are significantly different ($P \leq 0.05$); Keys: ARF= Raw unripe plantain flour 100g; BRF= Raw unripe plantain flour 70g and raw pigeon pea flour 30g; CRF= Raw unripe plantain flour 50g and raw pigeon pea flour 50g; AFU= Fermented unextruded unripe plantain flour 100g; BFU= Fermented unextruded unripe plantain flour 70g and pigeon pea 30g; CFU= Fermented unextruded unripe plantain flour 50g and pigeon pea 50g; AUE= Unfermented extruded unripe plantain flour 100g; BUE= Unfermented extruded unripe plantain flour 70g and pigeon pea 30g; CUE= Unfermented extruded unripe plantain flour 50g and pigeon pea 50g; AFE= Fermented extruded unripe plantain flour 100g; BFE= Fermented extruded unripe plantain flour 70g and pigeon pea 30g; CFE= Fermented extruded unripe plantain flour 50g and pigeon pea 50g

sample B and C (18.15 ± 0.02 and 18.03 ± 0.01 respectively) showed no significant difference but extruded-fermented sample A showed significant difference ($P \leq 0.05$). The raw blends had the highest phytate content with values that ranged from 21.35 ± 0.04 to 23.17 ± 0.01 for samples A to C. Fermented samples had the least phytate content with values that ranged from 11.53 ± 0.23 to 15.25 ± 0.00 for sample A to C.

4. DISCUSSION

The results of anti-nutrient components revealed that extrusion cooking reduced phytate levels. It could be expected that lowering this compound would enhance the bioavailability of minerals such as Zinc and Iron in the extrudates since phytic acid has been implicated in making these minerals unavailable as reported by Anuonye *et al.* [15]. Decrease in phytate content may be partly due to the formation of insoluble complexes between phytate and microbial degradation. It could also be due to the heat labile nature of phytic acid. The effect of fermentation on phytic acid may be due to the activity of enzyme like phytase produced by fermenting micro flora [16]. The reduction in phytate level of fermented blends could be interpreted as the main reason behind the observed increase in the concentrations of minerals in the fermented samples. Reduction in the phytate content was also reported by Hassan *et al.* [17] when he evaluated the effect of blanching and fermentation on cocoyam.

The level of tannin was significantly reduced in the unfermented extruded blends, fermented unextruded blends as well as fermented extruded blends. Decrease in tannin after fermentation may be as a result of microbial activity during fermentation like tannase and metabolites for microbial fermentation [18]. Tannins form insoluble complexes with proteins thereby decreasing protein digestibility [19]. Tannins also decrease palatability, cause damage to intestinal tract and enhance carcinogenesis [20]. Reduction of tannin in the fermented blends may be as a result of the fact that tannins are polyphenols and all polyphenolic compounds are water soluble in nature [21]. Therefore, reduction in tannin content of the fermented blends may be attributed to leaching out of phenols into the fermenting media [21]. Similar result was also reported by Hassan *et al.* [17]. Korus *et al.* [22] attributed the decrease in the phenolic content of beans to high temperature when he evaluated the effect of extrusion on the phenolic

composition and antioxidant activity of dry beans. Suazo *et al.* [23] reported that the concentration of phenol reduced with fermentation.

Saponin content was reduced in all the flour blends but significantly reduced in fermented extruded blends. Saponins have anti-hyper cholesterol, anti-inflammatory, cardiac depressant property and also appear to kill or inhibit cancer cells without killing the normal cells in the process [24]. The saponin content of the pigeon pea flour samples were reduced by each of the processing methods (fermentation and extrusion). The raw samples had the highest saponin values increasing from 1.32 ± 0.00 mg/g to 1.99 ± 0.00 mg/g while the fermented extruded blends had the lowest saponin values ranging from 0.57 ± 0.00 mg/g to 1.07 ± 0.02 mg/g. It has been reported that processing techniques such as boiling, sprouting and fermentation reduce anti-nutritional content of legume flours [25].

Alkaloids are important poisons produced by plants as active defense against aggressors [26]. There was significant difference ($P < 0.05$) in the alkaloid content of the unripe plantain and pigeon pea flour samples. The raw samples had the highest alkaloid content (1.47 ± 0.01 mg/g to 3.83 ± 0.00 mg/g) in comparison with the fermented extruded samples with a significant reduction ranging from 1.03 ± 0.01 mg/g to 0.55 ± 0.00 mg/g in the total alkaloid content. Alkaloids are nitrogen containing naturally occurring compounds found to have antimicrobial properties due to their ability to intercalate with DNA of the microorganisms [27]. The reduction of alkaloids due to boiling, sprouting and fermentation had been reported respectively by Nwanekezi *et al.* [25]. The presence of alkaloids in this study may lead to healing of wounds, varicose ulcers, hemorrhoids, frostbite and burn in herbal medicines as reported by Ndulaka and Obasi [28].

The rate at which these anti-nutrients affect the availability of nutrients by chelating the nutrients and making them available for utilization in the system will be relatively reduced. Since the higher the level of anti-nutrient, the lower the bioavailability of the nutrient and minerals contained therein [29].

5. CONCLUSION

This investigation shows that fermentation and extrusion of the blending of unripe plantain and pigeon pea has the potential of reducing the

antinutrient compositions of the samples thereby producing enriched complementary food for improving the health of malnourished children of developing countries.

ACKNOWLEDGEMENT

We express our gratitude to the Head of Department of Microbiology, Federal University of Technology Akure, Professor M. K. Oladunmoye and the technical staff of the department for their collective contributions.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Amadou I, Gbadamosi OS, Le GW. Millet based traditional processed foods and beverages: A review. *Cereal Foods World*. 2011;56:115-121.
2. Muchoki CN, Imungi JK, Lamuka PO. Changes in beta-carotene, ascorbic acid and sensory properties in fermented, solar-dried and stored cow-pea leaf vegetables. *African Journal of Food Agricultural Nutrition and Development*. 2010;7:16–26.
3. Iwe MO. The science and technology of soybean: Chemistry nutrition processing and utilization. 1st Edition. Rejoint Communication Services Limited. Enugu State. Nigeria. 2003;680.
4. Akomolafe OM, Aborisade AT. Effect of stimulated storage conditions on the quality of plantain (*Musa paradisiaca*) fruit. *International Journal of Agriculture Research*. 2007;2(12):1037-1042.
5. FAO. Production Yearbook for 2005. FAOSTAT Data, Food and Agriculture Organisation of the United Nations, Rome; 2005.
6. Adeniji TA, Sanni LO, Barimalaa LS, Hart AD. Determination of micronutrients and colour variability among new plantain and banana hybrid flour. *World Journal of Chemistry*. 2006;1(1):23-27.
7. Oboh HE, Erema VG. Glycemic indices of processed unripe plantain (*Musa paradisiaca*) meals. *African Journal of Food Science*. 2010;4(8):514-521.
8. Damaris AO. The potential of pigeon pea (*Cajanus cajan* (L.) Millsp.) in Africa. *Natural Resources Forum*. 2007;31(32): 297–305.
9. Harinder K, Sharma B. Studies on the baking properties of wheat: Pigeon pea flour blends. *Journal of Plant Foods for Human Nutrition*. 1999;54:217–226.
10. Ali M, Mostafa P, Fariba S, Shirin P, Reza F, Sedigheh S. Alkaloids and flavonoid of fenugreek seeds (*Trigonella foenum-graecum* L.) with antinociceptive and anti-inflammatory effects. *Journal of Food and Chemical Toxicology Elsevier*. 2012;50:2503-2507.
11. Amir MK, Rizwan, AQ, Faizan U, Syed AG, Asia N, Sumaira S, Muhammad KL, Muhammad YL, Shafiq UR, Ishtiaq H, Waheed M. Phytochemical analysis of selected medicinal plants of Margalla hills and surroundings. *Journal of Medicinal Plants Research*. 2011;5(25):6017-6023.
12. Singleton VI, Orthofer R, Lamuela-Raventos RM. Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Journal of Methods on Enzymology*. 1999;299:152-178.
13. Milan SS. Total phenolic content, flavonoid concentration and antioxidant activity of *Marrubium peregrinum* L. Extracts. *Kragujevac Journal of Science*. 2011;33:63-72.
14. Bansa A, Adeyemo SO. Evaluation of antibacterial properties of tannins isolated from *Dichrostachys cinerea*. *African Journal of Biotechnology*. 2010;6(15):1684- 5315.
15. Anuonye JC, Onu JO, Egwin E, Adeyemo SO. Nutrient and anti-nutrient composition of extruded acha/soybean Blend. *Journal of Processing*. 2009;34:680-691.
16. Amit V, Sewa R, Vikas B. Cereal phytases and their importance in improvement of micronutrients bioavailability. *Journal of Biotechnology*. 2017;7(42):1-7.
17. Hassan GF, Yusuf L, Adebolu TT, Onifade AK. Effect of fermentation on mineral and anti-nutritional composition of cocoyam. *Sky Journals of Food Science*. 2015;4(4):42-49.
18. Sorour MA, Mehanni AE, Taha EM, Rashwan AK. Changes of total phenolics, tannins, phytate and antioxidant activity of two sorghum cultivars as affected by processing. *Journal of Food and Dairy Science*. 2017;8(7):267-274
19. Uzoechina EO. Effects of separation on the phytochemical properties of legumes. *Journal of Natural Sciences Research*. 2007;66:361-362.

20. Anuonye JC. Some functional properties of extruded acha/soybean blends using response analysis. *African Journal of Food Science*. 2012;6(10):269-279
21. Udenzi EA, Ekwu FC, Isinguzo JN. Antinutrient factors of vegetable cowpea (*Sesquipedalis*) seeds during thermal processing. *Pakistan Journal of Nutrition*. 2007;6(10):194-197.
22. Korus J, Gumul D, Czechowska K. Phenolic composition and antioxidant activity of dry beans. *Journal of Food Technology and Biotechnology*. 2007; 45:139-146.
23. Suazo Y, Davidov-Pardo G, Arozarena I. Effect of fermentation and roasting on the phenolic concentration and antioxidant activity of cocoa from Nicaragua. *Journal of Food Quality*. 2014;37(1):50–56.
24. Ojinnaka MC, Odimegwu EN, Chidiebere FE. Comparative study on the nutrient and antinutrient composition of the seeds and leaves of *Uziza (Piper Guineense)*. *Journal of Environmental Science, Toxicology and Food Technology*. 2016;10(8):42-48.
25. Nwanekezi EC, Ehirim FN, Arukwe DC. Combined effects of different processing methods on vitamins and anti-nutrients contents of pigeon pea (*Cajanus Cajan*) flour. *Journal of Environmental Science, Toxicology and Food Technology*. 2017;11(4):73-81.
26. Kasproicz-Potocka M, Zaworska A, Gulewicz P, Nowak P, Frankiewicz A. The effect of fermentation of high alkaloid seeds of *Lupinus angustifolius* var. Karo by *Saccharomyces cerevisiae*, *Kluyveromyces lactis*, and *Candida utilis* on the chemical and microbial composition of products. *Journal of Food Processing and Preservation*. 2017;10(11):1-10.
27. Kasolo NJ, Bimenya SG, Ojok L, Ochieng J, Ogwal-Okeng WJ. Phytochemicals and uses of *Moringa oleifera* leaves in Ugandan rural communities. *Journal of Medicinal Plants Research*. 2010;4(9):753-757.
28. Ndulaka JC, Obasi NE. Production and quality evaluation of cookie-like products formulated from blends of wheat flour and garri from biofortified cassava varieties. *Research Journal of Food Science and Quality Control*. 2018;4(1):20-41.
29. Fagbemi AO, Ojokoh AO. Effects of fermentation and extrusion on the proximate and organoleptic properties of cowpea-plantain flour blends. *British Microbiology Research Journal*. 2016; 13(4):1-13.

© 2022 Similoluwa et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/69401>