



Amino Acids Analysis of the Four Mushroom Species, Antioxidant Activity and Immunostimulant Evaluations of their Polysaccharides Extracts

Khanittha Chawanoraset ^{a*}, Tanapak Inyod ^b, Patrawadee Kengkwasigh ^a,
Sarunya Laovitthayangoon ^a, Jarussang Kammultri ^{c,d},
Vachiraporn Ajavakom ^{c,d}, Runchana Funk ^a, Kittirat Prangchan ^a,
Thanapat Termarom ^b and Sirinan Thubthimthed ^a

^a Expert Centre of Innovative Herbal Products, Thailand Institute of Scientific and Technological Research, Pathumthani 12120, Thailand.

^b Expert Centre of Innovative Agriculture, Thailand Institute of Scientific and Technological Research, Pathumthani 12120, Thailand.

^c Industrial Metrology and Testing service Centre, Thailand Institute of Scientific and Technological Research, Pathumthani 12120, Thailand.

^d Department of Chemistry and Center of Excellence for Innovation in Chemistry, Faculty of Science, Ramkhamhaeng University, Bangkok 10240, Thailand.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/CJAST/2022/v41i1531718

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/87176>

Original Research Article

Received 06 March 2022

Accepted 13 May 2022

Published 21 May 2022

ABSTRACT

Mushrooms have been valued as edible and medical resources. This project aimed to evaluate of four mushrooms such as *Pleurotus ostreatus* (Fr.) Kummer, *Pleurotus opuntiae* (Dur.& Lev.) Sacc., *Pleurotus sajor-caju* (Fr.) Sing, and *Ganoderma lucidum* (Curtis) for amino acids, two extraction methods, antioxidant activity, total phenolic content, ¹H NMR analysis and *in vitro* immunostimulant activity, compared the cultivation between TISTR and Nakhon Prathom area. The results showed that glutamic acid and aspartic acid were abundant amino acids, while alanine, leucine, lysine, arginine, serine, phenyl alanine and threonine were found varied in those mushrooms. The most

*Corresponding author: E-mail: khanitha@tistr.or.th;

common amino acids found in *P. sajor-caju* were found in lower amounts in *G. lucidum*. There amino acids content was contrast with the antioxidant activity and total phenolic content. Meanwhile the ¹H NMR chemical shifts illustrated signals of a mixture of polysaccharides resonated at δ 3.0-4.0 ppm, and fatty acids at δ 0.70-2.30 ppm of the A extract as well as the two signals of anomeric protons α -glucan and β -glucan of the B extract appeared at δ 5.36 ppm and 4.47 ppm, respectively. The bioassay *in vitro* immunostimulant activity of all mushroom extracts showed good immunostimulant booster at concentration of 0.5 mg/mL. It was found that the A and B extracts of *G. lucidum* were the most potent immunostimulant activity followed by the A extract of *P. sajor-caju*, and they had higher activity than β -glucan standard with % phagocytic activity of 187.12 \pm 2.17, 166.23 \pm 0.56, 155.00 \pm 2.93, respectively. Future works will focus on their *in vivo* immunostimulant activity, healthy drinks and nutraceutical product development.

Keywords: *Pleurotus ostreatus* (Fr.) Kummer; *Pleurotus opuntiae* (Dur.& Lev.) Sacc.; *PLEUROTUS sajor-caju* (Fr.) Sing; and *Ganoderma lucidum* (Curtis); immunostimulant activity.

1. INTRODUCTION

Immune response is an important physiological process to identify and destroy foreign harmful substances as well as organisms to protect diseases, in which macrophages play key roles in phagocytosis, cytotoxicity and intracellular killing activities. Macrophages can resist to pathogens either directly through phagocytosis or indirectly by producing related factor such as nitric oxide (NO), interleukins (IL), tumor necrosis factor- α (TNF- α) and reactive oxygen species (ROS). [1] In clinical practice, immunomodulators are classified into three categories including immunosuppressants, immunostimulants and immuno adjuvants. Immune system modulations are used common as prophylactic medicine for increasing healthy people where modulations are used common as prophylactic medicine for increasing a number of healthy people whereas immunomodulators are synthetic or semi-synthetic components therefore, natural immunomodulators have been in growing interest for overall health-benefiting effects to humans with no or less toxicity [2,3]. Mushrooms, a fungi kingdom member with well-known medicinal properties has drawn significant interest due to their immunomodulatory and antioxidant attributes, anti-inflammatory, antimicrobial, antidiabetic, cardiovascular protective, hepatoprotective and anticancer potentials [4]. Several active components for example, essential amino acids, proteins, vitamins and mineral, fiber, and polysaccharides in mushrooms fruiting immunomodulatory responses [5,6]. Mushroom polysaccharides display antimicrobial properties against pathogenic bacteria and viruses and indicated the presence of rhamnose, xylose, fucose, arabinose, fructose, glucose, mannose, manitol, sucrose, maltose and trehalose

predominantly mushroom-based polysaccharides fractions [7]. There was reported for encapsulated polysaccharides extract in alginate gel beads for external influences protect and extend for their applications [8]. The polysaccharides seem to orchestrate antitumor properties via the activation of host immune response, inferred that the tumor cells have been no destroy directly but instead they were indirectly an impact by stress preventing on the body [9, 10]. Moreover, polysaccharides were compared in different mushroom species such as *A. bisporus*, *A. brasiliensis*, *Phellinus linteus* and *G. lucidum* and other active compounds, found a positive correlation between glucan level and antioxidation activity in *G. lucidum* [11]. The methanolic extracts of the wild *G. lucidum*, native to the Himalaya, including *P. ostreatus*, *P. erynii*, *P. nebrodensis* were clearly documented for their significant antioxidant potential [12, 13]. *G. lucidum* polysaccharides were reported for their worked well against MCF-7 cell lines [14]. Furthermore, the mechanism of polysaccharides for their antioxidant involving the hydrogen atom transfer (HAT) reactions and mostly occur in their neutral form, while the electron transfer (ET) mechanism usually occurs in acidic polysaccharides [15]. The *Pleurotus* spp. Possess a vast diversity of polysaccharides in particular heteropolysaccharides and glucans, constituting special biological activities for antioxidant, anti-inflammatory and immunomodulatory activities [16, 17]. There were predominantly consumed mushrooms in Asian countries and have been popularized worldwide according to their potential on essential amino acids, minerals, protein and biological active polysaccharides [18]. The World has been suffering from the 2019 COVID-19 pandemic and now there were treatments with medicine available approved from the U.S. Food and Drug

Administration (F.D.A.) and the first pill designed to treat symptomatic COVID-19 approved by the UK medicine regulator which these can be deployed as potential therapeutics against COVID-19 [19]. However, the problem with commercial medications is the increased risk of drug resistance development. Natural substances such as mushrooms reported previously, displayed clearly for their antiviral and anti-inflammatory activities and may hold natural remedies against COVID-19 [20]. Therefore, it is important to discover these active ingredients from mushroom bodies to examine their therapeutic potential for immunostimulant activity for leading mushroom-based drug discovery of natural active ingredients from high potential mushrooms in this study.

2. MATERIALS AND METHODS

2.1 Materials

2.1.1 Plant materials

Four mushroom species such as *Pleurotus ostreatus* (Fr.) Kummer, *Pleurotus opuntiae* (Dur.&Lev.) Sacc., *Pleurotus sajor-caju* (Fr.) Sing and *Ganoderma lucidum* (Curtis) were obtained from Dr. Tanapak Inyod, Expert Centre of Innovative Agriculture, Thailand Institute of Scientific and Technological Research, Ministry of Ministry of Higher Education, Science, Research and Innovation, Thailand. All the isolates were collected from the mushroom farm in Ratchaburi Province, Thailand. The mushroom pure cultures were transferred into PDA slants and maintained at room temperature. The three mushrooms species, *Pleurotus opuntiae* (Dur.&Lev.) Sacc., *Pleurotus sajor-caju* (Fr.) Sing and *Ganoderma lucidum* (Curtis) were purchased from the local mushroom farm, Tambon Srapattana Community Enterprise in Nakhon Prathom province, Thailand.

2.1.2 Mushroom production

The upper end of the sample bags were opened. Colonized (by mycelium) substrates were arranged on shelves in a greenhouse (temperature, 25-30°C; relative humidity, (85%). The cultures were constantly wet in order to maintain the required relative humidity (85%). The cultures were irrigated by spraying with water, once or twice a day. Primordia will be initiated after 7 days and mature mushrooms (noted as the point where the pileus started to

fold) from each bag were harvested at 10 days after open the bags.

2.1.3 Plant preparation

The mushrooms were individually collected, weighed, washed and torn to pieces by hands, then put each of them separately on aluminium foil sheet to dry in the hot air oven, temp 50°C for 24 hrs. The dried mushrooms were ground in to powder weighed and kept in zip lock bag at room temperature.

2.1.4 Plant extraction

2.1.4.1 Extraction A

The dried powder of *Pleurotus ostreatus* (Fr.) Kummer, *Pleurotus opuntiae* (Dur.&Lev.) Sacc., *Pleurotus sajor-caju* (Fr.) Sing and *Ganoderma lucidum* (Curtis) were individually extracted with Delonized water, temp 25°C, stirred for 7 hrs. The supernatant was collected after filtered and then freeze-dried under FreeZone 6, 12 and 18 Liter, Freeze dryer gave DI water extracts.

2.1.4.2 Extraction B

The dried powder of *Pleurotus Sajor-caju* (Fr.) Sing, *Pleurotus ostreatus* (Fr.) Kummer, *Pleurotus ostreatus* (Jacq. exFr.) Kummer and *Ganoderma lucidum* (Curtis) were individually extracted with with 80% (80:20 ethanol:water) at room temp. for 7 days, then filtered and dried under vacuum using evaporator gave ethanol extracts. mM. Measured their absorbance by using UV-VIS spectrophotometer at 517 nm.

2.2 Total Phenolic Content Analysis

Using Folin-ciocaltea colorimetry method; The stock standard solution of gallic acid solution was prepared conc. 5000 mg/mL in ethanol. The standard solution was pipetted to prepared six concentrations (50, 100, 150, 200, 250 and 300 mg/mL). The sodium carbonate (Na₂CO₃) was added, mixed and adjusted to 10 mL of DI water. The solutions were kept in dark for two hrs. and measured their absorbance by using UV-VIS spectrophotometer at 765 nm.

2.3 Anti-oxidation Activity Test

Free radical scavenging activity assay; The stock standard solution of 2,2-Diphenyl-1-picrylhydrazyl (DPPH) solutin was prepared conc. 607 µM in methanol. Prepared the DPPH

working by pipette DPPH stock solution 1 mL and added methanol 2.1 mL. Pipetted DPPH working solution 200 μ L into Well Plate and measured absorbance by Microplate Reader Sunrise. The stock standard solution of trolox was prepared into 10 mM. Measured their absorbance by using UV-VIS spectrophotometer at 517 nm.

2.4 HPLC Analysis Sample Preparation

HPLC: model LC-20A Series, Shimadzu, Japan, column; Shim-pack ISC-07/S 1504 Na.

Mobile phases: A:0.2N sodium citrate, pH 3.2; B:0.6N sodium citrate+0.2N boric acid, pH10; C:0.2N sodium hydroxide.

Flow rate:0.3 mL/min. **Injection volume:** 10 μ L. **Post column:** Reaction solution A and B with flow rate 0.3 mL/min.

Detector: Fluorescence detector, RF-515, Ex 348nm, Em 450nm. **Calibration curve and linearity:** Measure of amino acids standard for 5 concentrations by using HPLC-LC-20A, duplicate and read, relative percent different (%RPD) \leq 10. Peaks concentration (X) and peaks area (Y), read by Regression Coefficient (R^2) \geq 0.99.

2.4.1 Sample preparation

Weighed mushroom powder into vacuum reaction tube, added 6H HCl 4 mL then mixed by vortex mixer following by Aspirator. The sample solution was digested at 110 $^{\circ}$ C for 24 hrs., adjusted the volume to 10 mL with 0.1 N HCl after the liquid was at room temperature.

2.4.2 Nuclear magnetic resonance (NMR) spectra

The 1 H NMR spectra were measured with a Bruker ASCEND 400 FT-NMR spectrometer, operating at 400 MHz (1 H). The chemical shifts (δ_H and δ_C) were recorded in ppm. with reference to residual solvent signal $CDCl_3$ (δ_H 7.24), CD_3OD (δ_H 4.78, 3.31) where appropriated and coupling constants (J) were given in Hz.

2.5 Immunostimulant Activity

2.5.1 Cytotoxicity: For MTT assay

All cells on 96-well plate were washed out. MTT (10 μ L, 5 mg/ml) and the medium (190 μ L) were

added in each well. The cells were then incubated at 37 $^{\circ}$ C 5% CO_2 for additional 2 hours. The medium containing MTT was discarded, and MTT formazan that had been produced was extracted with 100 μ L of DMSO. The absorbance was read at 570 nm.

2.5.2 Nitrous Oxide (NO) production

For Griess assay, 100 μ L of Griess reagent was mixed with 100 μ L of collected medium in a 96-well plate. The mixture was incubated for 15 min. at room temperature. The absorbance was determined at 540 nm. with the spectrophotometry. The concentration of nitrite was calculated using a calibration standard curve constructed using sodium nitrite dissolved in DMEM.

2.5.3 Phagocytic activity

The RAW264.7 cells were seeded (1×10^5 cells/ml) in a 96-well plate with DMEM medium (10% FBS) and incubated at 37 $^{\circ}$ C in 5% CO_2 for 24 h. The cells were cultured with the eight mushroom extracts KC-MU1 TISTR, KC-MU2 TISTR, KC-MU3 Nakhon Prathom, KC-MU4 TISTR, KC-MU5 Nakhon Prathom, KC-MU6 Nakhon Prathom at 37 $^{\circ}$ C. DMEM and beta-glucan (50 and 100 μ g/ml) were used as negative and positive controls, respectively. The stimulated cells were washed twice with phosphate-buffered saline (PBS), and 100 μ L of 0.075% neutral red solution was added to each well, followed by incubation for 3 h at 37 $^{\circ}$ C. Following removal of the unphagocytized neutral red by PBS, destain buffer (ethanol:DW:acetic acid, 50:1:49) was added to each well at shaker for 1 h. The optical density (OD) value of each well was measured at 540 nm using a microplate reader. Cell phagocytosis (%) was calculated as follows: $OD_S / OD_C \times 100$, where OD_S and OD_C represent the OD values of the stimulated and control wells, respectively. The percentage of phagocytosis in the untreated cells was designated as 100%. All experiments were performed in triplicate.

2.6 Statistical Analysis

Significant differences between two normally distributed data were determined with Student's t-test. The values are reported as the means \pm SEM. Statistical analysis was performed using SPSS (Version 17.0).

3. RESULTS AND DISCUSSION

The result of amino acids determination of all mushroom extracts showed the most abundant of amino acids, including glutamic acid, aspartic acid following by leucine and alanine while phenyl alanine and threonine were found varied in Nang-Rom Hungari (KC-MU2), Nang-Fah-Phuthan-Dam (KC-MU4) and Ling-Zhi (KC-MU6) as can be seen in Table 1. Atri, N.S., *et al* evaluated on five wild edible fungal species of genus *Pleurotus* using Ultra Performance Liquid Chromatography (UPLC) found amino acids namely aspartic acid, arginine, alanine, proline and tyrosine were detected from the samples, aspartic acid (0.37-0.48%), arginine (0.19-0.23%), alanine (0.19-0.11%), Proline (0.02-0.09%) and tyrosine (0.14-0.20%) were found maximum in *Pleurotus pulmonarius*, *P. sajor-caju* (0.44%), *P.cystidiosus*, *P. floridanus* and *P. sapidus* [21] *P. sajor-caju* possesses different amount of dominant amino acids and found that aspartic acid as well as alanine were similar compared to the amino acids in this study.

The immunostimulant activities result of the eight mushroom extracts included A and B extracts were performed to evaluate cell viability and nitric oxide (NO) production and % Phagocytic activity, respectively using MTT and Griess assays. The results showed that the cell viability found that Ling-Zhi (KC-MU6) was the highest % viability at concentration 0.5 mg/mL following by Nang-Fah-Phutan-Dam (KC-MU4), Nang-Rom-Tao (KC-MU1) and Nang-Rom-Hungari (KC-MU2) in extract B, respectively. The cells viability in A extracts were similarly among the four extracts. Whereas nitric oxide production were varied in the range of 31.06±0.40 to 41.32±1.40 µM, at concentration 0.5 mg/mL in A and B extracts. Meanwhile the % Phagocytic activity of RAW 264.7 cells at concentration 0.5 mg/mL showed that Ling-Zhi (KC-MU6) was the most active. The phagocytic activity of the A extractions A were rinsing from KC-MU3, KC-MU2, KC-MU1, KC-MU5, KC-MU4 and KC-MU6 while the B extractions were KC-MU1, KC-MU2, KC-MU4, KC-MU3, KC-MU5 and KC-MU6, respectively. The previous pharmacological activities report of *P. sajor-caju* including antidiabetic, antibacterial, antiproliferation, antioxidant and immunological activities. Carbonero, *et al* [22] documented the *in vitro* immunological activity using Murine macrophage cell line Raw 264.7, NO and cytokine levels as well as NO, TNF-α levels determination, the results showed β-glucan

induced the production of NO and cytokines TNF-α and IL-1β dose of 0.1-300 µg/ml. While the *P. sajor-caju* both A and B extracts in this study illustrated the activity with the dose between 119-165 mg/ml and β-glucan prominent in B extract was the active compounds.

The antioxidation activity of the mushroom in A extracts were increased from KC-MU4, KC-MU1, KC-MU5, KC-MU3, KC-MU2 and KC-MU6, respectively. The activity in B extracts were risen from KC-MU4, KC-MU2, KC-MU5, KC-MU1, KC-MU3 and KC-MU6, respectively. The highest activity among all extracts was KC-MU6 (Ling-Zhi) with the extraction A which was related its total phenolic content found the highest amount in both Ling-Zhi Extractions. The lowest antioxidation activity and total phenolic content were KC-MU4.

The antioxidant activity using *in vitro* DPPH, hydroxyl radical and superoxide anion radical scavenging activities of aqueous extracts showed concentration dependent antioxidant activity in all the assays [22]. The aqueous extract with cold and hot water of *in vitro* DPPH, β-carotene linolenic acid, reducing power ability, phosphomolybdate assay, result of the cold water extract showed higher activity in all assays than hot the water extraction with doses of 50, 100, 250 and 500 µl [23]. Whereas the result of antioxidant activity in this research were in the range of 6.960.31 to 82.55±3.35 mg/ml in the doses of 50, 100, 250 and 300 mg/ml. The previous report of total phenolic content and antioxidant activity on DPPH assays of *P. ostreatus*, *P. sajor-caju*, *P. eryngii* and *P. cystidiosus* exhibited the phenolic content, 17.63±1.30 -7.17±2.24 mg GAE/g and the DPPH assay showed EC₅₀ values 5.00±0.06-9.13±0.17 mg/ml [24]. The results on the antioxidant and total phenolic content reported were higher than those in this experiment according to the methanol solvent can extract high amount of phenolic compounds, led to higher potent of antioxidant activity than water extract. The ¹H NMR spectrum illustrated the A extracts constituting of a mixture of polysaccharides, resonated chemical shift at δ 3.0-4.0 ppm and fatty acids at δ 0.70-2.30 ppm. While the B extracts showed the signals of α-glucan and β-glucan anomeric protons appeared at δ 5.36 ppm and 4.47 ppm, respectively. The results found the extract, containing of polysaccharides possesses considering a better immunostimulant activity than α-glucan and β-glucan composition.

Table 1. Amino acid analysis of the four mushrooms species, anti-oxidation, total phenolic content and immunostimulant activity of the polysaccharide extracts

Amino acid analysis of mushrooms	<i>Pleurotus ostreatus</i> (Fr.) Kummer (Nang Rom Tao)	<i>Pleurotus opuntiae</i> (Dur.&Lev.) Sacc. (Nang Rom Hungari)	<i>Pleurotus Sajor-caju</i> (Fr.) Sing (Nang Fah Phutan Dam)	<i>Ganoderma lucidum</i> (Curtis) (Ling Zhi)		
Amino acid Mg/100g	KC-MU1 TISTR	KC-MU2 TISTR	KC-MU3 Nakhon Prathom TISTR	KC-MU4 TISTR	KC-MU5 Nakhon Prathom	KC-MU6 Nakhon Prathom
Aspartic acid	1742.90	1956.23	1708.60	1980.42	1921.19	916.47
Threonine	837.36	956.79	878.69	1010.91	968.98	630.44
Serine	929.97	1081.02	944.46	1102.32	1024.69	573.80
Glutamic acid	2897.55	3350.86	3595.61	4541.77	3696.89	1133.20
Proline	681.51	793.16	707.93	846.88	832.59	419.01
Glycine	800.82	967.16	816.75	1009.54	906.83	493.73
Alanine	1110.29	1276.92	1141.46	1456.90	1261.88	625.97
Cystine	86.27	101.32	109.61	105.53	131.25	88.05
Valine	762.79	876.80	780.86	930.84	912.08	553.19
Methionine	231.40	254.66	256.93	246.91	303.75	106.52
Isoleucine	609.93	723.51	645.41	725.92	754.06	436.95
Leucine	1073.77	1278.95	1113.68	1276.90	1315.29	718.36
Tyrosine	429.35	458.90	438.17	466.82	523.83	260.41
Phenylalanine	736.68	830.29	927.64	906.64	1442.08	650.26
Histidine	390.62	445.09	387.67	453.53	426.41	227.32
Lysine	976.52	1152.92	931.75	1134.00	1060.24	477.29
Arginine	875.35	1154.38	852.41	979.40	1000.30	420.12
Tryptophan	234.78	273.06	214.46	268.90	224.70	107.23
Antioxidation						
Extraction A	56.88±2.46	18.30±0.48	32.01±0.50	71.37±1.61	41.01±1.06	6.96±0.31
Extraction B	58.01±2.10	73.93±1.75	36.47±4.00	82.55±3.35	63.56±2.37	8.26±0.52
Total phenolic content						
Extraction A	2.01±0.04	2.94±0.12	1.76 ± 0.09	0.99±0.14	1.49 ± 0.09	3.23±0.01
Extraction B	1.25±0.15	0.40±0.32	0.61 ± 0.41	0.37±0.52	0.39 ± 0.63	2.04±0.02
<i>In vitro</i> immunostimulant activity						
Extraction A	143.95±1.64	135.17±4.59	126.25±1.93 (0.25 mg/ml)	155.00±2.93	146.00±0.20 (0.25 mg/ml)	187.12±2.17
Extraction B	113.67±3.06	116.89±3.82	141.63±1.02	119.44±1.77	165.00±1.13	166.23±0.56

4. CONCLUSIONS

Our results demonstrated that the glutamic acid, aspartic acid and alanine exhibited the most plentiful in the four mushrooms. The content of amino acids was found less effect to the immunostimulant activity for example, the smallest amount of amino acids found in KC-MU6 which exhibited the highest activity compared to the others. On the other hand, the polysaccharides and α -glucan as well as β -glucan composition as a major component in the four mushroom play a key role on the potential to improve and stimulate the phagocytic ability of macrophage RAW 264.7 to produce NO, especially in Ling-Zhi mushroom. The four mushroom with rich polysaccharides than α -glucan as well as β -glucan in the extracts, considering for the greater activity, can be prepared and applied to the development and utilization of future new functional foods and healthy drinks.

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Li YH, Li ZK, Li CY. Study on the regulatory effect of crude polysaccharides from *Plantago asiatic* L. on the immune function of human macrophages and its mechanism. *J. Chin. Med. Mater.* 2020;43:2795-2798.
2. El Enshasy HA, Hatti-Kaul R. Trends Biotechnol. Mushroom immunomodulators unique molecules with unlimited applications. 2013;31:668-677.
3. Reis FS, Martins A, Vasconcelos MH, Morales P, Ferreira ICFR. Functional foods based on extracts or compounds derived from mushrooms. *Trends. Food Sci. Technol.* 2017;66:48-62.
4. Moradali MF, Mostafavi H, Ghods S, Hedjaroude GA. Immunomodulating and anticancer agents in the realm of macromycetes fungi. *Int. Immunopharmacol.* 2007;7:701-724.
5. Borchers AT, Krishnamurthy A, Keen CL, Meyers FJ, Gershwin ME. The immunobiology of mushrooms. *Exp. Biol. Med.* 2008; 233:259-276.
6. Nworu CS, Ihim SA, Okoye FBC, Esimone CO, Adikwu WU, Akah PA. Immunomodulatory and immunorestorative activities of β -D-glucan-rich extract and polysaccharide fraction of mushroom, *Pleurotus tuberregium*. *Pharm. Biol.* 2015;53:1555-1566.
7. Ruthes AC, Smiderle FR, Iacomini M. Mushroom heteropolysaccharides: A review on their sources, structure and biological effects. *Carbohydr. Polym.* 2016;136:358-375.
8. Babior BM, Woodman RC. Chronic granulomatous disease. *Semin. Hematol.* 1990;27:247-259.
9. Klaus A, Kozarski M, Vunduk J, Todorovic N, Jakovljevic D. Biological potential of extracts of the wild edible Basidiomycete mushroom *Grifola frondosa*. *Food Res. Int.* 2015;67:272-283.
10. Schwartz B, Hadar Y. Possible mechanisms of action of mushroom derived glucans on inflammatory bowel disease and associated cancer. *Ann. Transl. Med.* 2014;2:9.
11. Kozarski M, Klaus A, Niksc M, Jakovljevic D, Todorovic N, van Griensven LJLD. Antioxidative and immunomodulating activities of polysaccharides extracts of the medical mushrooms *Agaricus bisporus*, *Agaricus brasiliensis*, *Ganoderma lucidum* and *Phellinus linteus*. *Food Chem.* 2011;129:1667-1675.
12. Mohsin M, Negi PS, Ahmed Z. Determination of the antioxidant activity and polyphenol contents of wild Lingzhi or Reishi medicinal mushroom, *Ganoderma lucidum* (W. Curt. Fr.) P. Karst. (Higher Basidiomycetes) from central Himalayan hills of India. *Int. J. Med. Mushrooms.* 2011;13:535-544.
13. Mau J.-sai SY, Tseng YH, Huang SJ. Antioxidant properties of methanolic extracts from *Ganoderma tsugae*. *Food Chem.* 2005;93:641-649.

14. Hu H, Ahn N-S, Yang X, Lee Y-S, Kang K-S. *Ganoderma lucidum* extract induces cell cycle arrest and apoptosis in MCF-7 human breast cancer cell. *Int. J. Cancer* 2002;102:250-253.
15. Ferreira ICFR, Vaz JA, Vasconcelos MH. Compounds from wild mushrooms with antitumor potential. *Cancer Agents Med. Chem.* 2010;10:424-436.
16. Barbosa JR, dos Santos Freitas MM, da Silva Martins LH, de Carvalho RN. Polysaccharides of mushroom *Pleurotus* spp.: New extraction techniques, biological activities and development of new technologies. *Carbohydr. Polym.* 2020;229:115550.
17. Elsayed EA, El Enshasy H, Wadaan MAM, Aziz R. Mushrooms: A potential natural source of anti-inflammatory compounds for medical applications. *Mediat. Inflamm.* 2014;2014:805841.
18. Ruthes AC, Smiderle FR, Iacomini M. Mushroom heteropolysaccharides: A review on their sources, structure and biological effect. *Carbohydr. Polym.* 2016;136:358-375.
19. Stebbing J, Phelan A, Griffin I, Tucker C, Oechsle O, Smith D, Richardson P. COVID-19: Combining antiviral and anti-inflammatory treatments. *Lancet Infect. Dis.* 2020;20:400-402.
20. Lin LT, Hsu WC, Lin CC. Antiviral natural products and herbal medicines. *J. Tradit. Complement. Med.* 2014;4:24-35.
21. Atri NS, Sharma SK, Joshi R, Gulati A, Gulati A. Amino acid composition of five wild *Pleurotus* species chosen from North West India. *Euro. J. Biol. Sci.* 2012;4:31-34.
22. Carbonero ER, Ruthes AC, Freitas CS, Utrilla P, Galvez J, Silva EV. Chemical and biological properties of a highly branched β -glucan from edible mushroom *Pleurotus sajor-caju*. *Carbohydrate Polymers.* 2012;90:814-819.
23. Dashmuth SR, Shinde MN. Comparative study for antioxidant activity of cold and hot water extracts of *Pleurotus sajor-caju* and *Pleurotus florida*. *Int. J. Inno. Pharm. Sci. Res.* 2014;2:894-909
24. Kongkla K, Poemaim S. Total phenolic content and antioxidant activity of crude methanolic extract from five *Pleurotus* species. *Int. J. Agri. Tech.* 2016;12:1455-1462.

© 2022 Chawanoraset et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/87176>