



Effect of Nitrogen and Plant Growth Regulators on Nutrient Concentration and Uptake of Ajwain (*Trachyspermum ammi* L. Sprague)

**Sheeshpal Choudhary^{a*}, A. C. Shivran^{a#}, B.L. Dudwal^{a@}, Meena Choudhary^{a#},
Rajendra Jakhar^{a#} and Shiv Prasad Daroga^{a†}**

^a Department of Agronomy, Sri Karan Narendra College of Agriculture, Sri Karan Narendra Agriculture University, Jobner, Jaipur, Rajasthan-303329, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i1230962

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/84031>

Short Research Article

Received 05 January 2022

Accepted 01 March 2022

Published 05 April 2022

ABSTRACT

A field experiment was conducted during *rabi* season of 2016-17 entitled "Effect of nitrogen and plant growth regulators on nutrient concentration and uptake of ajwain (*Trachyspermum ammi* L. sprague)" at Agronomy farm, Sri Karan Narendra College of Agriculture, Jobner in Jaipur district of Rajasthan. The experiment comprising of total twenty treatment combinations consisting four levels of nitrogen (0, 30, 60, 90 kg ha⁻¹) and five levels of plant growth regulators (control, NAA 50 ppm at 40 day after sowing, NAA 50 ppm at 40 and 60 day after sowing, thiourea 500 ppm at 40 day after sowing, thiourea 500 ppm at 40 and 60 day after sowing) laid out in factorial randomized block design replicated thrice. The results showed that application of nitrogen up to 30 kg ha⁻¹ significantly higher nitrogen concentration in grain and straw over control. However, total nitrogen uptake was significantly increased with application of 60 kg N ha⁻¹ over lower levels, but remained at par with 90 kg N ha⁻¹. Foliar application of plant growth regulators significantly increased nitrogen concentration in grain, straw and total nitrogen uptake of ajwain over control.

^{*}Professor;

[@]Assistant Professor;

[#]Ph.D. Scholar;

[†]SRF;

*Corresponding author: E-mail: spkantwa@gmail.com;

Keywords: Ajwain; nitrogen and plant growth regulators.

1. INTRODUCTION

Ajwain is a belonging to the family Apiaceae. Ajwain is a annual herbaceous, aromatic, cross-pollinated plant with a white flower. Ajwain is widely diffused and cultivated throughout the world, including Iran, Pakistan, Afghanistan, India, and Europe, yet it is native to Egypt Shojaaddin et al [1]. Ajwain is commonly used to treat a variety of ailments in people and animals. Depending on the genotype or botanical kind, the seed contains 3 to 4% volatile oil. Besides volatile oil, the grains contain about moisture (8.9 percent), carbohydrate (38.6 percent), protein (15.4 percent), fat (18.1percent), crude fiber (11.9 percent) and minerals (7.1 percent). Phosphorus, potassium, calcium, iron, and sodium are significant minerals, whereas thiamine, nicotinic acid, carotene, and riboflavin are vital vitamins [2]. Thymol is used as an ingredient of deodorants, toothpastes, mouthwashes and many pharmaceutical preparations Malhotra and Vijay [3]. In India, during the year 2015-16, area under the ajwain crop was 24010 hectare and production was 17180 tonnes. India is the largest producer and exporter of the ajwain grain in the world. India exports ajwain to around 46 countries and the major importing countries are Pakistan, United States of America, Malaysia, Saudi Arabia, United Arab Emirates, Nepal, Canada and United Kingdom. In Rajasthan it is cultivated in the districts of Chittorgarh, Bhilwara, Udaipur, Rajsamand, Kota, Jhalawar covering an area of 11267 ha with production and productivity of 7666 tonnes and 680 kg/ha, respectively Anonymous [4].

2. MATERIALS AND METHODS

The experiment was conducted to study the effect of nitrogen and plant growth regulators on nutrient concentration and uptake of ajwain at Agronomy farm, Sri Karan Narendra College of Agriculture, Jobner, Jaipur Rajasthan during *rabi* season of 2016-17. The experimental soil was loamy sand with 8.2 pH and EC (1.25 ds/m), having low in organic carbon (0.14 percent), low available N (130.0 Kg ha⁻¹), medium in available P (18.9 Kg ha⁻¹), enough available K (175.6 kg ha⁻¹). The experiment consisted of 4 levels of nitrogen (0, 30, 60, 90 kg ha⁻¹) and 5 plant growth regulators (control, NAA 50 ppm at 40 day after sowing, NAA 50 ppm at 40 and 60 day after sowing, thiourea 500 ppm at 40 day after

sowing, thiourea 500 ppm at 40 and 60 day after sowing) in factorial randomized block design with 3 replications. A uniform dose of 40 kilogram phosphorus was applied through single super phosphate as basal and drilled about 5-7 cm deep through hand plough at the time of sowing. Nitrogen was applied through urea. One third nitrogen was applied at sowing as basal and remaining in 2 equal splits at 30 and 75 day after sowing as per treatments. The different weather parameters were recorded during crop growing period in the years. The maximum and minimum temperature recorded during *rabi* season were in the range of 20.4 to 43.4 °C and 2.8 to 25.1 °C in 2016-17. The total rainfall received during *rabi* season was 24.8 millimeter in 2016-17. The data were statistically analysed as per the procedure given by Panse and Sukhatme [5]. The crop was harvested at physiological maturity stage on 09 May, 2017.

3. RESULTS AND DISCUSSION

3.1 Effect of Nitrogen

An examination of data Table 1 revealed that application of nitrogen fertilization in crop resulted significant increase in nitrogen concentration in grain and straw of ajwain. Application of nitrogen up to 30 kg ha⁻¹ increase the nitrogen concentration in grain and straw by 14.74 and 43.93 percent and thus proved significantly superior over control. The nitrogen concentration in grain and straw increased dramatically with a non-significant difference as the nitrogen level increased. However, increasing nitrogen fertilisation levels resulted in a significant increase in total nitrogen uptake by crops. Nitrogen application up to 60 kg ha⁻¹ resulted in a total nitrogen intake of 62.58 kg N ha⁻¹, which was significantly greater than the control and 30 kg N ha⁻¹ but comparable to 90 kg N ha⁻¹. The magnitude of increase in total uptake was 14.84 and 67.59 percent over 30 kg ha⁻¹ and control with the application of nitrogen up to 60 kg ha⁻¹, respectively. The beneficial effect of nitrogen fertilisation on crop nitrogen content appears to be attributable to a better nutritional environment in both the root zone and the plant system. The availability of nitrogen nutrients in the crop root zone was increased due to an adequate supply of nitrogen early in the crop season. Boosted nutrient availability combined with enhanced metabolic processes at the cellular level may have increased nutrient

uptake and accumulation in different sections of the plant. Increased crop biomass output during harvest in terms of seed and straw yield, as well as increased nutrient concentrations, may have resulted in a large increase in nitrogen uptake by the crop as a result of nitrogen fertiliser. These results are in conformity with the findings of Patel et al. [6] in coriander and Naruka et al. [7] in

ajwain. Application of nitrogen 60 kg ha⁻¹ was found significantly higher over 30 kg N ha⁻¹ and control giving 10.87 and 28.38 percent higher grain yield, respectively. Further increase in level of nitrogen to 90 kg ha⁻¹ could not improve the grain yield of ajwain significantly over 60 kg N ha⁻¹. The similar results have also been reported by Muvel et al. [8] in ajwain (Table 2).

Table 1. Effect of nitrogen and plant growth regulators on nitrogen concentration in grain, straw, total nitrogen uptake of ajwain

Treatment	Nitrogen concentration (%)		Total nitrogen uptake (kg ha ⁻¹)
	Grain	Straw	
Nitrogen level (kg ha⁻¹)			
0	2.51	0.66	37.34
30	2.88	0.95	54.49
60	3.05	0.97	62.58
90	3.08	1.01	66.06
SEm±	0.07	0.02	1.44
CD (P=0.05)	0.20	0.06	4.12
Plant growth regulators			
Control	2.51	0.74	39.83
NAA 50 ppm spray at 40 day after sowing	2.86	0.91	53.55
NAA 50 ppm spray at 40 and 60 day after sowing	3.03	0.95	62.33
Thiourea 500 ppm spray at 40 day after sowing	2.94	0.93	55.58
Thiourea 500 ppm spray at 40 and 60 day after sowing	3.06	0.96	64.28
SEm±	0.08	0.02	1.61
CD (P=0.05)	0.22	0.07	4.60

Table 2. Effect of nitrogen and plant growth regulators on grain yield of ajwain

Treatment	Grain Yield (kg ha ⁻¹)
Nitrogen level (kg ha⁻¹)	
0	842
30	975
60	1081
90	1111
SEm±	25
CD (P=0.05)	70
Plant growth regulators	
Control	846
NAA 50 ppm spray at 40 day after sowing	975
NAA 50 ppm spray at 40 and 60 day after sowing	1084
Thiourea 500 ppm spray at 40 day after sowing	992
Thiourea 500 ppm spray at 40 and 60 day after sowing	1112
SEm±	27
CD (P=0.05)	79

3.2 Plant Growth Regulators

An investigation of data Table 1 revealed that foliar application of plant growth regulators significantly increased nitrogen concentration in grain and straw of ajwain over control, however all plant growth regulators remained at par to each other. The percent increase with thiourea 500 ppm spray at 40 and 60 day after sowing, NAA 50 ppm spray at 40 and 60 day after sowing, thiourea 500 ppm spray at 40 day after sowing, NAA 50 ppm spray at 40 day after sowing were 21.91, 20.71, 17.13 and 13.94 percent in grain and 29.72, 28.37, 25.67, 22.97 percent in straw over control, respectively. An examination of data further indicated that foliar application of plant growth regulators significantly increased total uptake of nitrogen over control, significantly higher total uptake of nitrogen (64.28 kg/ha) was obtained with foliar application of thiourea 500 ppm spray twice at 40 and 60 day after sowing, which was significantly higher to the tune of 15.65, 20.03 and 61.38 percent over thiourea 500 ppm spray at 40 day after sowing, NAA 50 ppm spray at 40 day after sowing and control, respectively, however, it remained at par with application of NAA 50 ppm spray at 40 and 60 day after sowing. Boosted nutrient availability combined with enhanced metabolic processes at the cellular level may have increased nutrient uptake and accumulation in different sections of the plant. Increased crop biomass output during harvest in terms of seed and straw yield, as well as increased nutrient concentrations, may have resulted in a large increase in nitrogen uptake by the crop as a result of nitrogen fertiliser. The similar results were also reported by Balai [9] and Meena [10] in coriander and Bochalía et al. [11] in fenugreek. Application of plant growth regulators significantly increased seed yield of ajwain over control Table 2. Significantly higher grain yield (1112 kg ha⁻¹) of ajwain was obtained with application of thiourea 500 ppm spray at 40 and 60 day after sowing, which was significantly higher to the tune of 12.09, 14.05 and 31.44 percent over thiourea 500 ppm spray at 40 day after sowing, NAA 50 ppm spray at 40 day after sowing and control, respectively, however, it remained at par with application of NAA 50 ppm spray at 40 and 60 day after sowing. The result corroborate with the finding of Bairwa and Kaushik [12] and Gour et al. [13] in fenugreek.

4. CONCLUSION

Based on one years of investigation, it may be concluded that application of nitrogen up to

30 kg ha⁻¹ significantly higher the nitrogen concentration in grain and straw over control. Foliar application of plant growth regulators significantly increased nitrogen concentration in grain, straw and total nitrogen uptake over control.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Shojaaddini M, Moharramipour S, Sahaf B. Fumigant toxicity of essential oil from *Carum copticum* against Indian meal moth, *Plodia interpunctella*. *Journal of Plant Protection Research*. 2008;48(4):411–19.
2. Pruthi JS. Minor spices and condiments, Crop management and post-harvest technology. Published by publication and Information Division, ICAR, Krishi Anusandhan Bhavan, Pusa, New Delhi-110012. 2001;124-133.
3. Malhotra SK, Vijay OP. II Particular herbs and spices: Ajwain. In: Peter, K.V.(ed.) *Handbook of Herbs and Spices*. Woodhead Publishing Ltd., Cambridge, UK. 2004;2:107-116.
4. Anonymous. *Vital agriculture statistics*. Directorate of Agriculture, Pant Krishi Bhawan, Jaipur (Rajasthan); 2018-19.
5. Panse VG, Sukhatme PV. *Statistical methods for Agricultural Workers*, ICAR Publication, New Delhi. 1985;361.
6. Patel CB, Amin AU, Patel AL. Effect of varying levels of nitrogen and sulphur on growth and yield of coriander (*Coriandrum sativum* L.). *The Bioscan*. 2013;8(4):1285-1289.
7. Naruka IS, Singh PP, Megha Barde, Rathore SS. Effect of spacing and nitrogen level on growth, yield and quality of ajwain (*Trachyspermum ammi* L. Sprague). *International Journal Seed Spices*. 2012;2(1):12-17.
8. Muvel R, Naruka IS, Chundawat RS, Shaktawat RPS, Rathore SS, Verma R. Production, productivity and quality of ajwain (*Trachyspermum ammi* L. Sprague) as affected by plant geometry and fertilizer levels *International Journal Seed Spices*. 2015;5(2):32-37.
9. Balai LR. Effect of thiourea on growth and productivity of coriander (*Coriandrum sativum* L.) varieties under normal and late

- sown conditions. Ph.D. (Ag.) Thesis, Rajasthan Agricultural University, Bikaner; 2005.
10. Meena SK. Effect of Plant Growth Regulators and Sulphur on Productivity of Coriander (*Coriandrum sativum* L.). M.Sc. (Ag.) Thesis, Swami Keshwanand Rajasthan Agricultural University, Bikaner; 2011.
 11. Bochalia GC, Tiwari RC, Ram B, Kantwa SR, Choudhari AC. Response of fenugreek (*Trigonella foenum-graecum* L.) genotypes to planting geometry, agro-chemicals and sulphur levels. Indian Journal of Agronomy. 2011;56(3):273-279.
 12. Bairwa RC, Kaushik MK. Effect of fertility levels and growth regulators on yield and yield attributes of fenugreek (*Trigonella foenum-graecum* L.) varieties. Research on Crops. 2007;8(2):355-360.
 13. Gour K, Patel BS, Mehta RS. Growth, yield and profitability of fenugreek (*Trigonella foenum-graecum* L.) as influenced by varying levels of growth regulators and vermin-wash. Journal of Spices and Aromatic Crops. 2010;19(1&2):65-67.

© 2022 Choudhary et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/84031>