



Efficacy of Newly Emerging Detergents, Laundry Bleach, and Toilet Soap against Bacteria Isolated from Fairly used Clothes, Male/Female Underwear

O. T. Osuntokun^{a*}, B. O. Adu^a, O. A. Thonda^b and O. M. Aladejana^b

^a Department of Microbiology, Faculty of Science, Adekunle Ajasin University, Akungba Akoko, Ondo State, Nigeria.

^b Department of Biological Science, Microbiology Unit, Kings, University, Odeomu, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/SAJRM/2022/v12i330273

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/86011>

Original Research Article

Received 08 February 2022

Accepted 16 April 2022

Published 26 April 2022

ABSTRACT

The aim of this study is to determine the potency of newly emerging detergents, laundry bleach, and toilet soap on bacteria isolates from fairly used clothes and Female Underwear. The current study included sample collection from used clothing and female underwear and compared the efficacy of detergents, toilet soap and laundry bleach (Jik), to reduce the microbial load in the used cloth. Organisms were isolated from the used cloth and female underwear, identified, and microbiologically characterized using the conventional method of microbiology assay. i.e, biochemical test, a compilation of cultural colonial and characteristics. Test Samples were washed using four (4) types of detergents and five (5) types of toilet soap and laundry bleach. The results of the colony-forming unit (CFU) before and after washing with detergent and toilet soap and laundry bleach were recorded, washing with detergents and toilet soap. The results showed that detergents were more effective in reducing bacterial load than toilet soap in all test samples. the addition of laundry bleach (Jik) also shows the effectiveness in reducing bacterial load in fairly used clothes and Female Underwear The antibiotics susceptibility test were determined using disc diffusion and agar well diffusion methods respectively. Eight (8) bacteria were identified include *Aerobacter aerogenes* (3), *Bacillus polymyxa* (5), *Staphylococcus aureus* (2), *Veillonella parvula* (5), *Escherichia freundii* (1), *Escherichia coli* (3), *Enterobacter aerogenes* (2) and *Aerobacter cloacae* (1). *Bacillus polymyxa*, and *Veillonella parvula* were the most common organism isolated and

*Corresponding author: Email: osuntokun4m@gmail.com, osuntokun4m@yahoo.com;

followed by *Escherichia coli* from fairly used clothes and Female Underwear. In this study, some of the Gram-positive(+ve) bacteria isolates were sensitive to commonly used antibiotics like Streptomycin, Tarivid, Reflacin, Ciprofloxacin, Augmentin, Gentamycin, Ceporex, Nalidixic acid, Septrin, and Amplicin. Gram-negative bacteria isolated were sensitive to Ciprofloxacin and all the isolated organisms were resistant to Ceporex. Gram-positive (-ve) bacteria were sensitive to Streptomycin and were resistant to Rifampicin. To determine the growth dynamics and killing kinetics of the isolated organisms, from fairly used clothes and Female Underwear, an Ultraviolet spectrophotometer were used to determine the wavelength and killing kinetics of isolated organisms from fairly used clothes and Female Underwear. The addition of antibiotics to the isolated organisms at the 48th-hour speed up the death rate of the isolates from fairly used clothes washed with different detergent Laundry Bleach and toilet soaps. It can be deduced that this study validates the efficacy of newly emerging detergents, Laundry bleach, and toilet soap against bacteria isolated from fairly used clothes. It is also recommended to wash fairly used clothes using different methods of soaking the fairly used clothes and female underwear in the Laundry Bleach for some days before finally washing in detergents and ironing before use. This method will drastically reduce and remove bacterial load in the fairly used clothes and female underwear.

Keywords: *Efficacy; newly emerging detergents; toilet soap; bacteria; fairly used clothes; female under-wear.*

1. INTRODUCTION

Fairly used clothes in Nigeria are known as “Okrika” and are clothing materials like shirts, trousers, towels, socks, panties, pillowcases, curtains, and bed sheets imported from the United States of America, United Kingdom, Asia, Germany, and the Netherlands after been used for a while by the initial buyer [1]. They are packed in bundles known as bales during importation. Nigerians’ desire for foreign goods (including wears) increases daily because of our economic situation and poverty under-tunes. These clothes were imported commonly from Asia and Europe, the largest exporter of second fairly used clothes were the united states of America (U.S.A) followed by Germany, the United Kingdom (U.K), and the Netherland while the largest importers of fairly used clothes are Sub-Sahara African, South East Asia and Eastern Europe [1].

Though the wears were cheaper, usually the driving force is the quality of the products which are adjudged better than the new ones, an erroneous belief. These were purchased without giving consideration to any attending health and risk implications. Clothes have the potential, just as any other hand contact site, to be a component in the chain of infection transmission during normal daily activities (2).

However, Some of the risks associated with fairly used clothing consist of; Tinea, Impetigo, Scabies, Body lice, Pubic lice, Head lice are

possible diseases that could be transported through fairly used clothes and their tendency to spread skin diseases.

Overtime man has been fighting for several centuries with the ever-persistent bacterial infections. The threat of bacterial infection was threatening with high mortality rates. This type of predicament which was faced by patients and also put the credibility of doctors to the task made the scientists of that time find ways to solve this problem [3].

At the turn of the century with the ever-growing number of disinfectants claiming to possess anti-bacterial properties, it has been left to the consumer to decide for oneself which is best. In the light of disinfectant used for cloth, availability, efficacy, and potency is our watchword. Local entrepreneurs haven’t been left behind by the “clean bus” and there are quite a number of products produced each with its own testimony of how effective they are against bacteria. Through this myriad of confusion, the consumer is the one to suffer most of our laundry but does not disinfect [4]. Increased life expectancy since the first half of the 20th century can be attributed to improved hygiene status resulting in decreased infectious disease incidence. Simple personal hygiene including soap utilization was the silent success of public health in the pre-disinfectant era [5].

In our day-to-day interactions with microorganisms, we employ a variety of ways to reduce otherwise their numbers so as to avoid

contamination with the microbes leading to eventual infection and illness. One major way of achieving these goals is the use of various types of disinfectants and detergents which are being sold in the market. Various types of disinfectants being sold have various activities ranging from –cidal to –static activities [6].

A detergent is a surfactant or a mixture of surfactants with cleansing properties when in dilute solutions. Disinfectants are specifically designed to kill bacteria, viruses, and mildews that are found on clothes and even kitchen and bathroom fixtures, and flourish on counters and drain gates in the sink [7]. Using disinfectants is important in households because they prevent the spread of bacteria [6].

Detergent is used to refer to synthetic cleaning compounds distinguishable from soap even though soap is also a detergent in its true sense. There are a large variety of detergents used in Nigeria, their potency should be questioned but the most commonly found are alkylbenzene sulfonates, a family of compounds that are similar to soap but are more soluble in hard water because the polar sulfonate (of detergent) is less likely than the polar carboxylate (soap) to bind to calcium and other ions found in hard water [8].

In domestic contexts, the term detergent by itself refers specifically to laundry detergent or dish detergent, as opposed to hand soap or other types of cleaning agents. Detergents, like soaps, work because they are amphiphilic; partly hydrophilic (polar), and partly hydrophobic (non-polar). Their dual nature facilitates the mixture of hydrophobic compounds (like oil and grease) with water. Because air is not hydrophilic, detergents are also foaming agents to varying degrees [9]. Toilet Soap on the other hand is a salt of a fatty acid used in a variety of cleansing and lubricating products. Soap when dissolved in water, possesses the ability to remove dirt from surfaces such as the human skin, textiles, and other solid. In a domestic setting, soaps are surfactants usually used for washing, bathing, and other types of housekeeping. In industrial settings, soaps are used as thickeners, components of some lubricants, and precursors to catalysts [10,11].

Triclosan one of the basic microbiological active ingredient of common soap and detergent has been shown to inhibit the growth of both Gram-positive and Gram-negative bacteria *in situ*, with varying effectiveness across bacterial species.

For, example, triclosan is relatively ineffective at inhibiting the growth of Gram-negative bacteria [12]. Although the antibacterial activity of triclosan involves some non-specific killing mechanisms, research finding suggests that the bacteriostatic action occurs by inhibiting a specific bacterial target, known as the enoyl-acyl carrier protein reductase. Triclosan shares this bacterial biosynthetic fatty acid pathway target with the antibiotic isoniazid [13,14,15].

Several investigators have reviewed studies examining the mechanisms of antiseptic resistance [13,14,16], there are few systematic reviews that have attempted to summarize the potential risks associated with triclosan in the context of the purported effectiveness of this antibacterial ingredient used in hygiene products in the community setting [16]. The efficacy of soap containing triclosan generally refers to the additional level of effectiveness beyond the ability of plain soap to simply remove transient organisms via surfactants and the mechanical action of the wash procedure.

The level of effectiveness may be measured at the microbiological level or at the population level, as added protection against bacterial contamination or the occurrence of common infectious illnesses. Risks, on the other hand, include the potential for bacteria to become unsusceptible to triclosan, for the emergence of cross-resistance to antibiotics, and for the ingredients to become toxic to the environment and to humans [16].

2. MATERIALS AND METHODS

2.1 Detergents, Laundry Bleach and Toilet Soap Used

Toilet soap used (4): - Test sample 1(Premier cool), Test sample 2 (Eva), Test sample 3(Lux), Test sample 4(Joy, Detergents used(4): - Test sample 1(Waw), Test sample 2(Canoe), Test sample 3(Good mama), Test sample 4(So easy). (Laundry Bleach [1] Jik.. All detergents, Laundry Bleach (Jik), and toilet soap used for this research were purchased from a Market in Akungba Akoko, Ondo State, Nigeria.

2.2 Sample Collection - Fairly used Clothes, Male and Female under Wear

Two categories of fairly used cloth samples, Towel, and Singlet were procured in May 2021

from Akungba Akoko market, located in Ondo State, Nigeria. The two categories of fairly used clothes, male and female underwear were labeled accordingly. The samples were then placed in sterile polythene bags and brought to the Microbiology Laboratory of Adekunle Ajasin University, Akungba-Akoko for analysis. Standard methods were used in the isolation and identification of the microorganisms.

2.3 Isolation of Microorganisms from Test Samples

To obtain microorganisms from fairly used clothes, male and female underwear (Before and after washing); the test tube was partly full of sterilized water. Dipped bacteria-laden swabs into the water, then transferred some of the bacteria collected into the water. To ensure maximum representation of the microorganisms in the samples during culturing and to obtain good results, a test run is were determined with the concentrated extract (After washing) and the raw sample (Before washing). Inoculate the dish by pouring 0.5m/L of the water into the dish and pouring already sterilized and cooled nutrient agar on it, then incubated 37°C for 24hrs [17].

2.4 Culturing of Isolate from Test Samples

Culturing of isolates from fairly used clothes, male and female underwear i,e test sample. Pour plate methods were used to inoculate the test sample. The technique is carried out for each of the sample extracts by inoculating 0.1 to 1 mL onto the dry agar plate and already prepared cooled sterile 20mL of nutrient agar was poured on it and are then incubated at room temperature and results were obtained after 24 hours.(18).

2.5 Bacteriological Analysis of Fairly used Clothes, Male and Female Underwear

Using the pour plate method of inoculation,0.5ml of the five-fold dilution of 10^5 and 10^3 of fairly used clothes, male and female underwear waters which has already been washed with toilet soap, laundry bleach, and detergents was aliquoted into sterile Petri dishes. The nutrient medium was prepared by dissolving 6.16grams of Agar into

220 liters of distilled water in a sterile conical flask, corked with cotton wool and aluminum foil, and then homogenized to dissolve. It was sterilized in an autoclave at a temperature of 121 for 15minutes, after the sterilization the medium was allowed to cool but not solidify, 20ml amount of the medium was then poured into different sterile Petri dishes containing 0.5ml of the inoculum and allowed to set, and then the plates were incubated for 24hrs. After 24hrs, the cultural characteristics on the plates were studied and recorded; resultant colonies were sub-cultured on fresh nutrient agar and then incubated for 24hrs. Pure isolates were preserved on a double-strength nutrient agar slant for further studies. [18].

2.6 Identification of Isolates from Fairly used Clothes, Male and Female Underwear

2.6.1 Macroscopic Examination and Biochemical Characteristics of Isolates

Cultural and macroscopic examinations were done to identify the pure isolate. identification of the isolates was based on the cellular morphology characteristics. In addition, various biochemical tests which include: Catalase, Indole, Motility, Gram staining, Fermentation of sugars (Sucrose, Lactose, Dextrose), Coagulate, Urease, Hydrogen sulphide, Gas production, and oxidase tests were done for conventional identification of the isolates [18].

2.6.2 Microscopic Examination and Gram Staining Techniques of Isolates

Gram staining is a common technique used to differentiate two large groups of bacteria based on their different cell constituents. The Gram stain procedure distinguishes between Gram-positive and Gram-negative groups by coloring these cells red or violet. The working solution of reagents used for the Gram staining technique was prepared according to the manufacturer's instructions. Staining was carried out by emulsifying approximately one isolated 18-24hours old colony in a drop of water placed at the center of a clean grease-free slide until a thin smear was made. The smear was air heat-fixed by passing the slide through a Bunsen burner flame and then air-dried. The heat-fixed smear was flooded with a basic aniline dye (crystal violet) for 60 seconds. This was flooded with

Lugol's iodine and allowed to remain for 60 seconds. This was then rinsed off with running tap water. The smear was decolorized with 70% ethanol which was immediately washed out to avoid total decolorization. The smear was counter-stained with Safranin for 60 seconds, washed off with running tap water, and blot-dried. The slide was then examined under an oil immersion objective microscope. Organisms that retained the purple colour of crystal violet- iodine complex (CV-1 complex) were recorded as Gram-positive, while those that appeared pink were Gram-negative. The slide was then air-dried and the bottom was dried. The slide was then observed under a compound microscope using an oil emulsion lens at a magnification of X100 [19,20,21].

2.6.3 Antibiotics sensitivity testing of isolates from fairly used clothes, male and female underwear

The antibiotics sensitivity test was performed in order to determine the sensitivity or resistance of the bacteria isolates to some selected antibiotics of certain concentrations. Kirby-Bauer National Committee for Clinical and Laboratory Standard (CLSI, 2014) modified disc diffusion technique was used to determine the antibiotics sensitivity of the isolates. Using a sterile inoculating loop, a colony of the organisms was taken and inoculated into a Nutrient agar broth in the test tubes and inoculated for 24hrs. A sterile swab stick was dipped into the test tubes to pick the organisms and the swab stick was rotated severally and pressed firmly on the side of the wall to remove excess inoculum and used to streak the entire dried surface of the already prepared, sterilized, and solidified nutrient agar in the Petri dishes, the streaking was repeated two (2) times by rotating the plate approximately each time to ensure that every part of the plate was inoculated. The plates were allowed to stay for 15minutes so as to allow any excess surface moisture to be absorbed before applying the drug impregnated discs. Pre-determined commercial (Oxoid), antibiotics disc of both gram-positive and gram-negative was fixed on the media using sterile forceps to ensure complete contact with agar. The plates were inverted and incubated

overnight at 37°C for 24hrs. After 24hrs of incubation, the plates were examined; the susceptibility of each isolate to each antibiotic was indicated by a clear zone of growth of inhibition around each disc. The diameter of the zone of inhibition was measured using a calibrated ruler from the underside of the plate and recorded in millimeters. The results were interpreted according to the Clinical laboratory standard institute [22]. The following antibiotics used were; For gram positive (+ve) Streptomycin (S,30mcg), Norfloxacin (NB, 10mcg), Chloramphenicol (CH, 30mcg), Ciprofloxacin (CPX, 10mcg), Erythromycin (E,30mcg), Levofloxacin (LEV, 20mcg), Gentamycin (CN, 10mcg), Ampiclox (APX, 20mcg), Rifampicin (RD, 20mcg), Amoxicillin (AMX, 20mcg) For gram negative (-ve) Tarivid (OFX, 10mcg), Reflacin (PEP,10mcg), Ciprofloxacin (CPX, 10mcg), Augmentin(AU, 30mcg,) Gentamycin (CN, 10mcg), Streptomycin (S, 30mcg), Ceporex (CEP, 10mcg), Nalidixic acid (NA,30mcg), Septrin (SXT, 30mcg), Ampicillin (PN, 30mcg). Each disc was pressed down to ensure complete contact with agar surface and incubated at 37°C for 24hrs. The zones of inhibition were measured using calibrated ruler in millimeters [18].

2.6.4 Growth dynamic / death rate(killing kinetics) of isolates using ultra violet spectrophotometer

Growth dynamic refers to the rate at which cells of microorganisms grow at a given time. The test were done to determine the rate of growth of the isolates as well as their killing time in due time. The colony were picked from the stocked culture slant and inoculated into a nutrient broth which was incubated for 37°C for 24hrs. A loopful of organisms were picked from the broth culture into nutrient broth in two sets which are set A and C respectively. The ultraviolet spectrophotometer was set at 480 wavelengths, warmed up for 15minutes and then the control were first read, the first reading were taken at zero hours and it continues after every 12hours for 8times, at the 5th reading which is the 48th hour of set B, ciprofloxacin was added to determine up the rate of the kill [23].

Table 1. Cultural Characteristics (Macroscopic Examination) of Isolated Bacteria from Fairly Used Clothes, Male /Female Underwear against Newly Emerging Detergents, Laundry Bleach and Toilet Soap

Newly Emerging Detergents, Laundry Bleach and Toilet Soap Test sample	Dilution Factor	Size	Shape	Colour	Texture	Opacity	Edges
Test sample 1 (Premier cool+jik)	10 ⁵	Big	Irregular	Creamish	Rough	Translucent	Rough
Test sample 1 (Premier cool+jik)	10 ³	Small	Regular	Creamish	Smooth	Opaque	Rough
Test sample 1 (Waw+jik)	10 ³	Small	Regular	Yellowish	Smooth	Translucent	Smooth
Test sample 1 (Waw+jik)	10 ⁵	Big	Regular	Milky	Smooth	Translucent	Smooth
Test sample 1(Premier cool)	10 ⁵	Small	Regular	Creamish	Smooth	Opaque	Smooth
Test sample 1(Premier cool)	10 ³	Small	Regular	Brownish	Rough	Opaque	Smooth
Test sample 1(Waw)	10 ⁵	Small	Regular	Whitish	Smooth	Opaque	Smooth
Test sample 1(Waw)	10 ³	Medium	Irregular	Golden yellow	Smooth	Translucent	Rough
Test sample 2 (Eva)	10 ⁵	Big	Irregular	Creamish	Smooth	Translucent	Smooth
Test sample 2 (Eva)	10 ³	Small	Irregular	Whitish	Smooth	Transparent	Smooth
Test sample 2 (Eva+jik)	10 ³	Medium	Regular	Yellowish	Smooth	Translucent	Smooth
Test sample 2(Canoe +jik)	10 ⁵	Big	Irregular	Creamish	Rough	Transparent	Smooth
Test sample 2(Canoe)	10 ⁵	Big	Irregular	Creamish	Smooth	Translucent	Smooth
Test sample 2(Canoe)	10 ³	Small	Regular	Whitish	Smooth	Opaque	Smooth
Test sample 2(Canoe+jik)	10 ³	Small	regular	Whitish	smooth	Opaque	Smooth
Test sample 3 (Good mama)	10 ⁵	Medium	Regular	Whitish	Smooth	Transparent	Smooth
Test sample 3 (Good mama+jik)	10 ³	Small	Regular	Yellowish	Rough	Translucent	Smooth
Test sample 3 (Lux+jik)	10 ⁵	Small	Regular	Whitish	Smooth	Opaque	Smooth
Test sample 3(Lux)	10 ³	Big	Regular	Yellowish	Smooth	Transparent	Smooth
Test sample 3(Lux)	10 ⁵	Medium	Irregular	Milky	Rough	Opaque	Rough
Test sample 3(Lux+jik)	10 ³	Small	Regular	Whitish	Smooth	Opaque	Smooth
Test sample 4 (Joy)	10 ⁵	Small	Irregular	Creamish	Rough	Transparent	Rough
Test sample 4 (So Easy+jik)	10 ³	Medium	Regular	Milky	Smooth	Transparent	Smooth
Test sample 4(Joy+jik)	10 ⁵	Big	Irregular	Creamish	Rough	Opaque	Rough
Test sample 4(So Easy)	10 ³	Medium	Regular	Creamish	Smooth	Transparent	Smooth

Key –NEWLY EMERGING DETERGENTS, LAUNDRY BLEACH AND TOILET SOAP

- 1.Toilet soap used (4): - Test sample 1(Premier cool), Test sample 2 (Eva) Test sample 3(Lux), Test sample 4(Joy),
- 2.Detergents used(4): Test sample 1(Waw), Test sample 2(Canoe), Test sample 3(Good mama), Test sample 4(So easy)
- 3.Laundry Bleach (1) Test sample (1) Jik.

3. RESULTS

The result obtained from this research work, the efficacy of newly emerging detergents, laundry bleach, and toilet soap against bacteria isolated from fairly used clothes, male and female underwear were collated and represented in the table and graphical forms.

Table .1; Shows cultural characteristics (Macroscopic examination) of isolated bacteria from fairly used clothes, male and female underwear washing with detergents, laundry bleach and toilet soaps. Cultural characteristics include the size, shape, colour, texture, opacity and edges. It was observed that some isolated bacteria were big, small, and medium in size, irregular and regular in shape, rough and smooth in texture, transparent, translucent and opaque in opacity, rough or smooth edges and the bacteria also exhibit different colour; whitish, orange, yellowish and creamy in colour.

1. Samples washed with Toilet soap (Test sample 1(Premier cool), Detergent (Test sample 1(Waw) with the addition of laundry bleach(Test sample 1(Jik), Test sample 4(Joy) with laundry bleach(Test sample 1(Jik), Test sample 2(Canoe) with laundry bleach(Test sample 1(Jik), Test sample 2(Canoe) and Test sample 2 (Eva).has a large-sized isolates.

2.Samples washed with Test sample 2 (Eva), Test sample 2(Canoe), Test sample 3(Lux) with laundry bleach(Test sample 1(Jik), Test sample 2(Canoe), with laundry bleach(Test sample 1 (Jik), Test sample 3(Good mama), with laundry bleach(Test sample 1(Jik), Test sample 4(Joy), Test sample 1(Premier cool) with laundry bleach(Test sample 1(Jik), Test sample 1(Waw), Test sample 1(Premier cool), has small-sized isolates.

Table 2; Shows the morphological characteristics(Macroscopic examination) of isolated bacteria from fairly used clothes, male and female underwear. In this table it was observed that the isolated bacteria has different morphological characteristics; transparent, translucent and opaque opacity, rough with creamy, whitish, milky, and yellowish in colour.

Table 3; Shows the result of Gram staining of the isolates from isolated bacteria from fairly used clothes, male and female underwear. It was observed from this table, all isolate after washing

with different detergent and toilet soap, were the result obtained.

Test sample 3 (Lux- 1×10^5 B), Test sample 3(Lux)+ laundry bleach(Test sample 1(Jik- 1×10^3), Test sample 1(Premier cool)+ laundry bleach(Test sample 1(Jik - 4×10^3 A), Test sample 1(Premier cool)+ laundry bleach(Test sample 1(Jik - 4×10^5 B), Test sample 1(Waw) + laundry bleach(Test sample 1(Jik - 2×10^5), Test sample 2(Canoe- 1×10^3), Test sample 4(Joy)+ laundry bleach(Test sample 1(Jik 8×10^5), Test sample 3(Good mama),+ laundry bleach(Test sample 1(Jik- 5×10^3), Test sample 3(Good mama - 2×10^5), Test sample 1(Waw)+ laundry bleach(Test sample 1(Jik- 6×10^3 A), Test sample 1(Waw- 3×10^5), Test sample 1(Waw)+ laundry bleach(Test sample 1(Jik- 6×10^5 B), Test sample 4(So easy)+ laundry bleach(Test sample 1(Jik - 7×10^3), Test sample 2 (Eva)+ laundry bleach(Test sample 1(Jik- 3×10^3) were negative to Gram staining, and Test sample 1(Premier cool- 4×10^5), Test sample 1(Premier cool + laundry bleach(Test sample 1(Jik- 4×10^5), Test sample 2.

(Eva- 3×10^3), Test sample 3(Lux- 1×10^3 A), Test sample 4(So easy- 4×10^5), Test sample 1(Waw- 3×10^3), were all Gram-positive.

Microscopic examination shows that all isolate obtained after washing with; Test sample 3(Lux- 1×10^5 B), Test sample 3(Lux)+ laundry bleach(Test sample 1(Jik- 1×10^3), Test sample 1(Waw) ,+ laundry bleach(Test sample 1(Jik- 6×10^5 B), Test sample 1(Premier cool- 4×10^5), Test sample 1(Premier cool) + laundry bleach(Test sample 1(Jik- 4×10^5 B) were Tiny rods shape, Test sample 1(Premier cool)+ laundry bleach(Test sample 1(Jik - 4×10^3 A), Test sample 4(Joy) + laundry bleach(Test sample 1(Jik - 8×10^5), Test sample 3(Good mama)+ laundry bleach(Test sample 1(Jik - 5×10^3), Test sample 2 (Eva- 3×10^3), Test sample 3(Lux- 1×10^3 A), Test sample 4(So easy- 4×10^3) has Long rods shape.

Test sample 3(Good mama - 2×10^5), Test sample 1(Waw)+ laundry bleach(Test sample 1(Jik- 6×10^3 A) has Short rods shape.

Test sample 1(Premier cool)+ laundry bleach(Test sample 1(Jik- 4×10^5 B), Test sample 2(Canoe - 1×10^3), Test sample 1(Waw - 3×10^5 B), Test sample 4(So easy)+ laundry bleach(Test sample 1(Jik- 7×10^3), Test sample 2 (Eva)+ laundry bleach(Test sample 1(Jik- 3×10^3), Test sample 1(Waw- 3×10^3 A) were all cocci in shape.

Table 2. Morphological Characteristics (Macroscopic Examination) of Isolated Bacteria from Fairly Used Clothes, Male /Female Underwear against Newly Emerging Detergents, Laundry Bleach and Toilet Soap

Newly Emerging Detergents, Laundry Bleach and Toilet Soap Test sample	Dilution Factor	Sub-Culture
Test sample 1 (Premier cool+jik)	10 ³	Swamy, opaque with creamy colour
Test sample 1 (Waw+jik)	10 ³	Motile, translucent with golden yellow colour
Test sample 1 (Waw+jik)	10 ⁵	Motile, translucent with milky colour
Test sample 1(Premier cool)	10 ⁵	Non-motile, opaque with creamy colour
Test sample 1(Premier cool)	10 ³	Swamy, opaque with brownish colour
Test sample 1(Premier cool+jik)	10 ⁵	Motile, translucent with creamy colour
Test sample 1(Waw)	10 ⁵	Motile, opaque with whitish colour
Test sample 1(Waw)	10 ³	Motile, translucent with golden yellow
Test sample 2 (Eva)	10 ⁵	Non-motile, translucent with creamy colour
Test sample 2 (Eva)	10 ³	Motile, transparent with whitish colour
Test sample 2 (Eva+jik)	10 ³	Motile, translucent with yellowish colour
Test sample 2(Canoe)	10 ⁵	Non-motile, translucent with creamy colour
Test sample 2(Canoe)	10 ³	Motile, opaque with whitish colour
Test sample 2(Canoe+jik)	10 ³	Swamy, opaque with creamy colour
Test sample 2(Canoe+jik)	10 ⁵	Swamy, transparent with creamy colour
Test sample 3 (Good mama)	10 ⁵	Motile, transparent with whitish colour
Test sample 3 (Good mama+jik)	10 ³	Motile, translucent with brownish colour
Test sample 3 (Lux+jik)	10 ⁵	Non-motile, opaque with whitish colour
Test sample 3 (Lux+jik)	10 ³	Motile, opaque with whitish colour
Test sample 3(Lux)	10 ³	Non-motile, transparent with yellowish colour
Test sample 3(Lux)	10 ⁵	Swamy organism, opaque with creamy colour
Test sample 4 (Joy)	10 ⁵	Swamy, translucent with creamy colour
Test sample 4 (So Easy+jik)	10 ³	Motile, transparent with milky colour
Test sample 4(Joy+jik)	10 ⁵	Motile, opaque with creamy colour
Test sample 4(So Easy)	10 ³	Motile, transparent with creamy colour

Table 3. Gram staining for identification of Isolated Bacteria from Fairly Used Clothes, Male/Female Underwear (Towel, Singlet and Panties) against Newly Emerging Detergents, Laundry Bleach and Toilet Soap

Newly Emerging Detergents, Laundry Bleach and Toilet Soap Test sample	Gram Staining	Microscopy Examination
Test sample 1 (Premier cool+Jik)4x10 ³ A	Long irregular rods	-ve
Test sample 1 (Premier cool+Jik)4x10 ⁵ B	Cluster cocci	-ve
Test sample 1(Premier cool) 4x10 ⁵	Tiny irregular rod	+ve
Test sample 1(Waw)3x10 ³ A	Cluster cocci	+ve
Test sample 1(Waw)3x10 ⁵ B	Cluster cocci	-ve
Test sample 1(Waw+Jik) 6x10 ³ A	Short rods	-ve
Test sample 1(Waw+Jik) 6x10 ⁵ B	Tiny irregular rods	-ve
Test sample 2 (Eva)3x10 ³	Long irregular rod	+ve
Test sample 2 (Eva+Jik)3x10 ³	Cluster cocci	-ve
Test sample 2(Canoe)1x10 ³	Irregular cocci	-ve
Test sample 3 (Good mama)2x10 ⁵	Short rods	-ve
Test sample 3 (Good mama+Jik)5x10 ³	Long irregular rods	-ve
Test sample 3 (Lux+Jik)1x10 ³	Tiny irregular rods	-ve
Test sample 3(Lux)1x10 ³	Long irregular rods	+ve
Test sample 3(Lux)1x10 ⁵	Short rods	-ve
Test sample 4 (So Easy+Jik)7x10 ³	Cocci	-ve
Test sample 4(Joy+Jik) 8x10 ⁵	Long rods	-ve
Test sample 4(So Easy) 4x10 ³	Long irregular rods	-ve

KEY: +(POSITIVE) - (NEGATIVE)

Table 4; Shows the biochemical tests carried out on the isolates. The tests include; the Motility test, Urease test, Indole test, Triple sugar fermentation (Sucrose, Lactose, Dextrose), Gas production, and Hydrogen sulphide test (H_2S). It was observed in this table that some isolates are positive and negative to motility test, it was observed that the isolates from different samples of newly emerging detergents, laundry bleach, and toilet soap, were the following observation, Test sample 3(Lux- 1×10^5 B), Test sample 3(Lux - 1×10^3 A), Test sample 1(Premier cool)+ Test sample (1) laundry bleach(Test sample 1(Jik- 4×10^5), Test sample 4(So easy) + laundry bleach(Test sample 1(Jik- 7×10^3 , Test sample 4(So easy - 4×10^3), Test sample 1(Waw - 3×10^3 A), Test sample 1(Waw- 3×10^5 B), Test sample 1(Premier cool + laundry bleach (Test sample 1(Jik- 4×10^3), Test sample 1(Premier cool - 4×10^5), Test sample 2 (Eva) + laundry bleach (Test sample 1(Jik- 3×10^3), Test sample 2 (Eva)- 3×10^5), Test sample 1(Waw + laundry bleach(Test sample 1(Jik - 6×10^3), Test sample 3(Good mama - 2×10^5). Test sample 3(Good mama + laundry bleach(Test sample 1(Jik- 5×10^3), Test sample 2(Canoe - 1×10^3), Test sample 4(Joy + laundry bleach(Test sample 1(Jik - 8×10^5) were Urease positive while, Test sample 3(Lux + laundry bleach(Test sample 1(Jik - 1×10^3) were Urease negative. It was also observed that Test sample 3(Lux- 1×10^5 B), Test sample 4(So easy + laundry bleach(Test sample 1(Jik - 7×10^3), Test sample 1(Waw- 3×10^3 A), Test sample 1(Waw - 3×10^5 B), Test sample 3 (Lux + laundry bleach(Test sample 1(Jik - 1×10^3), Test sample 1(Premier cool + laundry bleach(Test sample 1(Jik - 4×10^3), Test sample 1(Premier cool - 4×10^5), Test sample 2 (Eva - 3×10^5), Test sample 2 (Eva +jik 3×10^3 , Waw+jik 6×10^5 B, Waw+jik 6×10^3 A Good mama 2×10^5 , Canoe 1×10^3 , Joy + laundry bleach(Test sample 1(Jik- 8×10^5) were indole positive test while, Test sample 3(Good mama + laundry bleach(Test sample 1(Jik- 5×10^3), Test sample 4(So easy- 4×10^3 , Test sample 1(Premier cool - 4×10^5 , Test sample 3(Lux- 1×10^3 A) were indole negative test. It was also observed that Test sample 3(Lux - 1×10^5 B), Test sample 1(Premier cool),- 4×10^5), Test sample 3(Good mama - 2×10^5 , Test sample 1(Waw - 3×10^3 A), Test sample 2 (Eva + laundry bleach(Test sample 1(Jik- 3×10^3), Test sample 1(Premier cool + laundry bleach(Test sample 1 (Jik- 4×10^3 A), Test sample 1(Premier cool + laundry bleach(Test sample 1(Jik - 4×10^5 B, Test sample 3(Good mama + laundry bleach(Test sample 1(Jik- 5×10^3 , Test sample 1(Waw + laundry bleach(Test sample 1(Jik - 6×10^3 A), Test

sample 1(Waw + laundry bleach(Test sample 1 (Jik - 6×10^5 B), Test sample 4(So easy + laundry bleach(Test sample 1(Jik - 7×10^3) were positive for oxidase test while Test sample 4(Joy + laundry bleach(Test sample 1(Jik - 8×10^5), Test sample 3(Lux + laundry bleach(Test sample 1(Jik - 1×10^3). Test sample 4(So easy - 4×10^3), Test sample 1(Waw - 3×10^5 B), Test sample 2(Canoe - 1×10^3 , Test sample 2 (Eva - 3×10^3 and Test sample 3(Lux - 1×10^3 A) were negative for oxidase test.

Table 5; Sugar fermentation for identification of isolated bacteria from Fairly Used Clothes, Male/ Female Underwear (Towel, Singlet, and Panties) against Newly Emerging Detergents, Laundry Bleach, and Toilet Soap. All the isolates from the sample were positive for the triple sugar fermentation test except Test sample 3(Lux + laundry bleach(Test sample 1(Jik- 1×10^3) is negative for the triple sugar fermentation test. All isolates were positive for Hydrogen sulphide production test (H_2S). Also, Test sample 4(Joy + laundry bleach (Test sample 1(Jik- 8×10^5), Test sample 1(Waw- 3×10^3 A), Test sample 4(So easy - 4×10^3), Test sample 4(So easy + laundry bleach(Test sample 1(Jik- 7×10^3), Test sample 3(Lux - 1×10^3) were positive for gas production tests while the rest isolate was negative for gas production. All the isolates are positive for glucose test except Test sample 1(Waw + laundry bleach(Test sample 1(Jik- 6×10^5) were negative for the glucose test, all the isolates were also positive for mannitol test. It was also observed that Test sample 3(Lux - 1×10^5 B), Test sample 1(Premier cool),- 4×10^5), Good mama 2×10^5 , Waw 3×10^3 A, Eva + jik 3×10^3 , Premier cool+jik 4×10^3 A, Premier cool+jik 4×10^5 B, Test sample 3(Good mama + laundry bleach(Test sample 1(Jik- 5×10^3 , Test sample 1(Waw + laundry bleach(Test sample 1(Jik - 6×10^3 A), Test sample 1(Waw + laundry bleach(Test sample 1(Jik - 6×10^5 B), Test sample 4(So easy + laundry bleach(Test sample 1(Jik - 7×10^3), Test sample 3(Lux - 1×10^3) were positive for gas production tests while the rest isolate was negative for gas production. All the isolates are positive for the glucose test except Test sample 1(Waw + laundry bleach(Test sample 1(Jik- 6×10^5) were negative for the glucose test, all the isolates were also positive for mannitol test.

Table 6; Probable organism isolated from fairly used clothes, male/female underwear (towel, singlet, and panties) against Newly emerging detergents, laundry bleach, and toilet soap. The isolated organisms include; *Aerobacter*

aerogenes (3), *Bacillus polymyxa* (5), *Staphylococcus aureus* (2), *Veillonella parvula* (5), *Escherichia freundii*(1), *Escherichia coli* (3), *Enterobacter aerogenes* (2), *Aerobacter cloacae* (1).




Plate 1. Antibiotic susceptibility using disc diffusion method

This section of this research work shows the efficacy of newly emerging toilet soaps with the addition of laundry bleach against isolated bacteria from fairly used clothes, male and female underwear.

Fig 1; The result shows the efficacy of using toilet soap with the addition of laundry bleach (Jik) against the isolated bacteria from fairly used clothes, male and female underwear, this shows the effectiveness of toilet soap with the addition of laundry bleach against the isolated bacteria from fairly used clothes, male and female underwear on colony plate count. The results show that toilet soap with the addition of laundry bleach was most effective against the isolated bacteria from fairly used clothes, male and female underwear. From this study Test sample 4(Joy) toilet soap and addition of laundry bleach (jik) show the lowest colony counts of bacteria on a plate, before washing (10^5) 68 numbers of colony counts on a plate, (10^3) 128 numbers of colony counts, After washing with Test sample 4(Joy) toilet soap with the addition of laundry bleach (jik)(10^5) 19 number of colony counts, (10^3) 30 numbers of colony counts, After washing with laundry bleach (jik) (10^5) 5 numbers of colony counts, (10^3) 19 numbers of colony counts, which implies it is the most effective toilet soap among the toilet soaps used for the study. The table is arranged accordingly showing the efficacy of toilet soap against bacteria on fairly used clothes in ascending order.

Fig 2; The results show the efficacy of using detergents and laundry bleach (Jik) against the isolated bacteria from fairly used clothes, male/female underwear, the result shows how effective the detergent is against the isolated bacteria from fairly used clothes, male and female underwear. The result shows which detergent is most effective against the bacteria against the isolated bacteria from fairly used clothes, male and female underwear, From this study Test sample 4 (So Easy) detergent shows the lowest colony counts of bacteria on a plate, before washing (10^5) 68 numbers of the count, (10^3) 128 number of counts. After washing with Test sample 4 (So Easy) detergent (10^5) 20 number of counts, (10^3) 33 number of counts, with laundry bleach (jik) (10^5) 10 number of counts, (10^3) 22 number of counts, which implies Test sample 4 (So Easy) detergent is the most effective detergent on bacteria isolated from the sample among the detergents used for the study.

Fig 2 is arranged accordingly showing the efficacy of detergent against bacteria isolated on fairly used cloth in ascending order.

Fig 3; Shows the antibiotics susceptibility test of the identified Gram-negative bacteria isolated from fairly used clothes, male and female underwear. It was observed that *Escherichia coli* and *Aerobacter aerogenes* has the highest zone of inhibition to Reflacine (25mm) while *Escherichia freundii* and *Aerobacter cloacae* has no zone of inhibition to Reflacine (0mm). *Escherichia coli* has the highest zone of inhibition to Gentamycin (30mm), *Escherichia coli* and *Aerobacter aerogenes* has the highest zone of inhibition to Augmentin (28mm), *Aerobacter aerogenes* has the highest zone of inhibition to Ciprofloxacin (30mm), *Escherichia coli* and *Aerobacter aerogenes* has the highest zone of inhibition to Septrin (30mm), *Escherichia coli* has the highest zone of inhibition to Streptomycin (28mm), *Aerobacter aerogenes* has the highest zone of inhibition to Ceporex (30mm), *Escherichia coli* and *Aerobacter aerogenes* has the highest zone of inhibition to Tarivid (25mm), *Veillonella parvula* and *Enterobacter aerogenes* has the highest zone of inhibition to Nalidixic acid (20mm), *Aerobacter aerogenes* has the highest zone of inhibition to Amplicin (26mm).

Fig 4; Percentage zones of inhibition of antibiotics against *Aerobacter aerogenes* isolated from fairly used clothes, male and female underwear. It was observed that Nalidixic acid (NA) 13%, Tarivid (OFX)13%, Septrin

(SXT)15%,Ciprofloxacin(CPX) 18%, Augmentin (AU) 13%,Gentamycin (CN)14%, and Reflacine (PEF)14%

Fig 5; Percentage zones of inhibition of antibiotics against *Veillonella parvula* isolated from fairly used clothes, male and female underwear. It was observed that Nalidixic acid (NA) 13%, Tarivid (OFX) 13%, Septrin (SXT) 13%, Ciprofloxacin (CPX) 12%, Augmentin (AU) 12%, Gentamycin (CN) 12%,Reflacine (PEF) 12% and Amplicin (PN)13%.

Fig 6; Percentage zones of inhibition of antibiotics against *Enterobacter aerogenes* isolated from fairly used clothes, male and female underwear. It was observed that Nalidixic acid (NA) 12%, Tarivid(OFX)12%,Ceporex(CEP) 12%,Septrin (SXT)12%, Ciprofloxacin (CPX) 9%, Augmentin (AU)13%,Gentamycin (CN)9%, Reflacine(PEF)9% and Amplicin (PN)12%

Fig 7; Percentage zones of inhibition of antibiotics against *Escherichia coli* isolated from fairly used clothes, male and female underwear. It was observed that Tarivid(OFX)13%, Streptomycin (S)14%, Ciprofloxacin (CPX) 14%, Augmentin (AU)14%,Gentamycin (CN)16%, Reflacine (PEF)13% and Amplicin (PN)13%.

Fig 8; Shows the antibiotics susceptibility test of the identified Gram-positive organisms isolated from fairly used clothes, male and female underwear. It was observed that *Bacillus polymyxa* has the highest zone of inhibition to Ampiclox (20mm), *Bacillus polymyxa* has the highest zone of inhibition to Rifampicin (20mm), *Aerobacter aerogenes* has the highest zone of inhibition to Amoxil (23mm), *Bacillus polymyxa* and *Veillonella parvula* has the highest zone of inhibition to Streptomycin (20mm), *Aerobacter aerogenes* has the highest zone of inhibition to Norfloxacin (22mm), *Aerobacte raerogenes* has the highest zone of inhibition to Chloramphenicol (20mm), *Staphylococcus aureus* has the highest zone of inhibition to Ciprofloxacin (22mm), *Bacillus polymyxa* has the highest zone of inhibition of Erythromycin (20mm), *Bacillus polymyxa* has the highest zone of inhibition to Levofloxacin (20mm), *Bacillus polymyxa* and *Aerobacter aerogenes* has the highest inhibition of Gentamycin (20mm).

Fig 9; Percentage zones of inhibition of antibiotics against *Bacillus polymyxa* isolated from fairly used clothes, male and female underwear. It was observed that Gentamycin (CN) 13%, Levofloxacin (LEV) 9%,Ciprofloxacin (CPX) 9%,Chloramphenicol (CH) 10%,Norfloxacin (NB) 14%, Streptomycin (S) 11%, Amoxil. (AMX) 15%, Ampiclox (APX) 10%.

Fig 10; Percentage zones of inhibition of antibiotics against *Staphylococcus aureus* isolated from fairly used clothes, male and female underwear. It was observed that Gentamycin (CN)19%, Levofloxacin (LEV)20%,Erythromycin(E)20%,Ciprofloxacin (CPX) 24%,Streptomycin (S) 20%.

Fig 11; Percentage zones of inhibition of antibiotics against *Veillonella parvula* isolated from fairly used clothes, male and female underwear. It was observed that Gentamycin (CN)16%,

Erythromycin(E)9%,Ciprofloxacin(CPX)16%,Chloramphenicol (CH)14%,Norfloxacin (NB) 8%, Streptomycin(S)17%,Amoxil(AMX) 12%,Ampiclox (APX) 8%.

Fig 14; Shows the growth dynamics of bacteria isolated from fairly used clothes, male and female underwear using Ultraviolet spectrophotometer with the wavelength 480λ. It was observed that at 0hr *Escherichia freundii* has the highest growth rate of 0.9003λ and *Aerobacter cloacae* have the lowest growth rate of 0.070λ. At 84th hour *Aerobacter cloacae* has the highest death rate of 0.296λ and *Escherichia coli* has the lowest death rate of 0.095λ.

Fig 15; Shows the growth dynamics and killing kinetics of bacteria isolated from fairly used clothes, male and female underwear, and the addition of Ciprofloxacin antibiotics at the 48th hour using an Ultraviolet spectrophotometer. In this table, it was observed that at 0hr *Aerobacter aerogenes* have the highest growth rate of 0.142λ, and *Veillonella parvula* has the lowest growth rate of 0.059λ. At the 48th hour *Escherichia coli* has the highest death rate of 0.138λ and *Bacillus polymyxa* has the lowest death rate of 0.079λ.

Table 4. Biochemical Examinations for identification of Isolated Bacteria from Fairly Used Clothes, Male/ Female Underwear (Towel, Singlet and Panties) against Newly Emerging Detergents, Laundry Bleach and Toilet Soap.

Newly Emerging Detergents, Laundry Bleach and Toilet Soap Test sample	Motility	Urease	Indole	Catalase	Oxidase	Coagulate	H₂s	Gas Production
Test sample 1 (Premier cool+Jik) 4x 10 ⁵ B	+	+	-	+	+	-	+	-
Test sample 1 (Premier cool+Jik) 4x10 ³ A	-	+	+	-	+	-	+	-
Test sample 1 (Waw+Jik)6x10 ³ A	-	+	+	+	+	+	+	-
Test sample 1 (Waw+Jik)6x10 ⁵ B	-	+	+	+	+	+	+	-
Test sample 1(Premier cool)4x10 ⁵	+	+	+	+	+	+	+	-
Test sample 1(Waw)3x10 ³ A	-	+	+	+	+	+	+	+
Test sample 1(Waw)3x10 ⁵ B	-	+	+	+	-	+	+	-
Test sample 2 (Eva)3x10 ³	+	+	+	-	-	+	+	-
Test sample 2 (Eva+Jik)3x10 ³	+	+	+	+	+	-	+	-
Test sample 2(Canoe)1x10 ³	+	+	+	+	-	+	+	-
Test sample 3 (Good mama)2x10 ⁵	+	+	+	+	+	+	+	-
Test sample 3 (Good mama+Jik) 5x 10 ³	-	+	-	+	+	-	+	-
Test sample 3 (Lux+Jik)1x10 ³	-	-	+	+	-	+	+	-
Test sample 3 (Lux+Jik)2x10 ⁵	+	-	-	-	-	-	+	-
Test sample 3(Lux)1x10 ³ A	+	+	-	+	-	+	+	+
Test sample 3(Lux)1x10 ⁵ B	+	+	+	+	+	+	+	-
Test sample 4(Joy) 2x10 ³	-	-	-	+	+	+	+	-
Test sample 4(Joy+Jik)8x10 ⁵	-	+	+	+	-	-	+	+
Test sample 4(So Easy)4x10 ³	+	+	-	+	-	+	+	+
Test sample 4(So Easy)7x10 ³	+	+	+	+	+	+	+	+

Table 5. Sugar fermentation for identification of Isolated Bacteria From Fairly Used Clothes, Male/ Female Underwear (Towel, Singlet and Panties) against Newly Emerging Detergents, Laundry Bleach and Toilet Soap

Newly Emerging Detergents, Laundry Bleach and Toilet Soap Test sample	Lactose	Sucrose	Dextrose	Glucose	Mannitol
Test sample 1 (Premier cool+Jik) 4x10 ³ A	+	+	+	+	+
Test sample 1 (Premier cool+Jik) 4x10 ⁵ B	+	+	+	+	+
Test sample 1 (Waw+Jik)6x10 ⁵ B	+	+	+	-	+
Test sample 1 (Waw+Jik x10 ³ A	+	+	+	+	+
Test sample 1(Premier cool)4x10 ⁵	+	+	+	+	+
Test sample 1(Waw)3x10 ³ A	+	+	+	+	+
Test sample 1(Waw)3x10 ⁵ B	+	+	+	+	+
Test sample 2 (Eva)3x10 ³	+	+	+	+	+
Test sample 2 (Eva+Jik)2x10 ⁵	+	+	+	+	+
Test sample 2 (Eva+Jik)3x10 ³	+	+	+	+	+
Test sample 2(Canoe)1x10 ³	+	+	+	+	+
Test sample 3 (Good mama)2x10 ⁵	+	+	+	+	+
Test sample 3 (Good mama+Jik) 5x10 ³	+	+	+	+	+
Test sample 3 (Lux+Jik)1x10 ³	-	-	-	+	+
Test sample 3(Lux)1x10 ³ A	+	+	+	+	+
Test sample 3(Lux)1x10 ⁵ B	+	+	+	+	+
Test sample 4(Joy) 2x10 ³	+	+	+	+	+
Test sample 4(Joy+Jik)8x10 ⁵	+	+	+	+	+
Test sample 4(So Easy)4x10 ³	+	+	+	+	+
Test sample 4(So Easy)7x10 ³	+	+	+	+	+

Table 6. Probable organism Isolated From Fairly Used Clothes, Male/ Female Underwear (Towel, Singlet and Panties) against Newly Emerging Detergents, Laundry Bleach and Toilet Soap

Newly Emerging Detergents, Laundry Bleach and Toilet Soap Test sample	Organisms identified
Test sample 1(Premier cool + laundry bleach(Test sample 1(Jik - 4×10^5 B)	<i>Aerobacter aerogenes</i>
Test sample 1(Premier cool + laundry bleach(Test sample 1(Jik - 4×10^3 A)	<i>Veillonella parvula</i>
Test sample 1(Premier cool - 4×10^5)	<i>Bacillus polymyxa</i>
Test sample 1(Waw - 3×10^3 A)	<i>Staphylococcus aureus</i>
Test sample 1(Waw - 3×10^5 B)	<i>Veillonella parvula</i>
Test sample 1(Waw + laundry bleach(Test sample 1(Jik - 6×10^5 B)	<i>Escherichia coli</i>
Test sample 1(Waw + laundry bleach(Test sample 1(Jik - 6×10^3 A)	<i>Escherichia coli</i>
Test sample 2 (Eva + laundry bleach(Test sample 1(Jik - 3×10^3)	<i>Veillonella parvula</i>
Test sample 2(Canoe - 1×10^3)	<i>Veillonella parvula</i>
Test sample 3(Good mama + laundry bleach(Test sample 1(Jik - 5×10^3)	<i>Aerobacter cloacae</i>
Test sample 3(Good mama - 2×10^5)	<i>Escherichia freundii</i>
Test sample 3(Lux + laundry bleach(Test sample 1(Jik 1×10^3)	<i>Escherichia coli</i>
Test sample 3(Lux - 1×10^3 A)	<i>Bacillus polymyxa</i>
Test sample 3(Lux- 1×10^5 B)	<i>Aerobacter aerogenes</i>
Test sample 4(Joy + laundry bleach(Test sample 1(Jik - 8×10^5)	<i>Aerobacter aerogenes</i>
Test sample 4(So easy - 4×10^3)	<i>Bacillus polymyxa</i>
Test sample 4(So easy - 7×10^3)	<i>Veillonella parvula</i>

Key –NEWLY EMERGING DETERGENTS, LAUNDRY BLEACH AND TOILET SOAP

- 1.Toilet soap used (4): - Test sample 1(Premier cool), Test sample 2 (Eva) Test sample 3(Lux), Test sample 4(Joy),
- 2.Detergents used(4): Test sample 1(Waw), Test sample 2(Canoe), Test sample 3(Good mama), Test sample 4(So easy)
- 3.Laundry Bleach (1) Test sample (1) Jik..




Fig. 1. Efficacy of Newly Emerging Toilet Soaps With Addition of Laundry Bleach against Isolated Bacteria From Fairly Used Clothes, Male/ Female Underwear




Fig. 2. Efficacy of newly emrging Detergent With Addition of Laundry Bleach against Isolated Bacteria From Fairly Used Clothes, Male And Female Underwear




Fig. 3. Antimicrobial Susceptibility Test of Identified Gram Negative Bacterial Isolated From Fairly Used Clothes, Male and Female Underwear (Diameter of Zones Inhibition In Millimeter (Mm)

KEY: S- Streptomycin, OFX- Tarivid, PEP- Reflacine CPX- Ciprofloxacin,
 AU- Augmentin, CN- Gentamycin, CEP- Ceporex, NA- Nalidixic acid, SXT- Septrin, PN- Amplicin




Fig. 4. Percentage zones of inhibition of antibiotics against *Aerobacter aerogene* isolated from fairly used clothes, male and female underwear




Fig. 5. Percentage zones of inhibition of antibiotics against *Veillonella parvula* isolated from fairly used clothes, male and female underwear




Fig. 6. Percentage zones of inhibition of antibiotics against Enterobacter aerogenes isolated from fairly used clothes, male and female underwear




Fig .7. Percentage zones of inhibition of antibiotics against Escherichia coli isolated from fairly used clothes, male and female underwear




Fig. 8. Antimicrobial Susceptibility Test of Identified Gram-Positive Bacterial Diameter of Zones of Inhibition in Millimeter (Mm)

KEY: S- Streptomycin, NB- Norfloxacin, CH- Chloramphenicol, CPX- Ciprofloxacin, E- Erythromycin, LEV- Levofloxacin, CN- Gentamycin, APX- Ampiclox, RD- Rifampicin, AMX- Amoxil




Fig. 9. Percentage zones of inhibition of antibiotics against *Bacillus polymyxa* isolated from fairly used clothes, male and female underwear




Fig. 10. Percentage zones of inhibition of antibiotics against *Staphylococcus aureus* isolated from fairly used clothes, male and female underwear




Fig. 11. Percentage zones of inhibition of antibiotics against *Veillonella parvula* isolated from fairly used clothes, male and female underwear




Fig. 12. Efficacy of Toilet Soap test sample Against Bacteria Isolated From Fairly Used Clothes, Male /Female Underwear (Plate count 10^{-6})




Fig. 13. Efficacy of Detergent test sample Against Bacteria Isolated From Fairly Used Clothes, Male /Female Underwear (plate count 10^{-6})




Fig. 14. Growth Dynamics of Bacteria Isolated From Fairly Used Clothes, Male and Female Underwear Using Ultraviolet Spectrophotometer With The Wavelength 480λ. (With Control: 0.000)




Fig. 15. Growth Dynamics and Killing Kinetics of Bacteria Isolated From Fairly Used Clothes, Male/Female Underwear and Addition of Ciprofloxacin Antibiotics At 48th Hour Using Ultraviolet Spectrophotometer (With Control: 0.000).

4. DISCUSSION

The aim of this study is to determine the potency of newly emerging detergents, laundry bleach, and toilet soap on bacteria isolates from fairly used clothes and Female Underwear. The different samples of fairly used cloth examined in this study showed varying degrees of microbial load with bacteria. The level of patronage given to these fairly used clothes by all in the society makes it a matter of public health importance. Though there are no data to show the link between the wearing of fairly used clothes and any infectious disease or disease outbreaks, the organisms isolated have been incriminated in one disease condition or the other. Towel, Female singlet, female bra, and panties, which is one of the commonly purchased fairly used clothes, has the highest bacteria count from this study. This brings to surface facts, the need to know the source of these fairly used clothes. [12,24]. demonstrated the colonization of Towel and curtains in a Japanese hospital with pathogens like Methicillin-resistant *Staphylococcus aureus* (MRSA), *Clostridium difficile*, and *Vancomycin-resistant Enterococci* (VRE). If these Towels get to the markets as fairly used clothes, the lives of buyers/users become endangered and increased related health risks. The varied efficacy of the different detergents and toilet soap used for different fairly used cloth types recorded in this study provides a guide as to which detergent and toilet soap may be used with a particular fairly used cloth/wear.

From this research work, towel, adult wears, male boxer and female panties and bra, Test sample 3(Good mama) detergent performed better than other detergents, and J Test sample 4(Joy) and Test sample (1) Laundry Bleach (Jik) performed better than other soap the reason for this might be in the components of the detergents and toilet soap, the texture of the cloth materials and the primary level of microbial load. Towels with heavier textures are better re-washed with Laundry Bleach (Jik). However, there was a general reduction in the microbial load with the use of the detergents and toilet soap, this agrees with the studies [25].

Fairly used wears from this research work. *Staphylococcus aureus* was the most commonly isolated bacterium from the examined fairly used cloth samples. This organism has also been isolated from previous studies by [25,26] and [27]. *Staphylococcus aureus* has been

incriminated largely in various infections with multiple drug-resistant strains, skin infections such as boils and furuncles (28). This organism is also associated with pneumonia, bacteremia, and endocarditis [29,30]; *Staphylococcus aureus* isolation from these cloth samples no doubt is of public health concern.

Bacillus sp was the principal bacterium isolated by (1) and was also isolated from this study. *Bacillus* sp is not known as a human pathogen and is another health risk. Its isolation might be related to the fact that *Bacillus* sp generally can withstand harsh environmental conditions like heat, desiccation, and ultraviolet irradiation because of its ability to form endospores by which it can remain dormant for years [31;1], especially if the cloth is been transported from one region to the other, a very good example is the importation of this fairly used cloth from the United Kingdom and the United States of America.. *Bacillus polymyxa*, was implicated during this research work [23]. Other isolated organisms from the fairly used cloth and male and female underwear includes *Escherichia freundii*, *Veillonella parvula*, *Enterobacter aerogenes*, and *Aerobacter aerogene*.

Veillonella parvula is an anaerobic gram-negative coccus that is part of the normal human flora. It has rarely been identified as a pathogen in humans, and the most frequently reported infection caused by *Veillonella parvula* is osteomyelitis. a case of bacteremia unrelated to a central venous catheter and without an underlying source of infection [32]. They are usually recovered as part of polymicrobial infection and they are often regarded as a contaminant. However, they have been isolated in pure culture from various sterile body sites such as sinuses, lungs, liver, central nervous system, heart, and bone. The presence of this organism depends on the previous users of the fairly used cloth. But it is worthy of note, to find *Veillonella parvula* in the fairly used cloth during this research work [33,34,35,28,34].

Enterobacter aerogenes have been reported as important opportunistic and multi-resistant bacterial pathogens for humans during the last three decades in hospital wards. These Gram-negative bacteria have been largely described during several outbreaks of hospital-acquired infections in Europe and particularly in France where there is an influx of this imported fairly used cloth. The dissemination of *Enterobacter* sp. is associated with the presence of redundant

regulatory cascades that efficiently control the membrane permeability ensuring the bacterial protection and the expression of detoxifying enzymes involved in antibiotic degradation /inactivation. Emphasis should be taken to cognizant that *Enterobacter aerogenes* species are able to acquire numerous genetic mobile elements that strongly contribute to antibiotic resistance. Moreover, this particular fitness helps them to colonize several environments and hosts and rapidly and efficiently adapt their metabolism and physiology to external conditions and environmental stresses [36], to corroborate the finding of this research work, the isolated organism may travel from coast to coast i.e. horizontal transmission of mobile elements containing antibiotic resistance genes.

Escherichia freundii is another isolated organism, *Citrobacter freundii*, is gram-negative enterobacteria classified as a genus in the family Enterobacteriaceae. *Citrobacter freundii* species are often found in the stools of children, where they are generally regarded as normal components of the intestinal microflora. *Citrobacter freundii* may cause urinary tract infections, and are found in wounds, respiratory, meningitis, and sepsis. They can cause healthcare-associated infections, especially in pediatric and immuno-compromised patients. *Citrobacter freundii* have been reported previously in association with enteritis in humans, but thus far little effort has been made to clarify the nature of this association. It has been well established that *Escherichia coli* and some non-*E. coli* enterobacteria can cause diarrhea by producing either heat-stable (ST) or heat-labile (LT) enterotoxins [37].

Hence, isolation of these different organisms like *Staphylococcus aureus* from fairly used cloth samples calls for concern from this study which has been documented in serious wound infections, many strains of which are multidrug-resistant [38], *Staphylococcus aureus* was isolated in 17.1% of the cloth samples, this is in agreement with the work of [12]. *Enterobacter aerogenes* is associated with nosocomial infection, eye and skin infection, vagina infections especially among diabetics, pregnant women, and immunocompromised patients [39]. They are equally isolated from this study. Bacteria stay long on fabrics hence they are constantly present on fairly used cloth materials [40]. The counts of microbial load on used fairly used clothes might be due to factors like customers' hand touching to feel texture and

quality, environmental pollution, sewing, and packaging system or seller's carrier status. It is therefore advisable to display new wear in covered or glass-cased showrooms as against the open display commonly encountered to attract buyers common in Nigeria [41,42].

It was observed that Ciprofloxacin(CPX) has 18% percentage zones of inhibition of antibiotics against *Aerobacter aerogene*, Nalidixic acid (NA) and Tarivid(OFX) has 13% percentage zones of inhibition of antibiotics against *Veillonella parvula*, Augmentin (AU) has 13% percentage zones of inhibition of antibiotics against *Enterobacter aerogenes*, Augmentin (AU) has 14% percentage zones of inhibition of antibiotics against *Escherichia coli*, Amoxil (AMX) has 15% percentage zones of inhibition of antibiotics against *Bacillus polymyxa*, Ciprofloxacin (CPX) has 24% percentage zones of inhibition of antibiotics against *Staphylococcus aureus* and Streptomycin(S) has 17% percentage zones of inhibition of antibiotics against *Veillonella parvula*. The practice of formulating the toilet soap and detergent with specialized antibiotics should be upheld, this will give us leverage on the action of toilet soap /detergent on microbes isolated from the fairly used cloth and male /female underwear. The antibiotics will increase the killing action of the microbe thereby decontaminating the fairly used clothes to reduce the disease risk factors. One such example was demonstrated in the killing kinetics during the course of the research work, Ciprofloxacin, these antibiotics accelerate the death rate of the isolates before reading was taken by the Ultraviolet spectrophotometer. If antibiotics are added to the toilet and detergent then risen with the laundry bleach, the microbe will be eradicated permanently [18,23].

The emergence of new toilet soap, detergent laundry bleach will help to eradicate the so-called emerging normal flora organism if correctly formulated and effectively used Exposure of strains of bacteria, to lethal doses of hypochloric acid (Laundry bleach, toilet soap and detergent) causes a decrease in ATP production. Chlorine dioxide present in the detergent and soap acts on the permeability of the external membrane of the bacteria e.g. *E. coli*, through a primary lethal phenomenon that consists in a substantial leakage of K^+ ion; such leakage does not occur for macromolecules. Sub-lethal doses inhibit cellular respiration due to a nonspecific oxidizing effect [43]. All fairly used cloth should be washed with toilet soap and a mixture of detergent and a

high concentration of laundry(Jik) bleach before use, this will drastically reduce the microbial load on the cloth and prevent major health hazards, that may emanate from this fairly used cloth.

The direct effect of sunlight is of paramount importance in this study. The sun produces UV rays, this rays has a lethal effect on microbes at a preset wavelength after exposure for specific number of days. This is reason why, it is necessary to spread the fairly used cloth in sunlight for days before use. The UV rays will help to destroys existing and survived colony of microbe on fairly used cloth and underwear after washing with different soaps and detergent. The UVrays from the sun stimulates the rapid death rate of microbes on our farly used cloths and underware, this is deduces from this research work. The UV spectrophotometer were used, to produce artificial UV rays on the isolated organisms from the fairly used cloth and underwear, to determine the killing kinetics of microbe artificially, and the result was extraordinary at 480λ wavelength for 48 to 60 hours, No forms of microbe can survive as deduced from this study [18,44].

It was deduced that there is need to spread our fairly used clothing and underwear in the sun before and after used to reduce the number of existing colony of microbe, to avoid the scourge of microbial infection. Therefore, we should dissuade the mind of third world country, not all fairly used clothing are wholesome, and adequate precaution must be taking before use. Fairly used clothing and underwear must be washed with soaps, detergent, laundry bleach and finally spread in the sunlight.

5. CONCLUSION

Potential pathogens were isolated from the fairly used clothes with Towel, Singlet, bra, and panties have been implicated as the ones with the highest microbial load. These isolated microbes were incriminated in life-threatening diseases. Though the use of detergents, toilet soap, and laundry bleach in washing and re-washing the fairly used cloths reduces the microbial load drastically if not totally eradicated. fairly used clothes consumption is becoming a big industry in the third world like Nigeria because they are relatively cheaper and a little more durable quality better than the new cloth, it is very important and imperative, to educate the populace, about the danger and health implications of fairly used cloth, if not properly

washed and rewashed before use and different soap and detergent, To remove the intending danger of fairly used cloth.

6. RECOMMENDATION

From the result obtain from the efficacy of newly emerging detergents and toilet soap against bacterial isolated from fairly used clothes and underwear, I recommend that detergent or toilet soap can be used to wash any fairly used clothes purchased before use to reduce the microbial load present on the cloth and to prevent contacting of infectious disease, I also recommend that (Good mama)detergent and toilet soap with a high concentration of laundry bleach is best for washing of fairly used cloth.

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

ACKNOWLEDGEMENTS

The laboratory staff of Adekunle Ajasin University, Department of Microbiology, Faculty of Science, Akungba Akoko, Ondo State, Nigeria.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Agbulu CO, Gberikon GM, Ajine BO. Isolation and characterization of microorganisms associated with second hand female undergarments and children wear sold in Makurdi Metropolis. (Int. J. Curr. Microbiol. App. Sci. 2015;4(1):716-724 [2]).
2. Malnick S, Bardenstein R, Huszar M, Gabbay J, Borkow G. Pyjamas and sheets as a potential source of nosocomial pathogens. (Journal of Hospital Infection. 2008;70:89-92.

3. Scott TA, Batey SFD, Wienczek P, Wienczek G, Alt S, Francklyn CS. Immunity-Guided Identification of Threonyl-tRNA Synthetase as the Molecular Target of Obaflourin, a β -Lactone Antibiotic. *ACS Chem. Biol.* 2019;14 (12):2663–2671
4. Layton A, McKay L, Williams D, Garrett V, Gentry R, Saylor G. Development of Bacteroides 16S rRNA gene TaqMan-based real-time PCR assays for estimation of total, human, and bovine fecal pollution in water. *Appl Environ Microbiol.* 2006;72:4214– 4224.
5. Greene VW. Personal Hygiene and Life expectancy improvements since 1850; Historic and Epidemiological association. (*American Journal of Infection Control.* 2001;29:203-206).
6. Matiru V. Mapping of organizations engaged in climate change adaptation in Kenya: Final draft. Unpublished report prepared for the Worldwide Fund for Nature; 2009.
7. Frumkin H. Beyond toxicity human: health and the natural environment. *American Journal of Preventative Medicine.* 2001;20:234–240.
8. Hussain S, Khan F, Hussain HA, Nie L. Physiological and biochemical mechanisms of seed priming-induced chilling tolerance in rice cultivars. *Front. Plant Sci.* 2016;7:116.
9. Olajubu FA, Folorunsho VT, Olajede O. The Microbial Diversity of Fairly Used Wears Sold in Lagos Market, Nigeria. 2017. Department of Microbiology, Adekunle Ajasin University, Akungba-Akoko, Nigeria. 2017;12(2): 63-68
10. Aiello AE, Larson EL, Sedlak R. Hidden heroes of the health revolution sanitation and personal hygiene. *American Journal of Infection Control.* 2008;36:128-51
11. Pilonetto M, Rosa EA, Brofman PR, Baggio D, Calvário F, Schelp C, Nascimento A and Messias-Reason I. Hospital gowns as a vehicle for bacterial dissemination in an intensive care unit. *Brazilian Journal of Infectious Diseases.* 2004;8:206-10.
12. Shiomori T, Miyamoto H, Makishima K, Yoshida M, Fujiyoshi T, Uda T, Inaba T, Hiraki N. Evaluation of bed making-related airborne and surface methicillin resistant *Staphylococcus aureus* contamination. *Journal of Hospital Infection.* 2002;50:30-5.
13. Aiello AE, Larson EL. Causal inference: the case for hygiene and health. *American Journal of Infection Control.* 2002;30: 503-11.
14. Levy SB. Antibiotic resistance: consequences of inaction. *Clinical Infectious Diseases.* 2001;5124–9
15. Osuntokun OT, Mary TT. Comparative Study of Ethanolic Wild African Nutmeg (*Pycnanthus angolensis* (Welw.) Stem Bark Extract Potentials and Selected Conventional Toothpaste against Hidden Resident Mouth Cavity Microflora. *Asian Journal of Biochemistry, Genetics and Molecular Biology.* 2020;6(2):10-36.
16. Russell AD, Tattawasart U, Maillard JY, Furr JR. Possible link between bacterial resistance and use of antibiotics and biocides. *Antimicrobial Agents and Chemotherapy.* 1998;42:2151.
17. Osuntokun OT, Ibukun AF, Yusuf-Babatunde AM, Abiodun . Pre/post-plasmid profile analysis, killing- kinetics and secondary metabolites screening of *Adenopus breviflorus* (Benth) fruit extract against multiple drug resistant isolates using *Staphylococcus aureus* (MDRSA) as a case study. *Journal of Advanced Research in Biotechnology. J Adv Res Biotech.* 2019;4(1):1-17.
18. Osuntokun OT, Thonda OA, Akele EO, Adedokun LO, Adedayo SA, Bello OA. Pathogenic Bacteria Found on Surfaces of Canned Drinks and Wines Being Sold In Retail Shops in Ondo state, Nigeria, Health Implications, Food Safety and Quality Assessment. *South Asian Journal of Parasitology.* 2021;5(4):68-94.
19. Fawole MO, Oso BA. ISSN 1597-6343 Published by Faculty of Science; 2007.
20. Cheesbrough, M. (*District Laboratory Practice in Tropical Countries Cambridge University Press.* 2006:434.
21. Tankeshwar A. Citrate utilization test: principle procedure, expected results and positive organisms. *Microbe.* 2013:20-2.1
22. Clinical and Laboratory Standards Institute (CLSI). Performance standards for antimicrobial susceptibility testing; Twenty-sixth informational supplement. CLSI document M100-S26. Wayne, PA: Clinical and Laboratory Standards Institute; 2016.
23. Osuntokun OT, Azuh VO, Adejoro BF, Akele EO. Antimicrobial Spectrum, Growth/Killing Kinetics, Conventional/Molecular Assay of Characterizing Non-Leguminous Endophytic Bacteria and Fungi from *Helianthus annuus*, *Carica papaya* and

- Lycopersicon solanum. J Biomed Res Environ Sci. 2021;2(10):1018-1034.
24. Trillis F, Eckstein EC, Budavich R, Pultz MJ, Donskey CJ. Contamination of hospital curtains with health care associated pathogens. *Infection Control and Hospital Epidemiology*. 2008;29: 1074-6).
 25. Muthaini YM, Matiru VN, Bii C. Potential second hand clothes and the effectiveness of disinfection methods. *Kenyan Journal of Microbiology*. 2010;1: 1-19
 26. Sattler C, Armando C. Coagulase Positive Staphylococcal Infections (Staphylococcus aureus), In: Ralph D. Feign (Editor), *Textbook of Paediatric infectious diseases*, 5th Edition, Saunders Publishers. 2004:1099-1123.
 27. Gaspard P, Eschbach E, Gunther D, Gayet S, Bertrand X and Talon D. Methicillin resistant Staphylococcus aureus contamination of healthcare workers' uniforms in long-term care facilities. (*Journal of Hospital Infection*. 2009;71:170–5).
 28. Khadri H, Alzohairy M. Prevalence and antibiotic susceptibility pattern of methicillin-resistant and coagulase-negative staphylococci in a tertiary care hospital in India. *Int. J. Med. Med. Sci*. 2010;2(4):116-120
 29. Tekalign K, Bacha K. Prevalence and antibiotic susceptibility pattern of methicillin resistant Staphylococcus aureus among primary school children and pioneers in Jimma town, (South West Ethiopia. *Annals of Clinical Microbiology and Antimicrobials*. 2013;12:11)
 30. Fowler VG, Helen MHS, Boucher W, Corey RG, Abrutyn E. Daptomycin versus standard therapy for bacteremia and Endocarditis caused by Staphylococcus aureus. (*The new England Journal of Medicine*. 2006;355 (7):1812-17)
 31. Nester EW, Anderson GD, Roberts EC, Pearsall NN, Nester MT. (*A Human Perspective Microbiology*, 4th Edition. 2004:535-645.
 32. Barnhart RA, Weitekamp MR, Aber RC. Osteomyelitis caused by Veillonella. *Am J Med*. 1983;(5):902–904
 33. Chaturvedi VN, Methwani A, Chaturvedi P, Narang P. Bacterial flora in chronic tonsillitis. *Indian Pediatr*. 1989;(1):52–56.
 34. Koksai F, Yasar H, Samasti M. Antibiotic resistance patterns of coagulase-negative staphylococcus strains isolated from blood cultures of septicemic patients in Turkey. (*Microbiological Research*. 2009; 164:404-410.
 35. Huang SY, Tang RN, Chen SY. Coagulase-negative staphylococcal bacteremia in critically ill children: risk factors and antimicrobial susceptibility. (*J Microbiol Immunol Infect*. 2003;36:51-55)
 36. Hammerum AM, Hansen F, Nielsen HL, Jakobsen L, Stegger M, Andersen PS, Jensen P, Nielsen TK, Hansen LH, Hasman H, Fuglsang-Damgaard D. Hammerum AM(2016), *J Antimicrob Chemother*. *J Antimicrob Chemother*. 2016;(11):3117-3124.
 37. Back E, Mollby R, KaUser B, Stintzing G, Wadstrom T, Habte D. Enterotoxigenic E. coli and other gram-negative bacteria of infantile diarrhea: surface antigens, hemagglutinins, colonization factor antigen and loss of enterotoxigenicity. *J. Infect. Dis*. 1980;142:318-327. 3.
 38. Nkang AO, Okonko IO, Mejeha OK, Adewale OG, Udeze AO, Fowotade A, Fajobi EA, Adedeji AO, Babalola ET. (Assessment of antibiotics susceptibility profiles of some selected clinical isolates from laboratories in Nigeria. *J. Microbiol. Antimicrobial*. 2009;1(2): 19-26
 39. Kothavade RJ, Kura MM, Panthaki MH. Candida tropicalis, its prevalence, pathogenicity and increasing resistance to fluconazole. (*Journal of Medical Microbiology*. 2014;5(2):35-39)
 40. Muthaini YM, Matiru VN, Bii C. Potential second-hand clothes and the effectiveness of disinfection methods (*Kenyan Journal of Microbiology*. 2010;1: 1-19.
 41. Ertek M, Yazgi H, Erol S. Demonstration of in vitro antagonism between fusidic acid and quinolones. *J Int. Med. Res*. 2002;30:525-8.
 42. Pankaj K Tyagi, Shruti Tyagi. Bacterial contamination in kitchens of rural and urban areas in Meerut District of Uttar Pradesh (India), *African journal of Microbiology Research*. 2013:20-34
 43. Bettelheim KA, Evangelidis H, Pearce JL, Sowers E, Strockbine NA. Bettelheim KA, *ClinMicrobiol*. 1993;(3):760-1. DOI: 10.1128/jcm.31.3.760-761.1993. *J ClinMicrobiol*. 1993. PMID: 7681442 Free PMC article.
 44. Osuntokun OT. *Antimicrobial Spectrum, Growth/ killing kinetics, Conventional/Molecular assay and Ultraviolet Spectrophotometer Signatures*

of Characterizing Shigella Flexneri and Enterococcus Faecalis and Isolated from Swine House isolates”, International

Journal of Pharmacy and Infections Therapy Int J Phar Inft Thrp. 2021;4(1): 1-27.

© 2022 Osuntokun et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/86011>