



## Effect of Ethanol Leaf Extract of *Gongronema latifolium* (Bush Buck) on the Reproductive System of Male Albino Rats

Grace Sylvester Effiong<sup>a\*</sup>, Othuke Bensandy Odeghe<sup>b</sup>, Nse Udoka Ebe<sup>a</sup>,  
Sunday Edet Enidiok<sup>c</sup>, Uduak Onofiok Luke<sup>a</sup>, Emmanuel Onyi Nwuzor<sup>a</sup>  
and Esther Sunday Enidiok<sup>a</sup>

<sup>a</sup> Department of Biochemistry, Faculty of Science, University of Uyo, P. M. B. 1017, Uyo, Nigeria.

<sup>b</sup> Department of Biochemistry, Faculty of Science, Madonna University,  
P. M. B. 05 Elele, Rivers State, Nigeria.

<sup>c</sup> Department of Food Science and Technology, Faculty of Agriculture,  
University of Uyo, P. M. B. 1017, Uyo, Nigeria.

### Authors' contributions

This work was carried out in collaboration among all authors. Author GSE designed the study. Author NUE performed the statistical analysis. Author SEE wrote the protocol and wrote the first draft of the manuscript. Authors EON and ESE managed the analyses of the study. Authors OBO and UOL managed the literature searches. All authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/AJRB/2022/v10i230222

### Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/84266>

Original Research Article

Received 02 January 2022  
Accepted 04 March 2022  
Published 30 March 2022

### ABSTRACT

Medicinal plants are plants used for medicinal purposes and are commonly used in treating and preventing specific ailments and diseases that are generally considered to be harmful to humans. *Gongronema latifolium* is a well-known plant that is beneficial in preventing and treating certain diseases and ailment due to their phytochemical constituents. This study evaluates the effect of ethanol leaf extract of *Gongronema latifolium* on the reproductive system of male Albino rats using standard methods. Twenty-five male rats equally divided into five groups and five female rats were used. Group I and II served as the Normal and Positive controls and were orally administered with distilled water and subcutaneously standard drug; testosterone respectively. The other three groups were orally treated with *Gongronema latifolium* extract at low, middle and high dosage (200,300 and 400mg/kg body weight) respectively. After 7 days of treatment, the female rats were introduced into the male cages in the ratio 1:1 (male: female) to ascertain for the different

\*Corresponding author: E-mail: graceeffiong2007@yahoo.com, graceeffiong@uniuyo.edu.ng;

aphrodisiac parameters. Treatment continued for another 7 days after which the male animals were sacrificed and blood samples collected for hormonal assay. Results showed that all the aphrodisiac frequencies had a significant increase ( $p \leq 0.001$ ) in mounting frequency, intromission frequency and penile erectile frequency when compared with the controls while the mounting latency, intromission latency, ejaculatory latency and penile erectile latency were significantly reduced ( $p \leq 0.001$ ) in comparison with the controls in this study. Most of the reproductive hormones were significantly increased ( $p \leq 0.001$ ) in the extract treatment group when compared to the controls. Thus, this study suggests that the crude leaf extract of *Gongronema latifolium* may possess aphrodisiac properties and consequently on the reproductive system.

**Keywords:** Medicinal plants; *gongronema latifolium*; phytochemical; reproductive system; aphrodisiac; reproductive hormones.

## 1. INTRODUCTION

### 1.1 Medicinal Plant and Its Importance

Medicinal plants are those plants that are commonly used in treatment and prevention of specific ailments and diseases that are generally considered to be harmful to humans [1]. They have been known to be an important potential source of therapeutics or curative aids and are used with the intention of maintaining health, to be administered for a specific condition, or both, whether in modern medicine or in traditional medicine [2] and these are due to their phytochemical constituents which are chemically vital and active substances found in the plants that produce specific physiological action on the human body. The most important bioactive components of plants are; flavonoids, tannin, phenolic compounds and alkaloids [3]. Some medicinal plants, known as synergic medicinal plants, can complement, damage, or neutralise their potential negative effects in the body; others, known as official medicinal plants, are used in the treatment of complex cases such as cancer diseases; and still others, known as preventive herbal medicinal plants, have the ability to prevent the occurrence of certain diseases by reducing the side effects of synthetic treatment [4]. Medicinal plants have come to play a significant part in health care systems all around the world. This includes using medicinal plants not only to treat ailments, but also as a potential material for sustaining excellent health and circumstances. *Gongronema latifolium* is the botanical name for a local herb called 'utazi' by the Eastern part of Nigeria (Igbos) [5] 'arokeke' by the Western part of Nigeria (Yorubas) and 'utasi' by the Efik and Ibibios [6,7]. *Gongronema latifolium* is of West African Origin [8]. It can be found in Nigeria as well as other tropical nations like Guinea-Bissau, Western Cameroon, and Sierra Leone [9]. It's an edible rainforest plant native to Nigeria's south-eastern region. It's included as one of Nigeria's medicinally

important aromatic plants [10]. *Gongronema latifolium* is found across the tropical and subtropical parts of the world, particularly in West African countries (such as Nigeria, Cote d'Ivoire, Sierra Leone, Ghana, and Senegal) and America, with an average abundance in Northern and South Eastern Asia [11]. The common name for this leaf is Bush buck leaf and it belongs to the family of Asclepiadaceae [12,13]. *Gongronema latifolium* is a well-known medicinal plant in herbal medicine used in antimicrobial, anti-inflammatory and antioxidant activity as a result of its phytochemical properties. It belongs to the class of medicinal plants that are beneficial in preventing and treating certain diseases and ailments that are detrimental to human health. It is one of those leaves with numerous benefits for the human organism and this include both nutritional and health benefits such as; fertility benefits as it helps raise low sperm count, act as pain reliever, helps to lower sugar level, etc [14]. Pharmacological studies speculate that the plant has analgesic, antimicrobial, antibacterial, antiulcer, antioxidant, anti-asthmatic, antipyretic, hypoglycemic and anti-inflammatory properties [14]. Studies by [15] showed the presence of different types of alkaloids, flavonoids, total phenolic compound, lignan, terpenes, sterol, allicin, hydroxycinnamic acids, saponin and carotenoid in the leaves of *Gongronema latifolium* with some in an appreciable concentration. These results showed that *Gongronema latifolium* leaves possess a wide variety of phytochemicals which can be of pharmacological implication, both in terms of prevention and treatment/management of diseases. Important phytochemicals obtain from *Gongronema latifolium* plant and their health benefits are as indicated: Flavonoids are known to inhibit formation of plaques in arteries and so prevent arteriosclerosis, hypertension and other cardiovascular diseases [16,17]. They are also very important antioxidants that mop up reactive oxygen radicals known to be involved in many

conditions that cause cancers, diabetes, inflammatory diseases and neurodegenerative diseases [18]. Saponins lower cholesterol and glucose level. They are also involved in ulcer protection and certain antimicrobial activity [19]. Alkaloids are involved in antimicrobial and hypoglycemic activities [20]. Resins and essential oils have also been involved in antimicrobial [21], anti-inflammatory and antioxidant properties [22,23]. Cardiac glycosides (mainly cardenolides and bufadienolide groups) are useful for treatment of heart conditions [24].

The entire male reproductive system is dependent on hormones which are involved in fertility and sexuality and are usually produced in the testes [25]. The primary hormones involved in the functioning of the male reproductive system are follicle stimulating hormone (FSH), luteinizing hormone (LH) and testosterone. Sexuality is an important and fundamental factor in reproduction including conjugation, conception and procreation. Sexual dysfunction which is the inability to attain normal sexual activity, loss or partial erection, inability to keep erection, premature ejaculation, reduced libido or orgasm, arousal disorder and lack of detumescence [26] has been estimated to occur in approximately 30 million men globally [27]. Substances that have the ability to stimulant or increase sexual desire, performance, arousing sexual instinct, increased pleasure and enjoyment are referred to as aphrodisiacs [28] and they exert their sexual enhancing effects by increasing the flow of blood, promoting erection and by causing relaxation of corpus cavernosal smooth muscle [29]. In need to resolved issues of sexual dysfunction, some medicinal plants have gain attention as they are studied and discovered to have aphrodisiac properties and are of fewer or no side effects.

Testosterone is a medication (androgen family of medications) and naturally occurring steroid hormone [30]. Testosterone can be in form of a gel or patch that is applied to the skin, liquid that is injected into a muscle, tablet that is placed in the cheek or tablet that is taken by mouth. It is used to treat male hypogonadism. It may also be used to increase athletic ability in the form of doping [30].

*Gongronema latifolium* contains different active chemicals where some have been validated; thereby possessing good medicinal qualities. Sexual activities including reproductive hormones play a significant role in reproduction, therefore this present work determines the Effect of Ethanol Leaf Extract of *Gongronema latifolium*

(Bush Buck) on the Reproductive System of Male Albino Rats.

## 2. MATERIALS AND METHODS

### 2.1 Collection and Preparation of Plant Materials

*Gongronema latifolium* was purchased from Itam market in Akwa Ibom State, Nigeria in October 2019. The plant was identified and authenticated by Prof. (Mrs.) Uduak Eshiet of the Department of Botany and Ecological Studies, Faculty of Science, University of Uyo, Nigeria. It was given the Voucher Number UUPH 9(a) and was deposited at the Department of Pharmacognosy and Natural Medicine, Faculty of Pharmacy Herbarium, University of Uyo, Nigeria.

The wet method of extraction was used for the extraction. The leaves were plucked from the plants stalk, thereafter the leaves were washed and drained to remove the debris, and 500g weight of the leaves were cut into pieces and immersed in 2.5L of 50% ethanol, stirred and kept in a plastic container for 72 hours. At the end of the three days the mixture was filtered using a cheese cloth and then with a cotton which was immersed in a funnel. The filtrate was then put in two beakers and kept in a water bath at 30-40°C for two days, it yielded 25g of extract in a paste form and was stored in a refrigerator at 4°C until required for analysis.

### 2.2 Experimental Animals

Twenty-five male and five female Albino rats weighing between 150- 200g were used in this study. They were obtained from the animal house of the Faculty of Pharmacy, University of Uyo where the experiment was also carried out. The animals were kept in wooden cages with wire mesh top and maintained under standard conditions of humidity [50±5%] and temperature [28±2°C] and maintained in a 12 hours light and 12 hours dark cycle. They were given growers feed and water *ad libitum*.

### 2.3 Experimental Design

The animals were divided into five (5) groups of five (5) rats per group and labelled (I- V). The animals in group I served as the Normal Control (NC) and was administered distilled water only. The animals in group II which served as the Positive Control (PC) were administered testosterone subcutaneously at 400mg/kg. The animals in group III, IV and V were administered ethanol leaf extract of *Gongronema latifolium* at 200, 300 and 400mg/kg respectively. The

**Table 1. Experimental design**

Groups	No. of Animals	Treatment	Dosage
I (Normal control)	5	Distilled water	5 ml/kg
II (Positive control)	5	Testosterone	400 mg/kg
III (Low dose)	5	Ethanol extract of <i>Gongronema latifolium</i>	200 mg/Kg
IV (Medium dose)	5	Ethanol extract of <i>Gongronema latifolium</i>	300 mg/Kg
V (High dose)	5	Ethanol extract of <i>Gongronema latifolium</i>	400 mg/Kg

required dose of the extract was measured and administered orally to each rat based on their body weight as shown on {Table 1}. The extract was administered for 7 days before observation of aphrodisiac potency of the extract with five female rats in the ratio 1:1 (male: female) and administered for another 7 days after which the animals were sacrificed and blood samples collected for hormonal assay.

#### 2.4 Evaluation of the Aphrodisiac Parameters of the Extract on the Male Albino Rats

After 7 days of administration of the extract, each male rat was placed in a Plexiglass cage and given 10 minutes to acclimatize before the introduction of an estrous female rat. The test commenced when the female rats were introduced into the Plexiglass cage in the ratio 1:1. Observations were conducted in the dark phase of the light-dark cycle under dim light and very quiet conditions. Each test session was considered ended when either of the following took place:

1. At the end of 15 minutes, or
2. Immediately after the post ejaculation intromission, or
3. If ejaculation did not occur within 15 minutes, or
4. If ejaculation latency exceeded 15 minutes, or
5. If the mount latency or post ejaculatory interval exceeded 15 minutes.

Estrous was induced in the females by administration of 17- $\beta$ -estradiol (0.2 ml/kg) and progesterone (0.2 ml/kg), 48 hours and 8 hours respectively before the introduction of the female into the cage containing the male rats.

The following parameters were documented:

**Mount latency (ML):** This is the time interval from the introduction of the female into the cage until the first mount.

**Mount Frequency (MF):** This is the total number of mounts preceding ejaculation.

**Intromission latency (IL):** This is the time interval from the introduction of the female into the cage until the first intromission.

**Intromission frequency (IF):** This is the total number of intromissions preceding an ejaculation.

**Ejaculation latency (EL):** This is the time from the first intromission to ejaculation.

**Post-Ejaculatory Interval (PEI):** This is the time interval between an ejaculation and the next first mount [31,32].

#### 2.5 Sample Collection and Preparation

After 14 days of administration, the animals were anaesthetized with chloroform vapour and sacrificed. Whole blood was collected by cardiac puncture from each animal using sterile needle and 5ml syringe, emptied into a plain sample bottle and allowed to stand for 15minutes to clot before it was spun in a westerfuge centrifuge (model 1384) at 10,000g for 15minutes. Serum was separated with Pasteur pipette into another plain sample bottle and kept in the refrigerator at 4°C until needed for hormonal assay.

### 3. RESULTS

#### 3.1 Phytochemical Screening

The phytochemical screening of ethanol extract of *Gongronema latifolium* as shown in the {Table 2}, revealed the presence of the following secondary metabolites: alkaloids, flavonoids, saponins, tannins cardiac glycoside.

**Table 2. Phytochemical composition of *Gongronema latifolium* leaf extract**

Test	Observation	Inference
Alkaloids	Cream precipitate	+
Salkowski's test	Brown ring	+
Flavonoids	Yellow colouration	+
Saponins	Persistent foaming	+
Tannins	Dark green color	+
Free Anthraquinones	Pink colouration	-
Combined Anthraquinones	Pink colouration	-

Key: - absent, + positive

### 3.2 Effect of Extract on Sexual Behaviour Behaviours of the Male Albino Rats

The result as shown in {Table 3} revealed that Mounting latency (ML) showed a significant increase ( $p \leq 0.05$ ) in the group administered with standard drug ( $1.28 \pm 0.56$ ) when compare to the normal control ( $0.55 \pm 0.33$ ). There was a non-significant decrease in ML in low dose treatment group ( $0.00 \pm 0.00$ ), middle dose treatment group ( $0.15 \pm 0.06$ ) and high dose treatment group ( $0.47 \pm 0.29$ ) when compared to the normal control group. There was a significant decrease ( $P = .05$ ) in ML in the low dose treatment group ( $0.00 \pm 0.00$ ), middle dose treatment group ( $0.15 \pm 0.06$ ), and high dose treatment group ( $0.47 \pm 0.29$ ) when compared to the positive control group ( $1.28 \pm 0.56$ ).

Mounting frequency (MF) had a significant increase ( $P \leq .001$ ) in the middle dose treatment group ( $6.40 \pm 0.55$ ) when compared to the normal control ( $4.00 \pm 0.19$ ). There was also a significant increase ( $P \leq .001$ ) in the group administered with standard drug ( $9.40 \pm 0.80$ ) when compared to the normal control ( $4.00 \pm 0.19$ ). There was a non-significant decrease in the low dose treatment group ( $0.80 \pm 0.80$ ) and high dose treatment group ( $3.00 \pm 0.32$ ) when compared to the normal control. There was a significant decrease ( $P \leq .001$ ) in MF in the low dose treatment group ( $0.80 \pm 0.80$ ) and high dose treatment group ( $3.00 \pm 0.32$ ) when compared to the positive control group ( $9.40 \pm 0.80$ ). There was also a significant decrease ( $P \leq .001$ ) in the middle dose treatment group ( $6.40 \pm 0.55$ ) when compared to the positive control group.

The result also showed that Intromission latency (IL) had a significant increase ( $P = .05$ ) in the group administered with standard drug ( $1.24 \pm 0.52$ ) when compare to the normal control ( $0.73 \pm 0.47$ ). Also, there is a non-significant decrease in IL in low dose treatment group ( $0.00 \pm 0.00$ ), middle dose treatment group

( $0.34 \pm 0.20$ ) and high dose treatment group ( $0.00 \pm 0.00$ ) when compared to the normal control group. There was a non-significant decrease in IL in the low dose treatment group ( $0.00 \pm 0.00$ ), middle dose treatment group ( $0.34 \pm 0.20$ ), and high dose treatment group ( $0.00 \pm 0.00$ ) when compared to the positive control group ( $1.24 \pm 0.52$ ).

Also, from the result; Intromission frequency (IF) had a significant increase ( $P = .05$ ) in the group administered with standard drug ( $8.80 \pm 0.77$ ) when compare to the normal control ( $2.20 \pm 0.42$ ). There was also a significant increase ( $P = .05$ ) in the middle dose treatment group when compare to the normal control ( $2.20 \pm 0.42$ ). There was a non-significant decrease in the low dose treatment group ( $0.20 \pm 0.20$ ) and high dose treatment group ( $0.00 \pm 0.00$ ) when compare to the normal control. There was a significant decrease ( $P \leq .001$ ) in IF in low dose treatment group ( $0.20 \pm 0.20$ ) and high dose treatment group ( $0.00 \pm 0.00$ ) when compared to the positive control group ( $8.80 \pm 0.77$ ). Also, there was a significant decrease ( $p \leq 0.01$ ) in the middle dose treatment group ( $5.40 \pm 0.71$ ) when compared to the positive control group.

Ejaculatory latency (EL) had a significant increase ( $p \leq 0.01$ ) in the middle dose treatment group ( $5.12 \pm 0.20$ ) when compare to normal control ( $0.82 \pm 0.82$ ). There was also a significant increase ( $P = .05$ ) in the group administered with standard drug ( $5.56 \pm 0.80$ ) when compared to the normal control ( $0.82 \pm 0.82$ ). There was a non-significant decrease in the low dose treatment group ( $0.00 \pm 0.00$ ) and high dose treatment group ( $0.00 \pm 0.00$ ) when compared to the normal control. There was also a significant decrease ( $P \leq .001$ ) in EL in the low dose treatment group ( $0.00 \pm 0.00$ ) and high dose treatment group ( $0.00 \pm 0.00$ ) when compared to the positive control group ( $5.56 \pm 0.80$ ). Also, there was a non-significant decrease in the middle dose treatment group ( $5.12 \pm 0.20$ ) when compared to the positive control group.

Penile erectile latency (PEL) showed a significant increase ( $P \leq .001$ ) in the middle dose treatment group ( $6.27 \pm 0.85$ ) when compared to the normal control ( $1.40 \pm 0.40$ ). Also, there was a significant increase ( $P = .05$ ) in PEL in the group administered with standard drug ( $3.00 \pm 0.50$ ) when compared to the normal control group. There was a non-significant decrease in the low dose treatment group ( $0.00 \pm 0.00$ ) and high dose treatment group ( $0.00 \pm 0.00$ ) when compared to the normal control group.

Penile erectile frequency (PEF) had a non-significant increase in the middle dose treatment group ( $0.80 \pm 0.80$ ) when compared to the normal control group ( $0.40 \pm 0.40$ ). There was also a non-significant increase in PEF in the group treated with the standard drug ( $0.60 \pm 0.40$ ) when compared to the normal control group. There was a non-significant decrease in the low dose treatment group ( $0.00 \pm 0.00$ ) and high dose treatment group ( $0.00 \pm 0.00$ ) when compared to the normal control group. There was a non-significant decrease in PEF in the low dose treatment group ( $0.00 \pm 0.00$ ) and high dose treatment group ( $0.00 \pm 0.00$ ) when compared to the positive control group ( $0.60 \pm 0.40$ ) and also a non-significant increase in the middle dose treatment group ( $0.80 \pm 0.80$ ) when compared to the positive control group.

Post ejaculatory interval (PEI) had a significant increase ( $P \leq .001$ ) in the middle dose treatment group ( $9.87 \pm 0.94$ ) when compared to the normal control ( $1.71 \pm 0.71$ ). Also, there was a significant increase ( $P \leq .001$ ) in PEI in the group administered with standard drug, positive control group ( $10.15 \pm 0.50$ ) when compared to the normal control group. There was a non-significant decrease in the low dose treatment group ( $0.00 \pm 0.00$ ) and high dose treatment group ( $0.00 \pm 0.00$ ) when compared to the normal control group. There was a significant decrease ( $P \leq .001$ ) in PEI in the low dose treatment group ( $0.00 \pm 0.00$ ) and high dose treatment group ( $0.00 \pm 0.00$ ) when compared to the positive control group ( $10.15 \pm 0.50$ ). Also, there was a significant decrease ( $P \leq .001$ ) in the middle dose treatment group ( $9.87 \pm 0.94$ ) when compared to the positive control group.

### 3.3 Effect of the Extract on Reproductive Hormones of the Male Albino Rats

From the result as shown in {Table 4 and Fig. 1}, Testosterone showed a significant increase ( $P \leq .001$ ) in the low dose treatment group ( $1.73 \pm 0.20$ ), middle dose treatment group

( $1.68 \pm 0.17$ ) and also in the high dose treatment group ( $1.69 \pm 0.28$ ) when compared to the normal control group ( $0.20 \pm 0.05$ ). There was also a significant increase ( $P \leq .001$ ) of Testosterone in the low dose treatment group ( $1.73 \pm 0.20$ ), middle dose treatment group ( $1.68 \pm 0.17$ ) and high dose treatment group ( $1.69 \pm 0.28$ ) when compared to the positive control which is the standard drug ( $0.18 \pm 0.02$ ). There was a non-significant decrease in the group administered with the standard drug ( $0.18 \pm 0.02$ ) when compared to the normal control group.

Also from the result, LH - Luteinizing hormone showed a significant increase ( $P \leq .001$ ) in the low dose treatment group ( $2.58 \pm 0.18$ ) and also in the middle dose treatment group ( $2.57 \pm 0.11$ ) when compared to the normal control group ( $1.53 \pm 0.08$ ). Also, LH showed a significant increase ( $P = .05$ ) in the high dose treatment group ( $2.44 \pm 0.04$ ) when compared to the normal control group. There was a significant increase ( $p \leq 0.01$ ) of LH in the low dose treatment group ( $2.58 \pm 0.18$ ) and middle dose treatment group ( $2.57 \pm 0.11$ ) when compared to the positive control ( $1.51 \pm 0.16$ ). There was also a significant increase ( $P = .05$ ) of LH in high dose treatment group ( $2.44 \pm 0.04$ ) when compared to the positive control. There was a non-significant decrease in the group administered with the standard drug ( $1.51 \pm 0.16$ ) when compared to the normal control group {Table 4 and Fig. 2}.

On {Table 4 and Fig. 3}, FSH – Follicle-stimulating hormone showed a non-significant increase in the low dose treatment group ( $2.37 \pm 0.25$ ), middle dose treatment group ( $3.03 \pm 0.07$ ) and in the high dose treatment group ( $3.15 \pm 0.10$ ) when compared to the normal control group ( $2.19 \pm 0.04$ ). There was a non-significant decrease of FSH in the low dose treatment group ( $2.37 \pm 0.25$ ) when compared to the positive control ( $2.39 \pm 0.27$ ). Also, there was a non-significant increase in the middle dose treatment group ( $3.03 \pm 0.07$ ) and high dose treatment group ( $3.15 \pm 0.10$ ) when compared to the positive control. There was a non-significant increase in the group administered with the standard drug ( $2.39 \pm 0.27$ ) when compared to the normal control group.

## 4. DISCUSSION

The photochemical screening of the *Gongronema latifolium* extract showed the presence of alkaloids, flavonoids, saponins, tannins and cardiac glycoside and the absence of free anthraquinones and combined

anthraquinones which was also in line with the work of [33]. It has been reported that saponin constituents found in many plants may possess fertility potentiating properties and may be useful in treatment of impotency impotencies [34].

Therefore, saponin may have boost testosterone level as well as trigger sexual enhancing effect in this study. The presence of flavonoids in plants has been reported to have a role in alteration of hormonal level [35] and may be responsible for

**Table 3. Effect of *G. latifolium* leaf extract on sexual behavior of the male albino rats**


Treatment (mg/kg)	Normal control	200 (Low dose)	300 (Middle dose)	400 (High dose)	Standard drug (Testosterone)
Mounting Latency (ML)	0.55±0.33	0.00±0.00 <sup>d</sup>	0.15±0.06 <sup>d</sup>	0.47±0.29 <sup>d</sup>	1.28±0.56 <sup>a</sup>
Mounting Frequency (MF)	4.00±0.19	0.80±0.80 <sup>f</sup>	6.40±0.55 <sup>b,e</sup>	3.00±0.32 <sup>f</sup>	9.40±0.80 <sup>c</sup>
Intromission Latency (IL)	0.73±0.47	0.00±0.00	0.34±0.20	0.00±0.00	1.24±0.52 <sup>a</sup>
Intromission Frequency (IF)	2.20±0.42	0.20±0.20 <sup>f</sup>	5.40±0.71 <sup>c,e</sup>	0.00±0.00 <sup>f</sup>	8.80±0.77 <sup>c</sup>
Ejaculation Latency (EL)	0.82±0.82	0.00±0.00 <sup>f</sup>	5.12±0.20 <sup>b</sup>	0.00±0.00 <sup>f</sup>	5.56±0.80 <sup>c</sup>
Penile Erectile Latency (PEL)	1.40±0.40	0.00±0.00 <sup>d</sup>	6.27±0.85 <sup>c,e</sup>	0.00±0.00 <sup>d</sup>	3.00±0.50 <sup>a</sup>
Penile Erectile Frequency (PEF)	0.40±0.40	0.00±0.00	0.80±0.80	0.00±0.00	0.60±0.40
Post Ejaculatory Interval (PEI)	1.71±0.71	0.00±0.00 <sup>f</sup>	9.87±0.94 <sup>c</sup>	0.00±0.00 <sup>f</sup>	10.15±0.50 <sup>c</sup>

Data are expressed as mean ± SEM. Significant at <sup>a</sup>P = .05; <sup>b</sup>P ≤ .01; <sup>c</sup>P < .001 when compared to normal control. Significant at <sup>d</sup>P = .05; <sup>e</sup>P ≤ .01; <sup>f</sup>P ≤ .001 when compared to standard drug. n=5


**Table 4. Effect of *G. latifolium* leaf extract on male reproductive hormones**

Treatment (mg/kg)	Testosterone (ng/mL)	LH (mIU/mL)	FSH (mIU/mL)
Normal Control	0.20±0.05	1.53±0.08	2.19±0.04
200(Low dose)	1.73±0.20 <sup>c,f</sup>	2.58±0.18 <sup>b,e</sup>	2.37±0.25
300(Middle dose)	1.68±0.17 <sup>c,f</sup>	2.57±0.11 <sup>b,e</sup>	3.03±0.07
400(High dose)	1.69±0.28 <sup>c,f</sup>	2.44±0.04 <sup>a,d</sup>	3.15±0.10
Positive Control (Standard drug)	0.18±0.02	1.51±0.16	2.39±0.27


Data are expressed as mean ± SEM. Significant at <sup>a</sup>P = .05; <sup>b</sup>P ≤ .01; <sup>c</sup>P ≤ .001 when compared to normal control. Significant at <sup>d</sup>P = .05; <sup>e</sup>P ≤ .01; <sup>f</sup>P ≤ .001 when compared to the standard drug (positive control). n = 5



**Fig. 1. Effect of ethanol extract of *Gongronema latifolium* on testosterone level**



**Fig. 2. Effect of ethanol extract of *Gongronema latifolium* on lutenizing hormone level**



**Fig. 3. Effect of ethanol extract of *Gongronema latifolium* on Follicle stimulating hormone level**

enhanced male sexual behavior in this study. Reports have also shown alkaloids to may have been responsible for facilitation and enhancement of sexual behavior [36]. Therefore, alkaloid may as well trigger sexual activity in this study. These phytochemical substances are recognised to play essential roles in medicinal plants' bioactivity, and they are the source of their therapeutic benefits [14].

Aphrodisiacs are substances that have the ability to stimulate or increase sexual desire, performance, arousing sexual instinct, increased pleasure and enjoyment [28]. These are substances that can be used to modify impaired sexual functions. In the male rats, the extract caused a marked change in sexual behaviour behaviours. The result of these study showed that the administration of the ethanol leaf extract of *Gongronema latifolium* at the middle dose level of 300mg/kg body weight showed a significant decrease in mounting latency with a

corresponding increase in mounting frequency as compared to the controls suggesting sexual motivation which is not in line with the work of [37] but in line with result of [38]. There was also decrease in intromission latency and increase in intromission frequency suggesting increased copulation rate [39]. The result also showed that the effect of the extract on sexual behaviour behaviours of the rats as showed in Table 3 were dosage specific as its aphrodisiac effect was most effectively seen in the middle dose treatment group. Similar findings were recorded by [39] while working on the potential aphrodisiac activity of *Psoralea corylifolia* in male Albino rats.

## 5. CONCLUSIONS

These indices of libido when taken together pointed to the fact that the *Gongronema latifolium* extract may possess aphrodisiac properties. These effects were further enhanced by the action of the extract in increasing the



serum level of testosterone, follicle stimulating hormone and lutenizing hormone which are hormones involved in fertility and sexuality and are usually produced in the testes. All doses of the extract (Low, Middle and High dose) were seen to boost these hormones level which therefore shows there was a significant increase in these hormones. Hence, *Gongronema latifolium* extract might possess aphrodisiac potentials and may increase hormonal levels.

## ETHICAL APPROVAL

Ethical approval was gotten from the College of Health Sciences, University of Uyo, Akwa Ibom State, Nigeria.

## ACKNOWLEDGEMENT

The Authors wish to acknowledge the efforts and contributions of Mr Bala, Mrs Sifon and Mr. Nsikan in the technical aspect of the research.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

- Jide-Faleye F, Olatunya AM. Assessment of the antioxidant and antibacterial activities of *Petiveria alliacea* and *Viscum album* (MISTLETOE). International Journal of Advanced Research. 2016;4(5):1915-1919.  
DOI: 10.21474/IJAR01/539  
Available:http://dx.doi.org/10.21474/IJAR01/539
- 2.Smith-Hall C, Larsen HO, Pouliot M. People, plants and health: A conceptual framework for assessing changes in medicinal plant consumption. Journal of Ethnobiology and Ethnomedicine. 2012;8(1):8-43.  
DOI: 10.1186/1746-4269-8-43
- Sánchez-Gavilán I, Ramírez, E, de la Fuente, V. Bioactive Compounds in *Salicornia patula* Duval-Jouve: A Mediterranean Edible Euhalophyte. Foods. 2021;10:410.  
Available:https://doi.org/10.3390/foods10020410
- Ahn K. The worldwide trend of using botanical drugs and strategies for developing global drugs. BMB Reports. 2017;50(3):111–116.  
DOI: 10.5483/BMBRep.2017.50.3.221
- Katchy AN, Iyare, CO, Adeniyi, BD, Okonkwo, VO, Ikani, OR. Consumption of Aqueous Leaf Extract During Lactation May Improve Metabolic Homeostasis in Young Adult Offspring. Pakistan Journal of Biological Sciences. 2020;23(9):1201-1209.  
DOI: 10.3923/pjbs.2020.1201.1209
- Ugochukwu NH, Babady, NE. Antioxidant Effects of *Gongronema latifolium* in Hepatocytes of Rat Models of Non-Insulin Dependent Diabetes Mellitus. Fitoterapia. 2002;73:612-628.  
Available:http://dx.doi.org/10.1016/S0367-326X(02)00218-6
- Pulok K, Mukherjee AC, Thomas EB, Bhaskar DC, Amit KA, Suparna GC. Role of medicinal plants in inhibiting SARS-CoV-2 and in the management of post-COVID-19 complications Phytomedicine. 2022;98:153930.  
Availablehttps://doi.org/10.1016/j.phymed.2022.153930
- Aba PE. Evaluation of hepatic glycogen content, some haematological and biochemical parameters of alloxan-induced diabetic rats treated with combinations of glibenclamide and *G. latifolium* extract Journal of Complementary and Integrative Medicine. 2017;14(4):2016-2078.  
Available:https://doi.org/10.1515/jcim-2016-0078
- Beshel JA, Beshel PJ, Nku FN, Oshie C, Owu, DU, Nwokocha, Bórquez J. Blood pressure-reducing activity of *Gongronema latifolium* Benth. (*Apocynaceae*) and the identification of its main phytochemicals by UHPLC Q-Orbitrap mass spectrometry Journal of Basic and Clinical Physiology and Pharmacology. 2020;31(1):2018-0178.  
Availablehttps://doi.org/10.1515/jbcpp-2018-0178
- Utibeabasi UE, Okahale S, Ehiabhi, IAO, Olusegun E. Analyses of the Volatile Constituents and Antimicrobial Activities of *Gongronema latifolium* (Benth.) and *Gnetum africanum* L. Journal of essential oil-bearing plants JEOP. 2013;8(3):324-329.  
DOI: 10.1080/0972060X.2005.10643460.
- Agbo C, Baiyeri P, Obi IU. Indigenous knowledge and utilization of *Gongronema latifolium* Benth: A case study of women in University of Nigeria, Nsukka Bio-Research. 2005;3(2).

- DOI: 10.4314/br.v3i2.28596
12. Machebe NS, Agbo CU and Onuaguluchi CC. Oral administration of *Gongronema latifolium* leaf meal: Implications on carcass and haematological profile of broiler finishers raised in the humid tropics. African Journal of Biotechnology. 2011; 10(30):5800-5805.  
Available:<http://www.academicjournals.org/AJB> DOI: 10.5897/AJB11.074
  13. Morebise O. A Review on *Gongronema latifolium*, an Extremely Useful Plant with Great Prospects. European Journal of Medicinal Plants. 2015;10(1):1-9.  
DOI: 10.9734/EJMP/2015/19713
  14. Balogun ME, Obimma E, Besong E, Djobissie S. *Gongronema latifolium*: A Phytochemical, Nutritional and Pharmacological Review. Journal of Physiology and Pharmacology Advances. 2016;6(1):1.  
DOI: 10.5455/jppa.1969123104000
  15. Imoh C, Uchegbu FO. Phytochemical Analysis of *Gongronema latifolium* (Benth) Leaf Using Gas Chromatographic Flame Ionization Detector. Int'l J. Chem. and Biomol. Sc. 2015;1(2):60–68.  
Available:<http://www.aiscience.org/journal/ijcbs>
  16. Panche AN, Diwan AD, Chandra SR. Flavonoids: An overview. J Nutr Sci. 2016;5:47.  
Published online 2016 Dec 29.  
DOI: 10.1017/jns.2016.41  
PMCID: PMC5465813 PMID: 28620474.
  17. Wang X, Ding G, Liu B, Wang Q. Flavonoids and antioxidant activity of rare and endangered fern: *Isoetes sinensis*. PLoS ONE. 2020;15(5): e0232185.  
Available:<https://doi.org/10.1371/journal.pone.0232185>
  18. Lukong CB, Ifemeje JC, Chukwuka DT, Onah C. Effects of aqueous extracts of *Gongronema latifolium* leaves and *Piper guineense* seeds on liver and kidney biomarkers of albino rats. Trop. J. Appl. Nat. Sci. 2018;2(2):26-33.  
ISSN: 2449-2043.  
Available:<https://doi.org/10.25240/TJANS.2018.2.2.04>
  19. Cheek PR. Nutritional and Physiological Implications of Saponins: A Review, Canadian Journal of Animal Science. 1971;51(3):621-632.  
DOI: 10.4141/cjas71-082
  20. Punitha ISR, Shirwaika A, Shirwaikar A. Anti-diabetic activity of benzyl tetra isoquinoline alkaloid, berberine, in streptozotocin-nicotinamide induced type 2 diabetic rats. Diabetologia Croatica. 2005;34(4):117–1.
  21. Ohiagu FO, Chikezie PC, Maduka TDO, Enyoh CE, Chikezie CM. Bioactive compounds and medicinal usefulness of edible leaves of *Vernonia amygdalina*, *Ocimum gratissimum*, *Piper guineense* and *Gongronema latifolium*. SAJ Pharma Pharmacol 2021;7:101 SAJ Pharma Pharmacol 7: 101. ISSN: 2375-2262.
  22. Lemos AA., Lemos JA, Prado MA, Rodrigues MR, Silva MR.. Cockroaches as carriers of fungi of medical importance. Mycoses. 2006;49(1):23-5.  
DOI: 10.1111/j.1439-0507.2005.01179.x
  23. Hernández-Sánchez B, Santacruz E, Moore D and Sánchez C. 165 Hernández-Sánchez et al. Mexican Journal of Biotechnology 2021;6(1):165-189.  
Bioactive compounds from fungi with antiviral activities: Mechanism of action and biosynthetic pathways. Mexican Journal of Biotechnology, 2021;6(1):165-189.  
DOI: 10.29267/mxjb.2021.6.1.165
  24. Enemor VHA, Nnaemeka OJ1 and Okonkwo CJ. Minerals, Vitamins and Phytochemical Profile of *Gongronema latifolium*: Indices for Assessment of its Free Radical Scavenging, Nutritional, and Antinutritional Qualities. International Research Journal of Biological Sciences. 2014;3(1):17-21.  
ISSN 2278-3202.  
Available:<http://www.isca.in>, [www.isca.me](http://www.isca.me)
  25. Neave N. Hormones and Behaviour: A Psychological Approach. 2008; United Kingdom at the University Press, Cambridge, New York. 978-0-521-87145-7. ISBN 978-0-521-87145-7 hardback, ISBN 978-0-521-69201-4 paperback.  
DOI: 10.1017/CBO9780511808203
  26. AraújoVA, Freitas FV, Clóvis JMA, Lino-Neto NJ. Morphology of male reproductive system of two solitary bee species (Hymenoptera: Apidae). Systematics, Morphology and Physiology. Neotrop. entomol. 2010;39(4):10-71.  
Available:<https://doi.org/10.1590/S1519-566X2010000400020>
  27. Kotta S, Ansari SH, Ali J. Exploring scientifically proven herbal aphrodisiacs.

- Pharmacognosy Reviews. 2013;7(13):1-10.  
DOI: 10.4103/0973-7847.112832
28. Manortey S, Mensah PA, Acheampong GK, Dixit V. Evaluating Factors Associated with the Use of Aphrodisiacs among Adult Male Residents in Ashaiman Municipality, Ghana. Open Access Library Journal. 2018;5(9):1-13.  
DOI: 10.4236/oalib.1104876
  29. Thakur M, Chauhan NS, Bhargava S. A Comparative Study on Aphrodisiac Activity of Some Ayurvedic Herbs in Male Albino Rats. Archives of Sexual Behavior. 2009;38(6):1009-15.  
DOI: 10.1007/s10508-008-9444-8
  30. Mehrdad A, Ali R, Moravveji H, Mirkhani S, Namazi HW. Adherence to American Society of Health-System Pharmacists surgical antibiotic prophylaxis guidelines in Iran. Infect Control Hosp Epidemiol. 2006;27(8):876-8.  
Epub PMID: 16874651.  
DOI: 10.1086/506405
  31. Jian Feng C, Peng Ying Z, Cheng Wei X, Tao Tao H, Yun Gui B, Kao Shan C. Effect of aqueous extract of *Arctium lappa* L. (burdock) roots on the sexual behavior of male rats. BMC Complement Altern Med. 2012;12:1-8.  
[DOI:10.1186/1472-6882-12-8] [PMID] [PMCID]
  32. Fouche G, Afolayan AJ, Wintola OA, Khorombi TE, Senabe J. Effect of the aqueous extract of the aerial parts of *Monsonia angustifolia* E. Mey. Ex A. Rich., on the sexual behaviour of male Wistar rats. BMC Complement Altern Med. 2015;15:343.  
[DOI:10.1186/s12906-015-0880-4] [PMID] [PMCID]
  33. Asuzu CU. Bitter herbs of eastern Nigeria (*Gongronema latifolium*, *Vernonia amygdalina* and *Vitex doniana*): A REVIEW. Afr J Tradit Complement Altern Med. 2018;15(3):47-56.  
Available: <https://doi.org/10.21010/ajtcam.v15i3.647>
  34. Gauthaman K, Adaikan PG. The hormonal effects of *Tribulus terrestris* and its role in the management of male erectile dysfunction--an evaluation using primates, rabbit and rat. Phytomedicine. 2008;15(1-2):44-54.  
DOI: 10.1016/j.phymed.2007.11.011  
PMID: 18068966
  35. SA and Mishra SH. Aphrodisiac Studies of *Tricholepis glaberrima*. with Supportive Action from Antioxidant Enzymes. Pharmaceutical Biology. 2007;45(7):580-586.  
Available: <https://doi.org/10.1080/13880200701501326>
  36. Adimoelja A. Phytochemicals and the breakthrough of traditional herbs in the management of sexual dysfunction. International Journal of Andrology. 2000;23(2):82-4.  
DOI: 10.1046/j.1365-2605.2000.00020.x
  37. Neill DD, Vogel G, Hagler M, Hennessey A. Diminished sexual activity in a new animal model of endogenous depression. Neuroscience and Biobehavioral Reviews. 1990;14(1):73-76.  
Available: [https://doi.org/10.1016/S0149-7634\(05\)80162-9](https://doi.org/10.1016/S0149-7634(05)80162-9)
  38. Dasofunjo K, Asuk A, and Nku Ci. Evaluating the effect of ethanol leaf extract of *Gongronema latifolium* on some reproductive hormones of male Wistar rats. GSC Biological and Pharmaceutical Sciences. 2020;12(03):166-173.  
DOI: <https://doi.org/10.30574/gscbps.2020.12.3.0297>
  39. Varsha Z, Dinesh D, Vaibhao T, Shital P. Evaluation of potential aphrodisiac activity of *Moringa oleifera* seed in male albino rats. Int J of Pharmacy and Pharmaceutical Sci. 2013;5(4):683-689.

© 2022 Effiong et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/84266>