



Nephro-protective Efficacy of African Locust Bean Seed against Potassium Bromate-induced Renal Damage

I. O. Abali^a, N. M. Chika-Igwenyi^b, F. U. Agu^c, C. A. Onyeaghala^d, S. F. Orji^b, C. N. Ugwu^e, C. Igwenyi^b, C. L. Uche^f, O. I. N. Onyekachi^g, M. U. Nwobodo^b, C. E. Iwuoha^h and A. I. Airaodion^{i*}

^a Department of Surgery, Abia State University, Uturu, Nigeria.

^b Department of Internal Medicine, Alex Ekwueme Federal University Teaching Hospital, Abakaliki, Ebonyi State, Nigeria.

^c Department of Physiology, Gregory University, Uturu, Abia State, Nigeria.

^d Department of Internal Medicine, University of Port-Harcourt Teaching Hospital, Rivers State, Nigeria.

^e Department of Internal Medicine, Ebonyi State University, Abakaliki, Nigeria.

^f Department of Haematology, Abia State University, Uturu, Nigeria.

^g Department of Medical Microbiology, Ebonyi State University, Abakaliki, Nigeria.

^h Department of Community Medicine, Abia State University, Uturu, Nigeria.

ⁱ Department of Biochemistry, Federal University of Technology, Owerri, Imo State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJBGMB/2022/v12i3264

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/92032>

Original Research Article

Received 09 July 2022
Accepted 13 September 2022
Published 05 October 2022

ABSTRACT

Background: Potassium bromate (KBrO₃) has been utilized extensively to sterilize water, dye hair, oxidize food, add to bread dough as a maturing agent, and condition wheat dough, thereby entering into human body. It has been claimed to cause a number of toxicities including nephrotoxicity.

Aim: This study, therefore, sought to investigate the nephro-protective efficacy of African locust bean (ALB) seed against potassium bromate-induced renal damage.

Methodology: Using a Soxhlet extractor with ethanol as the solvent, ALB was extracted. Twenty-

*Corresponding author: E-mail: augustineairaodion@yahoo.com;

four mature male Wistar rats were randomly divided into groups A, B, C, and D after being acclimated in the lab. Oral distilled water was administered to Group A. Although groups C and D likewise received 100 and 200 mg/kg body weight of ALB, respectively, the animals in groups B, C, and D received 100 mg/kg body weight of potassium bromate. Rats received daily doses of freshly produced potassium bromate and ALB extract by oral gavage. Blood and kidney sample were taken after the prescribed 28-day course of medication. Standard techniques were used to assess renal biomarkers.

Results: When compared to the control group, potassium bromate treatment led to significant ($P < 0.05$) increases in the serum levels of creatinine, urea, uric acid, sodium (Na^+), potassium (K^+), chloride (Cl^-), and bicarbonate (HCO_3^-). Renal levels of tumor necrosis factor-alpha (TNF- α) and interleukin-6 (IL-6) were likewise elevated by KBrO_3 poisoning in comparison to the control group. However, combined administration of KBrO_3 and ALB seed extract resulted in significant dose-dependent reductions in the levels of all kidney biomarkers examined, with 200 mg/kg being the most effective dose.

Conclusion: This study revealed that potassium bromate indeed induced nephrotoxicity by unhooking renal biomarkers investigated in this study. It was further observed that seed extract of African locust bean (ALB) alleviated these adverse effects on the kidney by resisting the perturbations, thereby exhibiting nephron-protective effect. The biochemical composition of ALB has conferred it with this ability.

Keywords: African locust bean; nephron-protective efficacy; potassium bromate; renal damage.

1. INTRODUCTION

The usage of medicinal plants as a source of medication and to enhance human health dates back thousands of years. A number of plants contain bioactive components including phenolic and polyphenolic chemicals that control different immune pathways. Herbs that are high in flavonoids and phenolic compounds may also be anti-inflammatory and antioxidant [1]. African locust bean (*Parkia biglobosa*) is one of these plants. It is a perennial tree that belongs to the Leguminosae genus of legumes [2]. The plant's seeds are encased in an edible pulp that is yellowish, mealy, and sweet-tasting [3]. It is common knowledge that this plant contains a significant number of phenolic chemicals [4]. Epigallocatechin, epicatechin 3-O-gallate, and epigallocatechin 3-O-gallate were all present in the plant's bark [5]. Heart and saponin glycosides are present in leaf extract [6]. Protein and lactose are abundant in the fruit's pulp and seeds [5]. Phytate, tannin, oxalate, and hydrogen cyanide are a few antinutritional components found in seeds [2]. The extract of African locust bean (ALB) is antibacterial [4], antidiabetic [7], antifungal [8], anti-inflammatory [9], anti-diarrheal [10], anti-hypertensive [11], hypoglycemia [12], hypolipidemic [13], and hepatoprotective [14] in addition to having many beneficial effects. A recent study by Ezirim et al. [15] shown that ALB seed can alleviate testicular toxicity caused by potassium bromate.

To improve the quality of meals, certain compounds known as food additives are added during processing [16]. Food additives can generally be divided into two categories: those that are purposefully added to food and those that contaminate food in minute quantities as a result of handling and storage procedures [17]. A maximum limit has also been established for each additive by the Codex Alimentarius [18] Commission in consideration of any potential harmful consequences [17]. Potassium bromate (KBrO_3) is a white, crystalline salt that has been used for many years in the baking and confectionery industries to improve the flavor of goods [16]. It is classified as a flour treatment agent by FAO/WHO [19]. Since the International Agency for Research on Cancer (IARC) categorized potassium bromate as a Category 2B carcinogen, its official ban on use has been contested [20]. According to toxicological research, potassium bromate is hazardous to organs and has the capacity to rupture cell plasma membranes as an oxidizing agent and cause cells to release their internal contents into the extracellular environment [21,22]. In experimental animals and humans, potassium bromate was found to cause kidney toxicity [23]. It has also been linked to DNA damage and lipid peroxidation [24]. Additionally, it has been suggested that potassium bromate may cause cancer [25,26]. Following the injection of potassium bromate to the membrane lining the abdominal cavity of laboratory animals, an increase in the incidence of both malignant and

benign tumors was seen [27]. Additionally, giving animals potassium bromate caused a substantial rise in malignancies of the thyroid, kidneys, and other organs [28]. Therefore, the purpose of this study was to determine whether extract of ALB seed can prevent kidney damage from potassium bromate.

2. MATERIALS AND METHODS

2.1 Collection and Extraction of ALB

The seeds of ALB were acquired at a local market in Ibadan, Nigeria, and were identified by a botanist. They were processed into powder using a mechanical blender after being sun-dried. The extraction was finished using a Soxhlet device and ethanol as the solvent in accordance with the procedures outlined by Airaodion et al. [29,30]. The ethanol was evaporated in a rotary evaporator at 35°C with a yield of 2.55 g and a percentage yield of 10.20 percent. The extract was kept in the fridge until it was required.

2.2 Animal Treatment

The experiment involved twenty-four (24) mature male Wistar rats (*Rattus norvegicus*) weighing between 140 and 160 g. Prior to the study, they were acclimated in a lab setting for seven (7) days. The rats were kept in cages made of wire mesh, and they had unrestricted access to commercial rat food and water. The animals were housed at regular temperatures and humidity levels with 12-hour light and dark cycles. The Declaration of Helsinki and the regulations set by the Committee for the Purpose of Control and Supervision of Experiments on Animals were followed in conducting this investigation. Additionally, while using animals in research, NIH guideline was followed [31]. They were divided into groups A, B, C, and D at random. As usual, Group A received oral distilled water. The animals in groups B, C, and D received 100 mg/kg body weight of potassium bromate whereas groups C and D also received 100 and 200 mg/kg body weight of ALB, respectively. Rats were given daily oral gavages of ALB and fresh potassium bromate. Twenty-four hours after the last dose, the animals were slaughtered while being softly sedated with diethyl ether. Blood was drawn by puncturing the heart.

2.3 Renal Homogenates Preparation

The kidneys were removed from the animal after the sacrifice, cut in half, and maintained in an

ice-cold container to prepare the renal homogenates. The cortex and medulla were carefully cut apart using a sharp scalpel, and each was homogenized separately in a glass Teflon homogenizer in a solution of 2 mM Tris-HCl and 50 mM mannitol buffer at pH 7.0 to create a homogenate that was 10% (w/v). After being diluted to 5% with Tris-mannitol buffer, these homogenates were subjected to high speed homogenization (20,000 rpm) in an Ultra Turrex Kunkel homogenizer. The renal homogenate was used to determine interleukin-6 (IL-6) and tumor necrosis factor-alpha (TNF- α) concentrations.

2.4 Determination of Renal Biomarkers

Using a kit from Randox Laboratories Ltd. in the UK and the diacetyl monoxime method, serum urea was quantified. The level of uric acid was determined using a kit from Linear Chemicals Barcelona in Spain using the quinoneimine dye complex, whereas the level of creatinine was determined using kits from Randox Laboratories Ltd. in the UK based on its reaction with saturated picric acid to produce a yellow-red complex. Using kits from Teco Diagnostics in Anaheim, California, the serum concentrations of the ions sodium (Na⁺), potassium (K⁺), chloride (Cl⁻), and bicarbonate (HCO₃⁻) were all measured by spectrophotometric analysis. Rat ELISA kits with monoclonal antibodies specific for rat TNF- α and IL-6 were used, according to Mohamed and Saddek [32].

2.5 Statistical Analysis

Every piece of data is presented as mean \pm standard deviation. To examine the data by comparing the outcomes of the treatment groups to the control group, analysis of variance (ANOVA) was utilized in conjunction with a Post-Hoc test (Tukey's comparison test) using Graph Pad Prism software. All deviations with $p \leq 0.05$ were deemed significant.

3. RESULTS

When compared to the control group, potassium bromate treatment led to significant ($P \leq 0.05$) increases in the serum levels of creatinine, urea, uric acid (Table 1), sodium (Na⁺), potassium (K⁺), chloride (Cl⁻), and bicarbonate (HCO₃⁻) (Table 2). Renal levels of tumor necrosis factor-alpha (TNF- α) and interleukin-6 (IL-6) were likewise elevated by KBrO₃ poisoning in comparison to the control group (Table 3). However, combined

Table 1. Effect of ALB seed on serum creatinine, urea and uric acid concentrations of potassium bromate induced nephrotoxicity

Treatment Group	Creatinine (mg/dL)	Urea (mg/dL)	Uric Acid (mg/dL)
Control	0.89±0.01	19.64±1.29	4.78±0.78
100 mg/kg KBrO ₃ only	1.47±0.02	32.67±4.08	7.77±0.93
100 mg/kg KBrO ₃ + 100 mg/kg ALB	1.21±0.00	28.28±1.34	6.05±1.05
100 mg/kg KBrO ₃ + 200 mg/kg ALB	1.03±0.01	23.94±2.92	5.25±0.28
p-value	0.05	0.00	0.02

Results are presented as mean±Standard deviation (SD) with n = 6

Table 2. Effect of ALB seed on plasma electrolytes concentrations of potassium bromate induced nephrotoxicity

Treatment Group	Sodium (mEq/L)	Potassium (mmol/L)	Chloride (mmol/L)	Bicarbonate (mEq/L)
Control	137.34±9.28	3.96±0.82	97.78±6.24	26.93±3.26
100 mg/kg KBrO ₃ only	178.36±11.02	6.02±1.92	123.65±9.19	33.62±2.97
100 mg/kg KBrO ₃ + 100 mg/kg ALB	164.04±8.29	5.34±0.83	111.18±9.33	31.58±5.13
100 mg/kg KBrO ₃ + 200 mg/kg ALB	143.78±9.54	4.73±0.26	102.94±7.16	28.67±5.83
p-value	0.04	0.01	0.01	0.05

Results are presented as mean±SD with n = 6

Table 3. Effect of ALB Seed on tumor necrosis factor- α (TNF- α) and interleukins-6 (IL-6) of potassium bromate induced nephrotoxicity

Treatment Group	TNF- α (pg/mL)	IL-6 (pg/mL)
Control	3.67±0.48	5.93±0.74
100 mg/kg KBrO ₃ only	6.18±1.05	8.63±1.06
100 mg/kg KBrO ₃ + 100 mg/kg ALB	4.84±0.36	7.53±0.72
100 mg/kg KBrO ₃ + 200 mg/kg ALB	3.92±0.78	6.36±1.01
p-value	0.01	0.03

Results are presented as mean±SD with n = 6

administration of KBrO₃ and ALB seed extract resulted in significant dose-dependent reductions in the levels of all kidney biomarkers examined, with 200 mg/kg being the most effective dose.

4. DISCUSSION

Plants have long been utilized as folk medicine to treat a variety of illnesses. Even today, medical professionals and village doctors in many nations use unprocessed plant extracts to treat human illnesses [33,34]. To assess the effectiveness of various plants against chemically induced toxicity, experimental work has been done on them [35,36].

It may be possible to identify renal impairment brought on by KBrO₃ poisoning by measuring serum levels of urea, creatinine, and uric acid, which were previously utilized as early indicators of renal failure [37,38]. The significant increase in blood urea, creatinine, and uric acid in the KBrO₃-treated rats suggests the onset of renal failure and a reduced glomerular filtration rate.

Acute renal failure was the outcome of accidental potassium bromate poisoning, according to Adeleke and Asani [39].

Increases in blood creatinine and urea have been identified as key biomarkers of kidney disease and the degradation of the integrity of the renal tubules. These changes are damage indicators for inadequate glomerular filtration [40,41]. Increased levels of creatinine in the serum of rats treated with KBrO₃ supported earlier claims that KBrO₃ consumption results in renal injury [42,43].

Urea, which is largely made in the liver and released by the kidneys, is the main byproduct of protein catabolism [44]. It is the main method of getting harmful ammonia out of the body. The medical doctor can evaluate a patient's renal function using urea determination extremely well [37,38]. Increased urea levels are typically linked to nephritis, renal ischemia, obstruction of the urinary tract, and several extra-renal disorders.

Similar to other findings [45,46], potassium bromate-induced elevation in urea concentration were seen in this investigation. Additionally, renal failure has been connected to increased urea levels, according to Akomolafe et al. [47].

Degradation of purines and pyrimidines may potentially be a contributing factor to the rise in uric acid levels. According to a study by Mohamed and Saddek [32], KBrO_3 increased xanthine oxidase activity, which resulted in an excess of uric acid being produced. This process may have led to the action of KBrO_3 in the increase of uric acid. ALB extract reduced this KBrO_3 -induced effect on uric acid in a dose-dependent manner. The hypouricemic effects of ALB seed extract may be brought on by the presence of flavonoids, which have inhibitory effects on the enzyme activities of xanthine oxidase and xanthine dehydrogenase [48].

For animal species to survive, electrolytes are necessary [49]. A considerable increase in several electrolytes, including sodium (Na^+), potassium (K^+), chloride (Cl^-), and bicarbonate (HCO_3^-) ions, was also brought about in the current investigation by treatment with KBrO_3 . This is a sign of significant kidney injury, which was also seen in kidney parameter pathological differences. The same outcomes were described by previous studies [50,51]. The observed increases in the serum concentrations of urea, creatinine, and uric acid were supported by the elevated levels of Na^+ , K^+ , HCO_3^- , and Cl^- . The kidney is in charge of controlling different electrolytes and preserving homeostasis [52]. For instance, Na^+ and K^+ are important components of extracellular and intracellular fluids, respectively. Since the kidney controls both physiological states, elevated levels of these electrolytes may suggest renal failure, especially at the glomerular and tubular levels. The transit of various non-electrolytes and the maintenance of the water equilibrium level in physiological tissue are both significantly influenced by sodium and potassium [53,54]. Additionally, as HCO_3^- and Cl^- are reabsorbed in the proximal tubule, a high level of these electrolytes may indicate tubular renal damage. This outcome is consistent with the rise in all electrolytes observed by Kanadi et al. [55] after exposure of animals to KBrO_3 . The main anion in serum is chloride. Extracellular sodium and bicarbonate concentrations have an impact on extracellular chloride concentration [49]. However, concurrent treatment of KBrO_3 and ALB seed extract returned the serum levels of all the electrolytes

examined to normal, particularly in the group treated with 200 mg/kg, whose values were not substantially different from those in the normal control group. This raises the possibility of a preventive effect of ALB seed against renal damage brought on by KBrO_3 .

Interleukin-6 (IL-6) and tumor necrosis factor ($\text{TNF-}\alpha$) are two pro-inflammatory cytokines that are hypothesized to play a key role in the pathogenesis of chronic kidney diseases [56]. By activating transcription factors, which in turn cause the release of pro-inflammatory cytokines like IL-6 and $\text{TNF-}\alpha$, reactive oxygen species (ROS) can trigger inflammatory processes [57]. The findings of this investigation showed that KBrO_3 caused an inflammatory response that was manifested by an increase in renal $\text{TNF-}\alpha$ and IL-6 levels, indicating nephrotoxicity. This is comparable to the increase in $\text{TNF-}\alpha$ and IL-6 levels found in rats exposed to potassium bromate, as reported by Mohamed and Saddek [32]. Recently, Ugwu et al. [58] found that KBrO_3 caused oxidative stress in the rat kidney. The large rise in renal $\text{TNF-}\alpha$ and IL-6 levels may be a result of the kidneys producing more ROS. The overproduction of proinflammatory cytokines $\text{TNF-}\alpha$ and IL-6 in renal tissue was however suppressed by ALB seed therapy and KBrO_3 administration, which had a protective effect against pathological kidney changes. The ability of ALB seed to attenuates nephrotoxicity caused by potassium bromate intoxication could be as a result of its rich phytochemical constituents and antioxidant potential [58].

5. CONCLUSION

This study revealed that potassium bromate indeed induced nephrotoxicity by unhooking renal biomarkers investigated in this study. It was further observed that seed extract of African locust bean (ALB) alleviated these adverse effects on the kidney by resisting the perturbations, thereby exhibiting nephron-protective effect. The biochemical composition of ALB has conferred it with this ability.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Khan RA, Khan MR, Sahreen S, Shah NA, Ahmad B, Khan AM. Alteration of renal function by potassium bromate (KBrO₃): Protective effects of *Launaea procumbens*. African Journal of Pharmacy and Pharmacology. 2012;6(19):1400-1404.
2. Olabinri BM, Adetutu A, Olaleye MT, Oluwafunsho BO, Oyeniyi OO. A study of the antioxidative potentials of acetone and aqueous extracts of *Parkia biglobosa* and *Tetracarpidium conophorum* stem barks *in vitro*. International Journal of Medicine and Medical Sciences. 2013;5(8):368-373.
3. Aliero BL, Umaru MA, Suberu HA, Abubakar A. A handbook of common plants in Northwestern Nigeria, Sokoto University Press, Nigeria. 2001;130.
4. Miollogo-Kone H, Guissou IP, Nacoulma O, Traore AS. Comparative study of Leaf and stem bark extracts of *Parkia biglobosa* against enterobacteria. Afr. J. Trad. Compl. Altern. Med. 2008;5:238-243.
5. Alabi DA, Akinsulire OR, Sanyaolu MO. Qualitative determination of Chemical and Nutritional Composition of *ALB* (Jacq) Benth. Afr. J. Biotechnol. 2005;4(8):812-815.
6. Miollogo-Kone H, Lompo M, Kini F, Asimi S, Guissou IP, Nacoulma O. Evaluation of flavonoids and total phenolic contents of stem bark and leaves of *Parkia biglobosa* (Jacq) Benth (Mimosaceae) – free radical scavenging and antimicrobial activities. Res. J. Med. Sci. 2009; 3(2):70-74.
7. Odetola AA, Akinloye O, Egunjobi C, Adekunle WA, Ayoola AO. Possible antidiabetic and antihyperlipidaemic effects of fermented *Parkia biglobosa* (Jacq) extract in alloxan-induced diabetic rats. Clin Exp. Pharmacol. Physiol. 33(9):808-812.
8. Kouadio F, Kanko C, Juge M, Grimaud N, Jean A, N'Guessan YT, Petit JY. Analgesic and antiinflammatory activities of an extract from *Parkia biglobosa* used in traditional medicine in the Ivory Coast. Phytother. Res. 2000;14:635-637.
9. Fawole OB, Abioye OO. Comparative effectiveness of husk of *Parkia biglobosa* wild Bent. Mitta and Benomyl in controlling Fusarium wilt disease of Cowpea. J. Nig. Soc. Exper. Biol. 2002; 2(1):057-063.
10. Agunu A, Yusuf S, Gabriel AO, Abdulkadir UZ, Ezzeldin AM. Evaluation of five medicinal plants used in diarrhoea treatment in Nigeria. J. Ethnopharm. 2005;101:27-30.
11. Airaodion AI, Ogbuagu EO. Ameliorative effect of *Parkia biglobosa* (African locust bean) against egg-yolk induced hypertension. International Journal of Bio-Science and Bio-Technology. 2020; 12(5):17-25.
12. Airaodion AI, Airaodion EO, Ogbuagu EO, Ogbuagu U, Osemwowa EU. Effect of Oral Intake of African Locust Bean on Fasting Blood Sugar and Lipid Profile of Albino Rats. Asian Journal of Research in Biochemistry. 2019;4(4): 1-9.
13. Ugwu CN, Iwuoha CE, Chika-Igwenyi NM, Onyeaghala CA, Orji SF, Igwenyi C, Uche CL, Onyekachi OIN, Nwobodo MU, Abali IO, Airaodion AI. Chemotherapeutic Propensity of Africa locust bean (*Parkia biglobosa*) Seed on Lipid Profile against Potassium Bromate-induced Cardiotoxicity. Journal of Applied Life Sciences International. 2022;25(8):24-31.
14. Airaodion AI, Ogbuagu EO, Ogbuagu U, Adeniji AR, Agunbiade AP, and Airaodion EO. Hepatoprotective effect of *Parkia biglobosa* on acute ethanol-induced oxidative stress in Wistar rats. International Research Journal of Gastroenterology and Hepatology. 2019c;2(1): 1-11.
15. Ezirim EO, Uche CL, Abali IO, Iwuoha CE, Chika-Igwenyi NM, Onyeaghala CA, Orji SF, Ugwu CN, Ugwu NI, Igwenyi C, Airaodion AI. Therapeutic Potential of *Parkia biglobosa* Seed against Potassium Bromate-induced Testicular Toxicity. International Journal of Research and Reports in Gynaecology. 2022;5(3):78-89.
16. Abdulmumeen HA, Risikat AN, Sururah AR. Food: Its preservatives, additives and applications. International Journal of Chemical and Biochemical Sciences. 2012;1:36-47.
17. Achukwu PU, Ufelle SA, Ukaejiofo EO, Nwachukwu DN. The effect of potassium bromate on some haematological parameters of Wistar rats. Nigeria Journal of Physiology and Science. 2009;24(1):59–61.
18. CODEX Alimentarius Commission. Joint FAO/WHO Food Standards Programme. Forty-first Session Rome, Italy, 2 – 6 July 2018. Report of the 50th session of the CODEX Committee on Food Additives, Xiamen, China. 26 – 30 March 2018.
19. Oladije JE, Apeh DO, Idih FM, Shaibu IE, Shuaibu HT, Ameh DO. Effects of Ethanol

- Extract of Pterocarpus Erinaceus Leaves and Potassium Bromate Administration on Some Biochemical and Haematological Indices in Albino Rats. EAS Journal of Pharmacy and Pharmacology. 2020; 2(5):8-14.
20. IARC. Potassium Bromate – Summary of Data Reported and Evaluation. 1999;73:481.
Available:<http://www.inchem.org/document/s/iarc/vol73/73-17.html>
 21. Akanji MA, Nafiu MO, Yakubu MT. Enzyme activities and histopathology of selected Tissues in rats treated with potassium bromate. African Journal of Biomedical Research. 2008;11:87–95.
 22. Olajide JE, Ojogbane, BE, Ayeni C. Protective activity of acetylsalicylic acid on biochemical markers of hepatic and renal functions in rats administered with potassium bromate. Biolife. 2015;3(1):317–326.
 23. Uchida HA, Sugiyama H, Kanehisa S, Harada K, Fujiwara K, Ono T, Makino H. An elderly patient with severe acute renal failure due to sodium bromate intoxication. Internal Medicine. 2006;45(3): 151-154.
 24. Adekoya AO, Adekoya BJ, Desalu OO, Aderibigbe A. Patterns of lipid profile in adult nephritic syndrome patients in Nigeria. International Journal of Biology and Medical Research. 2011;2(4):954–960.
 25. Sai K, Umemura T, Takagi A, Hasegawa R, Kurokawa Y. The protective role of glutathione, cysteine and vitamin C against oxidative DNA damage induced in rat kidney by potassium bromate. Jpn J. Cancer Res. 1992;83:45–51.
 26. Chipman JK, Davies JK, Parson JL, Nair J, O'Neill G, Fawell JK. DNA oxidation by potassium bromate; a direct mechanism or linked to lipid peroxidation. Toxicol. 1998; 126:93-102.
 27. Kurokawa Y, Aoki S, Matsushima Y, Takamura N, Imazawa T, Hayashi Y. Dose-response studies on the carcinogenicity of potassium bromate in F344 rats after long-term oral administration. Journal of the National Cancer Institute. 2016;77(4):977-982.
 28. DeAngelo AB, George MH, Kilburn SR, Moore TM, Wolf DC. Carcinogenicity of potassium bromate administered in the drinking water to male B6C3F1 mice and F344/N rats. Toxicologic Pathology. 2018; 26(5):587-594.
 29. Airaodion AI, Ngwogu AC, Ekenjoku JA, Ngwogu KO. Hepatoprotective potency of ethanolic extract of *Garcinia kola* (Heckel) seed against acute ethanol-induced oxidative stress in Wistar rats. International Research Journal of Gastroenterology and Hepatology. 2020;3(2):1-10.
 30. Airaodion AI, Ogbuagu EO, Ogbuagu U, Awosanya OO, Airaodion EO. Effect of Methanolic extract of *Corchorus olitorius* leaves on hypoglycemic and hypolipidaemic activities in albino rats. Asian Plant Research Journal. 2019;2(4):1-13.
 31. National Research Council. Guide for the Care and Use of Laboratory Animals, 8th ed. The National Academies Press: Washington, DC, USA; 2011.
 32. Mohamed EAK, Saddek EA. The protective effect of taurine and/or vanillin against renal, testicular, and hematological alterations induced by potassium bromate toxicity in rats. The Journal of Basic and Applied Zoology. 2019;80(3):1-11.
 33. Airaodion AI, Ekenjoku JA, Ngwogu KO, Ngwogu AC. Consumption of coconut (*Cocos nucifera* L.) water improved fertility parameters in male Wistar rats. Asian Journal of Pregnancy and Childbirth. 2019; 2(3):1-7.
 34. Airaodion AI, Ogbuagu EO, Ekenjoku JA, Okoroukwu VN, Ogbuagu U, Airaodion EO. Antifertility Effect of Ethanolic Leaf Extract of *Carica papaya* in Male Wistar Rats. Merit Research Journal of Medicine and Medical Science. 2019;3(9):121-129.
 35. Airaodion AI, Ogbuagu EO, Ekenjoku JA, Ogbuagu U, Okoroukwu VN. Antidiabetic effect of ethanolic extract of *Carica papaya* leaves in alloxan-induced diabetic rats. American Journal of Biomedical Science & Research. 2019;5(3):227-234.
 36. Airaodion AI, Akaninyene IU, Ngwogu KO, Ekenjoku JA, Ngwogu AC. Hypolipidaemic and antidiabetic potency of *Allium cepa* (onions) bulb in alloxan-induced diabetic rats. Acta Scientific Nutritional Health. 2020;4(3):1-8.
 37. Ogbuagu EO, Airaodion AI, Ogbuagu U, Nweke IN, Unekwe PC. Nephrotoxicity of ethanol extract of *Xylopiya aethiopicum* fruit in Wistar rats. International Journal of Advances in Nephrology Research. 2021; 4(1):1-16.

38. Airaodion AI, Ngwogu KO, Ngwogu AC, Megwas AU, Ekenjoku JA. Nephrotoxicity of Monosodium Glutamate (MSG) in Wistar Rats. *International Journal of Advances in Nephrology Research*. 2020;3(1):1-10.
39. Adeleke SI, Asani MO. Accidental potassium bromate poisoning causing acute renal failure. *Online J Health Allied Scs*. 2009;8(1):11
40. Airaodion AI, Alabi OJ, Ogbuagu EO, Atiba FA, Olawoyin DS. Nephro and hepatotoxic effect of air-freshener in Wistar rats. *Asian Journal of Research in Nephrology*. 2020; 3(2):1-9.
41. Airaodion AI, Megwas AU, Ekenjoku JA, Ngwogu KO, Ngwogu AC. Nephro- and hepato-toxicity of common household insecticides used in Nigeria. *International Research Journal of Gastroenterology and Hepatology*. 2020;3(2):21-28.
42. Bao L, Yao XS, Tsi D, Yau CC, Chia CS, Nagai H. Protective effects of bilberry (*Vaccinium myrtillus L.*) extract on KBrO₃-induced kidney damage in mice. *J. Agric. Food Chem*. 2008;56: 420–425
43. Altoom NG, Ajarem J, Allam AA, Maodaa SN, Abdel-Maksoud MA. Deleterious effects of potassium bromate administration on renal and hepatic tissues of Swiss mice. *Saudi Journal of Biological Sciences*. 2018;25:278–284.
44. Airaodion AI, Ogbuagu EO. Effect of *Cyperus esculentus L.* (tiger nut) milk on hepatic and renal indices of Wistar rat. *Asian Journal of Research in Nephrology*. 2020;3(2):10-16.
45. Akanji MA, Nafiu MO, Yakubu MT. Enzyme and histopathology of selected tissues in rats treated with potassium bromate. *Afr. J. Biomed. Res*. 2008;11:87-95.
46. Oyewole OI. Chemopreventive role of vitamin C and E on potassium bromate induced renal oxidative damage in rat. *Journal of Medicine and Medical Sciences*. 2011;2(11):1189-1192.
47. Akomolafe SF, Olasehinde TA, Adewale, OO, Ajayi OB. Curcumin improves biomolecules associated with renal function and attenuates oxidative injury and histopathological changes in potassium-induced toxicity in rats' kidney. *Biological Trace Element Research*. (Epub ahead of print; 2020.
48. Rahmat RF, Adnan S, Anugrahwaty R, Alami EPS, Siregar B. Red onion growth monitoring system in hydroponics environment. In *Journal of Physics: Conference Series*. 2019;1235(1): 012117.
49. Gazuwa SY, Dabak JD, Jaryum KH Oluwa I. Relationship between dose and duration of administration of potassium bromate on selected electrolytes and hepatorenal parameters in male albino Wistar rats. *European Journal of Nutrition & Food Safety*. 2019;11(4): 214-220.
50. Khan N. *Nigella sativa* (black cumin) ameliorates potassium bromate-induced early events of carcinogenesis: diminution of oxidative stress. *Human & Experimental Toxicology*. 2003;22:193-203.
51. Kermani NM, Abushofa FA, Aldaek AM, Jaat FG. Protective Effect of *Allium cepa L.* (onion) Against Potassium Bromate-Induced Hematological, Biochemical and Histopathological Alterations in Rats. *International Journal of Innovative Science and Research Technology*. 2020;5(11): 201-207.
52. John AD, Larry AW. The autonomic nervous system and renal physiology. *International Journal of Nephrology and Renovascular Disease*. 2013;6:149-160.
53. Airaodion AI, Onabanjo KS. Effect of fortifying zobo (*Hibiscus sabdarifa*) drink with pineapple and watermelon on the mineral compositions and microbial quality. *Merit Research Journal of Food Science and Technology*. 2022;7(1):001-007.
54. Airaodion AI, Ayita EB, Oluba SO, Emaleku SA, Osunmuyiwa OJ. Chemical composition and nutraceutical potential of velvet tamarind fruit pulp. *Asian Journal of Biochemistry, Genetics and Molecular Biology*. 2021;3(5):22-31.
55. Kanadi MA, Wudil AM, Alhassan AJ, Ngwen AL, Yaradua AI, Nasir A. Dose-dependent chemopreventive effect of methanol extract of *Carica papaya* seed on potassium bromate-induced nephrotoxicity in rats. *Asian Journal of Biochemistry, Genetics and Molecular Biology*. 2019; 2(1):1-12.
56. Stenvinkel P, Ketteler M, Johnson RJ. IL-10, IL-6, and TNF alpha: central factors in the altered cytokine network of uremia—the good, the bad, and the ugly. *Kidney International*. 2005;67:1216–1233.
57. Valério DA, Georgetti SR, Magro DA, Casagrande R, Cunha TM, Vicentini FT.

- Quercetin reduces inflammatory pain: Inhibition of oxidative stress and cytokine production. *Journal of Natural Products*. 2009;7211975–11979.
58. Ugwu CN, Abali IO, Iwuoha CE, Chika-Igwenyi NM, Onyeaghala CA, Orji SF, Igwenyi C, Uche CL, Onyekachi OIN, Nwobodo MU, Airaodion AI. Ameliorative effect of *Parkia biglobosa* (African locust bean) seed against potassium bromate-induced oxidative stress. *Merit Research Journal of Medicine and Medical Science*. 2022; 10(8):213-219.

© 2022 Abali et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/92032>