



Screening of *Mucuna pruriens* (L.) DC. and *Pongamia pinnata* (L.) Pierre (Fabaceae) Minerals on Biochemical Parameters of *Oryctolagus cuniculus* as Antivenom

Obou Constantin Okou^{a,b*}, Djako Sosthène Thierry Akré^a,
Allali Eugène Koffi^a, Djané Hamed^a
and Allico Joseph Djaman^b

^a Department of Biochemistry and Microbiology, Jean Lorougnon Guédé University, Côte d'Ivoire.

^b Department of Biology-Health, Félix Houphouët-Boigny University Abidjan, Côte d'Ivoire.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJOB/2023/v17i4329

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/97893>

Original Research Article

Received: 25/01/2023

Accepted: 28/03/2023

Published: 10/04/2023

ABSTRACT

Aims: The objective of this study was to evaluate the antivenom action of minerals from two plants on the biochemical parameters of rabbits.

Methodology: For its realization, thirty-six (36) rabbits including nineteen (19) males and seventeen (17) females were divided into three portions (E1, E2 and E3). In each portion, there are different batches with three rabbits per batch. So, for E1, each rabbit of the control lot and those of

*Corresponding author: E-mail: constinokob@gmail.com;

the experimental lot were collected separately in the red tubes (dry tubes). However, those from the experimental batch were injected 20-30 minutes after the injection of *Naja nigricollis* venom. For E2, each rabbit in a given experimental batch was first scarified with a specific potion initially prepared and then separately harvested the following day. Then, each animal was separately envenomed and then separately sampled. In E3, the same procedure of E2 was observed with the only difference that this time, each animal of a given experimental lot was first separately envenomed and then scarified with a precise potion previously prepared and subsequently collected three days later. All the blood samples taken in the various red tubes were transported in a cooler for the determination of biochemical parameters.

Results: The venom has a dose-dependent action on the red blood cells *in vitro*, whereas the *in vivo* tests highlighted in a general way, the stabilizing and preventive action of the minerals from the potion P2 (*Mucuna pruriens*), P3 (*Pongamia pinnata*), P4, P5 and P6, and the curative action of P1 and P3 (*Pongamia pinnata*) on some biochemical parameters. Indeed, their action was manifested by the survival of some envenomed animals with a greater activity for P3.

Conclusion: For the development of an antivenom, it would be advisable to use more *Pongamia pinnata* because of its effectiveness both preventively and curatively.

Keywords: Antivenomous; minerals; *Mucuna pruriens*; *Pongamia pinnata*; biochemical parameters.

1. INTRODUCTION

“The annual number of ophidian bites exceeds 5 million and the number of people who die following an ophidian bite is 125,000” [1,2]. “Worldwide, 500,000 to 5,000,000 people are estimated to be victims of snakebite envenomation, of which 50,000 to 150,000 die and 400,000 suffer serious functional sequelae: amputation, renal failure, neurological sequelae” [3,2,4].

Ophidian envenomations in Africa represent a public health problem, due to their severity, but also because of the difficulties of their management [2]. “In fact, in urban Africa, envenomed patients can be treated with serotherapy (immunotherapy). It comes from the specific study of venoms from rigorous biochemical and toxicological analysis” [5]. “However, immunotherapy, which remains the only specific treatment for ophidian envenomation, is very expensive, available only in a few pharmacies and is constantly in short supply” [6]. “For rural populations, venomous snake bites are a medical, social and economic problem because of their low income and the distance from medical centers [7]. In the Maghreb as well as in sub-Saharan countries, the venomous snakes considered more dangerous are represented by the Viperidae and Elapidae” [8,9].

“*Naja melanoleuca*, *Naja nigricollis*, *Naja haje*... are part of the Elapidae found in Africa [10]. The *Naja nigricollis* has a venom containing toxins that can quickly immobilize a prey (neurotoxic)”

[11]. “These toxins are polypeptides and proteins, of small size (PM < 30 kDa) and represent 50 to 70% of the dry weight of Elapidae venoms” [11]. “These proteins can intervene alone or in synergy with other components of the venom to cause serious disruptions of biological systems of the body. These disturbances are at the origin of the edema, hematological, neurological, renal, pulmonary disorders and tissue necrosis characteristic of ophidian envenomation” [12,13,14]. These major facts push African populations to adopt traditional medicine as a therapeutic alternative.

“Numerous researches keep showing that medicinal plants contain various biologically active chemical principles that exert different pharmacological activities: antioxidant, anti-inflammatory, analgesic activities, etc” [15,16,17,18]. “Among these bioactive substances, some of them may have effects on undoubtedly biochemical parameters” [18].

“According to the World Health Organization (WHO) in 2013, about 80% of the populations of developing countries use traditional medicine and in particular herbal medicine for their health care needs. The African floral heritage is very rich in medicinal plants whose effectiveness is proven. As a matter of fact, it has been shown that the continent is full of nearly 5,000 medicinal species” [19,20,18].

“In West Africa, particularly in Benin, 80% of people bitten by snakes claim to use traditional treatment rather than modern Western medicine”

[21,7]. "In Côte d'Ivoire, studies conducted by some researchers have reported noteworthy results regarding the ethnomedical approach to treating snakebites with medicinal plants. According to these authors, the root of *Securidaca longepedunculata* Fresen. (Polygalaceae) can be used in the case of envenomations by Elapidae" [22-24]. In addition, in the Bouaké region (Côte d'Ivoire), some populations use two Fabaceae, namely *Mucuna pruriens* and *Pongamia pinnata* to treat cases of ophidian envenomations.

It is in the concern to exploit rationally this patrimony, to give a scientific basis to the use of these plants and to bring its contribution to the discovery of new heads of series of drugs that this present study was led. It aimed to know the antivenom effect *In vivo* of *Pongamia pinnata* and *Mucuna pruriens* on some biochemical parameters of rabbits.

2. MATERIALS AND METHODS

2.1 Materials

2.1.1 Biological materials

The plant material consisted of *Mucuna pruriens* and *Pongamia pinnata* (Fabaceae). They were harvested in the month of December 2019 in Bouaké (Central Côte d'Ivoire).

2.1.2 Animal material

For this study, thirty-six rabbits, including nineteen (19) males and seventeen (17) females of the Hyplus breed, aged two and a half months were purchased from a breeder in the locality of Daloa (Côte d'Ivoire). After the acclimatization period in the animal house of the Laboratory of Animal Physiology of the University Jean Lorougnon Guédé (UJLoG) of Daloa (Côte d'Ivoire), the weights of the rabbits varied between 1.45 and 2.4 Kg. Besides this animal model, viper skulls and *Naja nigricollis* venom were also used. The viper skulls were provided by a medicodrugist while the *Naja nigricollis* venom was supplied by the Pasteur Institute of Adiopodoumé (Côte d'Ivoire).

2.2 Methods

2.2.1 Method of preparation of minerals

For its realization, the various plants were harvested in Bouaké, washed, cut up and then

dried in the sun at room temperature for one week. Thereafter, the plant parts/organs were dried in an oven at a temperature of 70°C for three days. After this drying time, the organs (plant and animal) obtained were incinerated in a muffle furnace for 13 hours at 550°C. The ashes collected were weighed with a precision balance. They were smooth except for the viper skull which was rough. The colors varied from gray to brown.

The combination of the ashes from the various biological products resulted in the following potions:

- P1 was composed of the ashes of the two plants and the skull of viper;
- P2, P3 and P4 were constituted respectively and only of ash of *Mucuna pruriens*, *Pongamia pinnata* and skull of viper;
- P5 was formed by the ashes of *Mucuna pruriens* and *Pongamia pinnata*;
- P6 was consisted of the ashes of skull of viper and *Mucuna pruriens*;
- P7 was comprised of the ashes of viper skull and *Pongamia pinnata*.

2.2.2 Calculation of incineration efficiency

The formula below was used to calculate the dry matter weight of the organs used.

$$\text{Ac :Ash content} = \frac{\text{Mass of ash}}{\text{Dry matter}} \times 100$$

2.2.3 Method of scarification of experimental batches

To scarify the experimental batches, the following potions:

- P1 was used for batch 4;
- P2 was utilized for lot 5;
- P3 was solicited for lots 6;
- P4 has been used for batch 7;
- P5 was served for lot 8;
- P6 was required for lot 9;
- P7 has been used for lot 10.

There were two males and one female in each experimental batch. However, prior to scarification, the affected areas (toes of the left paw and tarsus of the right paw) were bared with a pair of scissors. After that, a separate amount of 0.45 mg of the previously prepared potion was applied to each affected area of each given

batch. The experimental tests continued for three days after the scarification (for the preventive test) and the curative test.

2.2.4 Blood collection method

In general, blood samples were taken at the Agrovalorisation Laboratory of the Jean Lorougnon Guédé University from the short saphenous vein and/or the femoral vein. The restraint method was performed by three individuals. The areas where these veins are located were previously exposed with a pair of scissors. The vacutainers in which the needles were inserted were used to collect the samples using red tubes (dry tubes). The tubes obtained were kept in a cooler containing ice and then transported to the laboratory for analysis.

2.2.5 *In vitro* hemolysis test method

To carry out this test, two control batches (batch 1 and batch 2) were formed. Batch 1 consisted of three males and batch 2 of three females. Among these control batches, a blood sample was taken from one of them in order to perform the *in vitro* hemolysis test of *Naja nigricollis* venom. To perform this test, ten (10) tubes were used including one (1) control tube and nine (9) experimental tubes. The stock solution was prepared in tube 1 by dissolving 1.6 mg of venom crystals in 1 mL of physiological water. In the remaining nine (9) tubes (tubes 2 to tubes 10), a volume of 0.5 mL of physiological water was added. The concentration ranges of venom were prepared using the double dilution technique of geometric reason 1/2. It consisted in taking 0.5 mL of the stock solution (tube 1) and transferring it to 0.5 mL of physiological water in tube 2 and then homogenizing it. This procedure was repeated until tube 9. From tube 9, a volume of 0.5 mL was taken and then discarded afterwards. Thus, the concentrations varied in the tubes from

1.6 mg/mL to $6.26 \cdot 10^{-3}$ mg/mL. To these nine (9) experimental tubes and to the control tube (tube 10), 5 drops of rabbit whole blood collected and then manually homogenized were added. After mixing, the whole preparation was incubated at room temperature for 30 to 40 minutes for microscopic observation. This observation was performed at magnification 40 (X40). To do this, the preparations of tube 1 (SM); 2; 3 and tube 10 (control tube) were diluted separately to 1/5th. Then each dilution was spread between slide and coverslip by putting the dilution of tube 10 (control) and an experimental dilution (for example tube 1). This last operation was also done for tube 2 and tube 3.

2.2.6 Method of carrying out the experimental tests

For the experimental tests, 2 mg of venom crystals were dissolved in 0.5 mL of physiological solution to obtain a concentration of 4 mg/mL. This is the concentration that was injected into the rabbits. Indeed, according to [25], the median lethal dose for a 2 kg rabbit is 2 mg/Kg of body weight in intra-muscular injection.

3. RESULTS

3.1 Results of the Hemolytic Power of the Venom *In vitro*

Figs .1 (a and b); 2 (a and b) and 3 (a and b) are the results of the effects of different concentrations of the venom tested *in vitro* on red blood cells. These figures showed that in general, the density of red blood cells varied compared to the control depending on the concentrations tested. Thus, it was higher when the concentration was low (0.4 mg/mL) and low when the concentration was high (1.6 mg/mL).




Fig. 1. Hemolysing power of venom on red blood cells (at magnification 40)
1.a: control red blood cells; 1.b: red blood cells at 0.4 mg/mL venom concentration



Fig. 2. Hemolysing power of venom on red blood cells (at magnification 40)
 2.a: control red blood cells; 2.b: red blood cells at 0.8 mg/mL venom concentration



Fig. 3. Hemolysing power of venom on red blood cells (at magnification 40)
 3.a: control red blood cells; 3.b: red blood cells at a concentration of 1.6 mg/mL venom

3.2 Results of Biochemical Parameters of Control Rabbits

The results of the biochemical parameters of the control rabbit batches were recorded in Table 1 below. In this table, the reference value of biochemical parameters of:

- blood glucose was 193.8 ± 24.77 mg/dL for the male and 152.32 ± 17.35 mg/dL for the female;
- creatine levels were 18.38 ± 1.44 mg/L for males and 15.87 ± 1.83 mg/L for females;
- uremia were 45.93 ± 3.55 mg/L for males and 39.66 ± 4.57 mg/L for females;
- GOT value was 87.1 ± 75.99 IU/L for males and 124.57 ± 46.92 IU/L for females;
- GPT value was 81.12 ± 44.09 mg/L for the male and 13.76 ± 15.81 mg/L for the female;
- triglycerides were 344.71 ± 127.18 mg/dL for the male and 154.16 ± 37.64 mg/dL for the female;
- total cholesterol were 94.12 ± 23.08 mg/dL for males and 112.96 ± 16.26 mg/dL for females;
- HDL cholesterol for males was 26.16 ± 33.86 mg/dL and for females 68.07 ± 12.69 mg/dL;
- LDL cholesterol readings were 122.67 ± 184.78 mg/dL for males and 20.87 ± 12.88 mg/dL for females;

- sodium content was 132.87 ± 7.72 mmol/L for males and 141.9 ± 1.75 mmol/L for females;
- chlorine were 107.43 ± 10.22 mmol/L for males and 101.5 ± 0.62 mmol/L for females,
- potassium levels were 3.93 ± 0.13 mmol/L for males and 4 ± 0 mmol/L for females.

3.3 Results of the Biochemical Parameters of the Envenomed Rabbits

In Table 2 are the results of the biochemical parameters of the envenomed rabbit lot. Analysis of these results revealed that:

- in the M1 male, creatinine, uremia, triglycerides and potassium (K) were above their respective reference values, whereas sodium was below its reference value. Apart from these parameters the others were within their respective norms.
- in male M2, blood glucose and triglycerides were respectively superior to their norms, meanwhile transaminases (GOT and GPT), sodium (Na) and potassium (K) were inferior to their respective reference values. Independently of these, the other parameters were normal.
- at the F1 level, blood glucose, creatinine, urea, transaminases, LDLc and ionogram (Na, Cl and K) were elevated compared to their respective norms, in contrast to HDLc. As for triglycerides and total cholesterol, their values were normal.

Table 1. Biochemical parameters of control batches

Lots	Individuals	Gly	Creat.	Urea	Transaminases		TG	CT	HDLc	LDLc	Na	Ionogram Cl	K
					GOT	GPT							
Biochemical parameters of the male control batch													
Lot 1	MW 07	185.2	17.38	43.45	121.41	126.52	198.02	69.33	9.09	20.63	124.5	116.4	3.78
	MW 16	221.72	17.74	44.35	139.88	78.37	411.98	98.04	4.23	11.42	134.4	96.3	4
	MW 10	174.47	20.03	50	0	38.47	424.12	114.98	65.15	335.97	139.7	109.6	4
	Average	193.8	18.38	45.93	87.1	81.12	344.71	94.12	26.16	122.67	132.87	107.43	3.93
	Standard deviation	24.77	1.44	3.55	75.99	44.09	127.18	23.08	33.86	184.78	7.72	10.22	0.13
Reference value		169.03-218.57	16.94-19.82	42.38-49.48	11.11-163.09	37.03-125.21	217.53-471.89	71.04-117.2	-7.7-60.02	-62.11-307.45	125.15-140.59	97.21-117.65	3.8-4.06
Biochemical parameters of the female control batch													
Lot 2	FW 01	134.02	14	35	80.78	0.56	120.62	106.19	64.83	17.24	143.6	100.8	4
	FW 10	154.4	15.94	39.85	118.83	31.29	194.87	131.47	57.32	35.18	142	101.7	4
	FW 12	168.53	17.66	44.14	174.09	9.44	147	101.21	82.07	10.2	140.1	102	4
	Average	152.32	15.87	39.66	124.57	13.76	154.16	112.96	68.07	20.87	141.9	101.5	4
	Standard deviation	17.35	1.83	4.57	46.92	15.81	37.64	16.26	12.69	12.88	1.75	0.62	0
Reference value		134.97-169.67	14.04-17.7	35.09-44.23	77.65-171.49	-2.05-29.57	116.52-191.8	96.7-129.22	55.38-80.76	7.99-33.75	140.15-143.65	100.88-102.12	4

Gly : Glycemia(mg/dL) ; Creat. : Creatinine (mg/L) ; Urea (mg/L) ; Transaminases (GOT and GPT in UI/L) ; TG : Triglycerides(mg/dL) ; TC : Total Cholesterol(mg/dL) ; HDLc : HDL cholesterol (mg/dL) ; LDLc: HDL cholesterol (mg/dL) ; Na : Sodium (mmol/L) ; Cl : Chloride(mmol/L) ; K : Potassium (mmol/L). MW:Male Witness;FW:Female Witness

Table 2. Biochemical parameters of the envenomed batch

Lots		Gly	Creat.	Urea	Transaminases		TG	TC	HDLc	LDLc	Na	Ionogramm Cl	K	Legend
					GOT	GPT								
MW	Ave	193.8	18.38	45.93	87.1	81.12	344.71	94.12	26.16	122.67	132.87	107.43	3.93	Male
	Sd	24.77	1.44	3,55	75.99	44.09	127.18	23.08	33.86	184.78	7.72	10.22	0.13	Lower value
Reference value		169.03-218.57	16.94-19.82	42.38-49.48	11.11-163.09	37.03-125.21	217.53-471.89	71.04-117.2	-7.7-60.02	-62.11-307.45	125.15-140.59	97.21-117.65	3.8-4.06	Higher value
FW	Ave	152.32	15.87	39.66	124.57	13.76	154.16	112.96	68.07	20.87	141.9	101.5	4	Female
	Sd	17.35	1.83	4.57	46.92	15.81	37.64	16.26	12.69	12.88	1.75	0.62	0	Lower value
Reference value		134.97-169.67	14.04-17.7	35.09-44.23	77.65-171.49	-2.05-29.57	116.52-191.8	96.7-129.22	55.38-80.76	7.99-33.75	140.15-143.65	100.88-102.12	4	Higher value
State														
Lot 3	M1	181.22	11.82	29.63	58.02	48.83	120.55	90.81	9.08	58.01	171.1	110.02	3.1	Death
	M2	154.15	19.39	48.51	219.22	172.52	77.73	104.27	20.09	67.73	165.4	106.42	7.54	Death
	F1	172.2	20.64	51.6	381.85	171.08	174.01	116	29.96	49.33	229.3	130.01	9.72	Death

Glyc : Glycemia(mg/dL) ; Creat : creatinine(mg/L) ; Urea (mg/L) ; Transaminases (GOT and GPT en UI/L) ; TG : Triglycerides(mg/dL) ; TC : Total Cholestérol (mg/dL) ; HDLc : HDL cholesterol (mg/dL) ; LDLc : LDL cholesterol (mg/dL) ; Na : Sodium (mmol/L) ; Cl : Chlorine(mmol/L) ; K : Potassium (mmol/L). MW: Male witness;FW: Female witness; Ave:Average;Sd: Standard deviation
Example: M1= Male number 1; F1= Female number 1

3.4 Results of Mineral Tests on Biochemical Parameters of Rabbits Treated as a Preventive Measure before Envenomation

The results of the biochemical parameters of the preventively treated rabbits were reported in Table 3. Analysis of the biochemical parameters in this table in comparison with the reference values in Table 1 revealed that:

- at the level of glycemia, it was normal in a male from lot 4 to lot 10 and in the female from lot 5, lot 9 and lot 10; while it was increased in a male from lot 4 and the female from lot 4, 6, 7 and 8 contrary to a male from lot 5 to 10.
- at the level of creatinine, it was normal in the two males of lot 4 and 7, one male of lot 5 and 8, the female of lot 5, 7 and 9; whereas it was elevated in the female of lot 4, 6 and 8 in contrast to a male of lot 5, 8 and a female of lot 10, and two males of lot 6, 9 and 10.
- with regard to urea, the respect of its norm was observed in two males of lot 4 and 7, one male of lot 5 and 8, and in the female of lot 7, 9 and 10; meanwhile, it was high in the female of lot 4, 6 and 8 in opposition to that of a male of lot 5 and 8, of two males of lot 6, 9 and 10, and of a female of lot 5
- at the GOT level, it was normal in the three individuals of lot 4, two males of lot 6, 7 and 9, one male of lot 8 and 10; at the time it was augmented in two males of lot 5, in one male of lot 8 and 10, and in the female of lot 5, 6, 7, 8, 9 and 10.
- in GPT, it was greater in the female of lots 4, 5, 6, 7, 8, 9 and 10, and in one male of lots 5 and 8. Apart from these observations, it was normal in the individuals of the other batches.
- triglycerides were lower in one male from lots 4, 5, 6, 9 and 10, and in two males from lots 7 and 8, and vice versa in the female from lots 4, 5, 6, 7 and 9. Except for these findings, they met their standard in the other cases.
- total cholesterol was high in the females of lots 4, 6, 7 and 9, and in one male of lots 5 and 10, although the opposite was true in one male of lots 6 and 7. Excluding these remarks, it was normal in the animals of the various constituted batches.
- the HDLc level was decreased in the female of lots 4, 5, 6, 7, 8, 9 and 10 in contrast to a male of lot 5. In addition to

these observations, the respect of its norm was noticed in the animals of the different batches made.

- LDLc was elevated in one male from batch 4 and in the female from batches 4, 5, 6, 7, 8, 9 and 10, but was normal in the other individuals from the various batches.
- in terms of sodium, it was increased in the female of batches 7 to 10, and two males of batches 7 to 10; however, it was low in a female of batches 4 to 6, and a male of batch 5. Except for these remarks, it was normal in the other animals.
- at the level of chlorine, it was reduced in the female of lot 7; whilst it was increased in the female of lots 4, 5, 6, 8, 9 and 10, and a male of lot 7. Besides these remarks, it obeyed its norm in the other individuals of the different batches realized.
- in terms of potassium, it was normal in the two males of lot 4, and one male of lots 5, 9 and 10. It was lower in the female of lots 4 to 7, one male of lots 5, 7 and 8, and the two males of lot 6, in contrast to one male of lots 7 to 10, and in the female of lots 8 to 10.

3.5 Results of Mineral Tests on Biochemical Parameters of Rabbits Treated as a Preventive Measure and then Envenomed

The biochemical parameters of the rabbits treated as a preventive measure and then envenomed in Table 4 made it possible to note in comparison with lot 3 in Table 2 (untreated envenomed lot) that :

- in general, the batches are characterized by the death of all animals in batches 4, 7 and 8, and the survival of one male in batch 10, one male and one female in batch 5, two males in batch 9, and all individuals in batch 6.
- there was considerable variation in blood glucose, creatinine, urea and total cholesterol in all animals of the different batches made. However, exceptions were observed for transaminases, triglycerides, HDLc, LDLc and ionograms in the different batches. Thus at the level of:
 - GOT, apart from the animals of lot 4 that obeyed the results of Table 2, the GOT values obtained in the other lots differed from those of Table 2;
 - GPT, besides the animals of lot 4 and in a certain sense those of lot 9 and 10, the GPT

values got in the other cases were different from those of Table 2;

- triglycerides, except for the animals of lot 8 which respected the observations of Table 2, the females of lots 6, 7, 9 and 10 allowed to have variable triglyceride values contrary to the different males of these lots. As for the remaining batches, their triglyceride values were variable;
- HDLc, excluding lots 7, 8 and 10 and to some extent lot 5 which followed the remarks in Table 2, the HDLc values achieved in the other cases varied;
- LDLc, the results were exactly the same as in Table 2, with one exception, a male from lot 4;
- ionogram, with the certain exception of the animals of lot 4 and those of lot 10 which respected the results of Table 2, the animals of the other lots allowed to have discordant results.

When the biochemical parameters of the preventively treated and then envenomed rabbits in Table 4 were compared to those in Table 3 (rabbits scarified preventively prior to envenomation), it was noted that in the:

- lot 4; outside of triglycerides and chlorine, which were the same in both tables, the other parameters (blood glucose, creatinine, urea, GOT, GPT, total cholesterol, HDLc, LDLc, sodium and potassium) varied either in one animal or in two or even in three for this lot;
- in lot 5; except for GOT and LDLc that were identical in both tables, the other parameters differed for the same lot from one animal of the same sex to another;
- from lot 6; all biochemical parameters studied for a given animal of a constituted lot changed in both tables;
- from lot 7; with the exception of triglycerides, HDLc and sodium which were the same in both tables, the other biochemical parameters studied varied in one or two or three animals for that lot from one table to another;
- of lot 8; other than the triglycerides, HDLc, LDLc and sodium which did not change between the two tables, the other biochemical parameters studied were modified in one or two or three animals for this lot from one table to another;

- from lot 9; apart from LDLc and sodium which were the same in both tables, the other biochemical parameters studied were not in one or two or three animals for this lot from one table to another;
- of lot 10; independently of HDLc, LDLc and ionogram (sodium, chlorine and potassium) which were the same in both tables, the other biochemical parameters studied were not identical in one animal or two or three for this lot from one table to another.

3.6 Results of Mineral Tests on Biochemical Parameters of Rabbits Treated Curatively after Envenomation

The biochemical parameters of the envenomed and then curatively treated rabbits in Table 5 in comparison with lot 3 in Table 2 (untreated envenomed lot) showed that:

- in lot 11, the increase of GPT and ionogram (sodium, chlorine and potassium) was noticed in two females of the lot as well as those of lot 3 (envenomed not treated). Similarly, it was found that the two individuals in Table 5 did not survive like the one in Table 2. The elevation of GPT and ionogram were followed by a regression of total cholesterol in both individuals. Apart from these common parameters, the others varied. As for the surviving female, these biochemical parameters such as blood glucose, creatinine, transaminases (GPT and GOT), LDLc and sodium were rising like those of batch 3 (untreated envenomed). Except for these parameters, the others changed.
- in batch 12, elevated creatinine, urea, GPT, LDLc, sodium and chlorine in one female were followed by a decrease in HDLc. These observed parameters were identical to those of batch 3 (untreated envenomed). It was also observed that one of the two females in Table 5 did not survive like the one in Table 2. In contrast to this finding, the second female was found alive. Regardless of these parameters, the others changed. In the second female of the batch that survived, there was an increase in GPT, sodium and potassium as well as those of batch 3 (envenomed untreated). In addition to these parameters, the others varied.

Table 3. Biochemical parameters of rabbits treated with minerals as a preventive measure

Lots	Individuals	Gly	Creat.	Urea	Transaminases		TG	TC	HDLc	LDLc	Ionogram		
					GOT	GPT					Na	Cl	K
Lot 4 (P1)	M.3	180.09	17.6	44	40.59	66.17	177.34	104.59	42.25	26.88	137.6	112.8	3.8
	M.13	222.91	17.66	44.15	99.37	99.86	312.9	96.05	33.93	259.76	132.9	115.3	3.84
	F.3	192.27	19.14	47.85	133.67	181.21	335.66	137.4	20.85	49.42	132.9	109.1	3.95
Lot 5 (P2)	M.14	125.11	18.57	46.42	370.57	106.26	350.59	245.9	141.86	33.93	123.7	107.4	3.81
	M.9	187.08	7.1	17.75	253.07	212.66	77.96	76.72	40.15	21.03	132.6	112.1	3.66
	F.11	157.81	15.26	34.66	201.12	179.57	202.69	122.34	15.32	62.9	135.9	115.6	3.81
Lot 6 (P3)	M.8	179.3	16.24	40.6	103.78	112.65	349.48	102.06	11.9	317.17	127.8	114.7	3.78
	M.11	119.6	14.19	35.47	111.43	50.37	147.56	62.86	13.05	121.94	125.7	117.1	3.41
	F.7	190.88	23.84	59.6	231.24	192.35	281.81	138.31	23.96	230.19	129.4	117.6	3.65
Lot 7 (P4)	M.1	131.61	17.01	42.52	69.39	84.95	146.13	64.88	7.23	28.22	183.3	127.5	6.93
	M.6	206.25	19.03	47.57	71.55	66.34	116.55	116.09	13.06	79.72	183.6	110	2.91
	F.6	188.94	15.8	39.5	175.82	109.94	319.96	163.16	14.94	84.23	165.3	98.7	3.36
Lot 8 (P5)	M.5	181.2	11.82	29.53	57.97	48.83	120.15	90.81	8.88	57.9	171.1	110.6	3.1
	M.12	154.25	19.39	48.47	219.32	172.52	75.73	104.27	25.89	67.23	165.4	105.2	7.54
	F.14	172.19	20.64	51.6	381.85	171.08	173.81	116	30.91	50.33	229.3	129.6	9.83
Lot 9 (P6)	M.15	141.39	14.53	36.32	71.68	76.64	234.56	98.29	13.39	37.99	188.3	111.5	4.03
	M.2	186.8	16.83	42.07	129.51	122.88	211.36	93.53	13.33	37.93	171.8	112.9	6.27
	F.13	142.83	14.83	37.05	604.59	245.89	287.09	155.16	24.08	76.66	181.9	118.1	5.66
Lot 10 (P7)	M.7	170.88	16.68	41.7	104.16	79.86	158.49	90.82	13.15	79.86	160.37	113.33	4.2
	M.4	132.7	16.7	41.75	170.54	89.28	243.76	136.36	54.16	33.38	165.1	115.47	4.92
	F.2	158.78	16.34	40.85	342.09	484.52	133.92	118.42	16.73	74.91	168.4	114.1	5.84
Reference value	M	169.03-218.57	16.94-19.82	42.38-49.48	11.11-163.09	37.03-125.21	217.53-471.89	71.04-117.2	-7.7-60.02	-62.11-307.45	125.15-140.59	97.21-117.65	3.8-4.06
	F	134.97-169.67	14.04-17.7	35.09-44.23	77.65-171.49	-2.05-29.57	116.52-191.8	96.7-129.22	55.38-80.76	7.99-33.75	140.15-143.65	100.88-102.12	4

Gly: Glycemia (mg/dL); Creat.: Creatinine (mg/L); Urea (mg/L); Transaminases (GOT and GPT in U/L); TG: Triglycerides (mg/dL); TC: Total Cholesterol (mg/dL); HDLc: HDL cholesterol (mg/dL); LDLc: LDL cholesterol (mg/dL); Na: Sodium (mmol/L); Cl: Chloride (mmol/L); K: Potassium (mmol/L). M: Male; F: Female
 Example: M7= Male number 7; F2= Female number 2

Table 4. Biochemical parameters of rabbits treated as a preventive measure and then envenomed

Lots	Individuals	Gly	Creat.	Urea	Transaminases		TG	TC	HDLc	LDLc	Ionogram			State
					GOT	GPT					Na	Cl	K	
Lot 4 (P1)	M.3	339.32	33.87	13.55	83.69	75.12	151.98	54.08	61.23	166.55	156.5	107.8	8.22	Death
	M.13	350.66	44.6	17.84	203.54	143.31	273.09	55.14	132.6	360.67	144.9	97.4	1.27	Death
	F.3	308.17	48.75	19.5	320.61	327.61	428.6	99.51	98.25	267.24	197	118.4	8.57	Death
Lot 5 (P2)	M.14	39.47	18.7	46.75	675.38	0	237.11	209.47	35.51	96.59	154.8	100.5	4.31	Death
	M.9	147.35	17.04	42.6	295.44	271.03	112.04	28.85	9.05	24.62	163.1	104.2	3.8	Survivor
	F.11	82.38	16.95	42.37	225.4	0	86.36	99.06	80.09	217.84	142.3	98.7	1.81	Survivor
Lot 6 (P3)	M.8	115.06	13.76	34.4	426.76	0	105.8	61.85	71.92	195.62	164.3	99.3	4.72	Survivor
	M.11	324.68	39.15	15.66	90.16	106.74	96.66	44.86	60.79	165.35	147.5	102	5.69	Survivor
	F.7	78.19	17.18	42.95	121.08	563.75	116.16	116.53	23.94	65.12	153.4	101.2	5.16	Survivor
Lot 7 (P4)	M.1	85.53	16.57	41.42	146.05	251.56	59.11	62.03	51.64	140.46	145.2	99.1	2.26	Death
	M.6	341.19	38.52	15.41	175.89	134.53	140.2	47.55	59.56	162	146.1	101.8	4.86	Death
	F.6	82.1	17.86	44.65	0	0	264.21	100.19	15.04	40.91	150.3	98.9	4.71	Death
Lot 8 (P5)	M.5	93.48	16.49	41.22	709.54	0	45.24	58.87	48.63	132.27	154.9	105.2	2.16	Death
	M.12	148.47	17.76	44.4	227.57	321.61	84.41	53.91	12.01	32.67	150.3	107.9	4.1	Death
	F.14	117.64	12.61	31.52	369.78	0	174.93	47.24	15.78	42.92	149.7	101.2	2.7	Death
Lot 9 (P6)	M.15	106.07	16.97	42.42	175.2	0	78.81	76.29	64.96	176.69	153.2	102.1	4.53	Survivor
	M.2	85.13	23.25	58.12	0	632.38	60.41	80.66	64.42	175.22	142.4	100.6	3.38	Survivor
	F.13	350.66	44.6	17.84	203.54	143.31	273.09	55.14	132.6	360.67	144.9	97.4	1.27	Death
Lot 10 (P7)	M.7	51.29	53.43	133.87	0.2	445.12	124.06	24.94	23.41	63.67	147.3	102.6	3.91	Survivor
	M.4	84.36	19.12	47.8	245.5	219.36	55.19	46.84	34.81	94.68	153.6	102.8	4.08	Death
	F.2	103.42	13.75	34.37	84.84	524.72	84.76	40.25	16.56	45.04	148.3	103.8	4.41	Death

Gly : Glycemia (mg/dL) ; Creat. : Creatinine (mg/L) ; Urea (mg/L) ; Transaminases (GOT and GPT in UI/L) ; TG : Triglycerides (mg/dL) ; TC : Total Cholesterol (mg/dL) ; HDLc : HDL cholesterol (mg/dL) ; LDLc:cholesterol (mg/dL) ; Na : Sodium (mmol/L) ; Cl : Chloride (mmol/L) ; K : Potassium (mmol/L). M:Male;F: Female
Example: M1= Male number 1; F14= Female number 14

Table 5. Biochemical parameters of envenomed and curatively treated rabbits

Lots	Individuals	Gly.	Creat.	Urea	Transaminases		TG	CT	HDLc	LDLc	Ionograms			State
					GOT	GPT					Na	Cl	K	
Lot 11 (P1)	F.9	152.94	13.1	32.53	159.15	155.83	126.06	52.96	14.31	13.44	166.4	104.8	8.11	Death
	F.5	322.64	24.8	9.92	278.13	196.18	229.63	93.59	101.53	53.87	152.4	96.9	2.38	Survivor
	F.8	87.1	19.15	47.87	150.02	130.5	103.7	85.29	76.11	11.56	153.5	106.6	4.49	Death
Lot 12 (P3)	F.15	97.34	8.83	22.87	100.95	440.48	215.59	95.43	15.99	36.32	185.2	122.7	6.65	Survivor
	F.16	58.19	20.1	49.8	100	73	96.91	77.89	23.8	34.71	154.9	106.1	3.4	Death
	F.17	99.93	22.4	55.7	168	209	233.88	137.97	59.8	11.4	186.4	101.2	4.1	Survivor
Lot 3 envenomed	F1	172.2	20.64	51.6	381.85	171.08	174.01	116	29.96	49.33	229.3	130.01	9.72	Death

Gly : Glycemia (mg/dL) ; Creat. : Creatinine (mg/L) ; Urea (mg/L) ; Transaminases (GOT and GPT in UI/L) ; TG : Triglycerides (mg/dL) ; TC : Total Cholesterol (mg/dL) ; HDLc : HDL cholesterol (mg/dL) ; LDLc:cholesterol (mg/dL) ; Na : Sodium (mmol/L) ; Cl : Chloride (mmol/L) ; K : Potassium (mmol/L).F: Female
Example: F17= Female number 17

4. DISCUSSION

4.1 Results of the Hemolytic Power of the Venom *In vitro*

"The results of the *in vitro* hemolysis test showed that in general the venom of *Naja nigricollis* has a hemolytic effect on rabbit whole blood. However, these effects are dependent on its concentration. In fact, hemolysis is more important when the concentration is high. This explains why the density of red blood cells is low with the concentration of 1.6 mg/mL; whereas it is higher with the lowest concentration used (0.4 mg/mL). This gradual action of venom would imply that this effect would be a function of the quantity injected. Thus, venom has a dose-dependent action on red blood cells *in vitro*. These results are in agreement with those of" [26]. This author indicates that the effect of Elapidae venom is proportional to the quantity of toxin molecules introduced into the organism.

4.2 Results of Biochemical Parameters of Control Rabbits

Comparison of the result of the reference values in Table 1 showed that with the exception of GOT, total cholesterol, HDLc and sodium, the biochemical parameters of males were generally higher than those of females. Besides these exceptions, the biochemical parameters obtained varied from male to female animals. This observation was confirmed by [27] who stated that "the variation in blood content of these different biochemical elements is a function of sex, reproductive cycle, age and growth of rabbits". "Nevertheless, all the biochemical parameters obtained from the control batches during this study were with few exceptions identical to those of" [28].

4.3 Results of the Biochemical Parameters of the Envenomed Rabbits

At M1 and F1, there is an elevation in both creatinine and urea values. According to [29,30,31] and [32], an increase in serum urea concentration associated with creatinine would indicate renal failure. Therefore, it can be inferred that the M1 and F1 males would obey this assertion. There was also an increment in triglycerides in M1 and M2. According to [33,34] and [32], "this rising value of triglycerides would be consistent with hypertriglyceridemia and

would be a risk factor for predicting cardiovascular disease. The reason is that triglycerides are the main constituents of cell membranes and the increase in their concentration would probably be the result of apoptosis". Accordingly, M1 and M2 would obey this principle. LDLc is also elevated in F1. This increase could promote the risk of coronary mortality because according to [35,36] and [32], its reduction would induce the opposite effect. Regarding the elevation of transaminases in F1, it could indicate hepatocellular damage or bile flow disturbance according to [37] and [32]. The high value of ionogram (Na, Cl and K) in F1 and blood glucose in F1 and M2 would mean a heart problem and diabetes respectively according to [38]. In M2, the decrease in transaminases would reveal a vitamin B6 deficiency according to [39]. Concerning dyskalemia (potassium variation) in M1 and M2, they would be the basis of neuromuscular and cardiac signs according to [40] or even hemodynamic according to [41]. These authors believe that potassium plays a major role in maintaining the resting membrane potential. These authors generally stipulate that dyskalemia (hyperkalemia or hypokalemia) is life-threatening. HDLc decay in F1 would designate according to [42,37] and [32], "hypocholesterolemia. It can be observed in liver damage, severe infections, anemia, treatment with particular drugs (hormones such as clofibrate and androsterone), mental retardation and congenital acyl transferase deficiency. Thus, these different pathologies mentioned above would be related to the action of the venom on the vital organs of the various animals involved, which would lead to their death".

4.4 Results of Mineral tests on Biochemical Parameters of Rabbits Treated as a Preventive Measure before Envenomation

"The increase in blood glucose in M13 and F3 of lot 4, F7 of lot 6, F6 of lot 7 and F14 of lot 8 treated respectively with P1 (ash of both plants and viper skull), P3 (ash of *Pongamia pinnata*), P4 (ash of viper skull) and P5 (ash of *Mucuna pruriens* and *Pongamia pinnata*) would indicate, according to [38] and [32], hyperglycemia". These authors suggest that hyperglycemia is due to insulin-dependent or non-insulin-dependent diabetes, pancreatic diseases: acute or chronic pancreatitis, endocrine diseases: pheochromocytoma, hypercorticism, corticosteroid therapy, and hypothyroidism. The decrease in M14 of lot 5,

M11 of lot 6, M1 of lot 7, M12 of lot 8, M15 of lot 9 and M4 of lot 10 treated respectively with P2 (*Mucuna pruriens* ash), P3 (*Pongamia pinnata* ash), P4 (viper skull ash), P5 (ash of *Mucuna pruriens* and *Pongamia pinnata*), P6 (ash of viper skull and *Mucuna pruriens*) and P7 (ash of viper skull and *Pongamia pinnata*) would reflect hypoglycemia according to the same authors. It would be due to malnutrition or prolonged fasting, secretion by the body of an excess of insulin: insulinoma, polyadenomatosis, endocrine insufficiency: adrenal, pituitary, and a hepatic disorder: acute hepatitis. Also, according to [43] and [32]; "blood glucose is also low in anorexia and high in diabetes. Hence, it can be said that anorexia would be observed in M14 of batch 5, M11 of batch 6, M1 of batch 7, M12 of batch 8, M15 of batch 9 and M4 of batch 10; whereas diabetes would be seen in M13 and F3 of batch 4, F7 of batch 6, F6 of batch 7 and F14 of batch 8".

"The elevation of creatinine in F3 of batch 4, F7 of batch 6, F6 batch 7 and F14 of batch 8 treated with P1 (ash of both plants and viper skull), P3 (ash of *Pongamia pinnata*) respectively, P4 (ash of viper skull) and P5 (ash of *Mucuna pruriens* and *Pongamia pinnata*) could be explained by the reduction of creatinuria i.e. the elimination of creatinine by the kidneys because this excretion by the kidneys is more specific to creatinine than urea" [31,32]. "This decrease in creatinuria would thus show a dysfunction of the kidney. Regarding the decline in creatinine in M9 of batch 5, M8 and M11 of batch 6, M5 of batch 8, M15 and M2 of batch 9, and M7, M4 and F2 of batch 10 treated with P2 (*Mucuna pruriens* ash), P3 (*Pongamia pinnata* ash) respectively, P5 (*Mucuna pruriens* and *Pongamia pinnata* ash), P6 (viper skull and *Mucuna pruriens* ash) and P7 (viper skull and *Pongamia pinnata* ash), it could be a sign of cachexia" [44,45,31,32].

The higher urea in F3 of batch 4, F7 of batch 6 and F14 of batch 8 treated with P1 (ash of both plants and viper skull), P3 (ash of *Pongamia pinnata*) and P5 (ash of *Mucuna pruriens* and *Pongamia pinnata*), respectively, would indirectly reveal renal and liver function dysfunction [46,37,32]. "Relatively to its low in M9 and F11 of lot 5, M8 and M11 of lot 6, M5 of lot 8, M15 and M2 of lot 9, M7 and M4 of lot 10 treated with P2 (*Mucuna pruriens* ash), P3 (*Pongamia pinnata* ash) respectively, P5 (ash of *Mucuna pruriens* and *Pongamia pinnata*), P6 (ash of viper skull and *Mucuna pruriens*) and P7 (ash of viper skull and *Pongamia pinnata*) would show hypo-

uremia. This hypo-uremia would reflect severe liver failure or alternatively a deficiency in urea cycle enzyme" [37,32].

According to [29,30,31,32], the rise in serum urea concentration associated with that of creatinine would indicate renal failure. As such, it is possible to deduce that F3 of batch 4, F7 of batch 6 and F14 of batch 8 treated with P1 (ash of both plants and viper skull), P3 (ash of *Pongamia pinnata*) and P5 (ash of *Mucuna pruriens* and *Pongamia pinnata*) respectively would obey this assertion.

"The enhancement of GOT in M14, M9 and F11 from lot 5, F7 from lot 6, F6 from lot 7, M12 and F14 from lot 8, F13 from lot 9 and M4 and F2 from lot 10 treated P2 (*Mucuna pruriens* ash), P3 (*Pongamia pinnata* ash), P4 (viper skull ash), P5 (*Mucuna pruriens* and *Pongamia pinnata* ash) respectively, P6 (ashes of viper skull and *Mucuna pruriens*) and P7 (ashes of viper skull and *Pongamia pinnata*) would show the destruction of hepatocytes although in addition to the liver, this enzyme can be found in the heart, skeletal muscle, lungs and kidneys" [47,48,32].

"The increase in GPT in F3 of lot 4, M9 and F11 of lot 5, F7 of lot 6, F6 of lot 7, M12 and F14 of lot 8, F13 of lot 9 and F2 of lot 10 treated with P1 (ash of both plants and viper skull), P2 (ash of *Mucuna pruriens*), P3 (ash of *Pongamia pinnata*), P4 (ash of viper skull), respectively, P5 (ash of *Mucuna pruriens* and *Pongamia pinnata*), P6 (ash of viper skull and *Mucuna pruriens*) and P7 (ash of viper skull and *Pongamia pinnata*) would reveal hepatic cell necrosis because GPT is a cytosolic enzyme secreted in liver cells. It is released into the bloodstream in case of hepatic cell necrosis" [49,47, 48,32].

According to [37] and [32], "elevated transaminases (GOT and GPT) could reflect hepatocellular damage or bile flow disturbance. Therefore, elevated transaminases (GOT and GPT) would be observed in acute hepatitis. Hence, M9 and F11 of batch 5, F7 of batch 6, F6 of batch 7, M12 and F14 of batch 8, F13 of batch 9 and F2 of batch 10 treated respectively with P2 (*Mucuna pruriens* ash), P3 (*Pongamia pinnata* ash), P4 (viper skull ash), P5 (ash of *Mucuna pruriens* and *Pongamia pinnata*), P6 (ash of viper skull and *Mucuna pruriens*) and P7 (ash of viper skull and *Pongamia pinnata*) would obey this observation".

Decrease in triglycerides was observed in M3 of batch 4, M9 of batch 5, M11 of batch 6, M1 and M6 of batch 7, M5 and M12 of batch 8, M2 of batch 9 and M7 of batch 10 treated with P1 (ash of both plants and viper skull) respectively, P2 (ash of *Mucuna pruriens*), P3 (ash of *Pongamia pinnata*), P4 (ash of viper skull), P5 (ash of *Mucuna pruriens* and *Pongamia pinnata*), P6 (viper skull ash and *Mucuna pruriens*) and P7 (viper skull ash and *Pongamia pinnata*) in opposition to their elevation in F3 of lot 4, F11 of lot 5, F7 of lot 6, F6 of lot 7 and F13 of lot 9 treated with P1 (ash of both plants and viper skull), P2 (ash of *Mucuna pruriens*), P3 (ash of *Pongamia pinnata*), P4 (ash of viper skull) and P6 (ash of viper skull and *Mucuna pruriens*) respectively. This increase in triglycerides is consistent with hypertriglyceridemia in females from the above batches. According to [33,34] and [32], this hypertriglyceridemia would be a risk factor for predicting cardiovascular disease. In effect, triglycerides are the main constituents of cell membranes and the elevation of their concentration would likely be the result of apoptosis. So, F3 from batch 4, F11 from batch 5, F7 from batch 6, F6 from batch 7 and F13 from batch 9 would be exposed to cardiovascular diseases in contrast to M3 from batch 4, M9 from batch 5, M11 from batch 6, M1 and M6 from batch 7, M5 and M12 from batch 8, M2 from batch 9 and M7 from batch 10 treated respectively with P1 (ash from both plants and from viper skull), P2 (ash of *Mucuna pruriens*), P3 (ash of *Pongamia pinnata*), P4 (ash of viper skull), P5 (ash of *Mucunapruriens* and *Pongamia pinnata*), P6 (ash of viper skull and *Mucuna pruriens*) and P7 (ash of viper skull and *Pongamia pinnata*) respectively.

The increase in total cholesterol was noticed in F3 of batch 4, M14 of batch 5, F7 of batch 6, F6 of batch 7, F13 of batch 9 and M4 of batch 10 treated with P1 (ash of both plants and viper skull), P2 (ash of *Mucuna pruriens*), P3 (ash of *Pongamia pinnata*) respectively, P4 (viper skull ash), P6 (viper skull and *Mucuna pruriens* ash) and P7 (viper skull and *Millettia pinnata* ash), in contrary to their regression in M11 of lot 6 and M1 of lot 7 treated with P3 (*Pongamia pinnata* ash) and P4 (viper skull ash), respectively. The reduction designates according to [42,37] and [32], "hypocholesterolemia. It can be observed in liver damage, severe infections, anemia, treatment with particular drugs (hormones such as clofibrate and androsterone), mental retardation and congenital acyl transferase deficiency. As for their elevation, according to the

authors, it indicates hypercholesterolemia. It is observed in hepatic disorders, renal disorders, pancreatic disorders and thyroid disorders". According to [36] and [32], "hypercholesterolemia is characterized by chronic inflammation of the arterial wall that develops in response to damage affecting the vascular endothelium. Atherosclerotic lesions develop in large and medium caliber arteries, which are approximately 3 to 0.5 cm in diameter".

The lowering of HDLc found in F3 of batch 4, F11 of batch 5, F7 of batch 6, F6 of batch 7, F14 of batch 8, F13 of batch 9, and F2 of batch 10 treated with P1 (ash of both plants and viper skull), P2 (ash of *Mucuna pruriens*), P3 (ash of *Pongamia pinnata*), respectively, P4 (viper skull ash), P5 (*Mucuna pruriens* and *Pongamia pinnata* ash), P6 (viper skull and *Mucuna pruriens* ash), and P7 (viper skull and *Pongamia pinnata* ash) was in antagonism to M14 from lot 5 treated with P2 (*Mucuna pruriens* ash). According to [49,50] and [31], "a 1% increase in HDLc is associated with a 3 to 4% decrease in coronary risk, and a level below 0.40 g/L in humans is considered a risk factor for cardiovascular disease". Similarly, the work of [51] and [32] showed "that the increase in HDLc levels would indicate a protective factor for the heart muscle, thus revealing its beneficial effect against cardiovascular complications, particularly atherosclerosis. Therefore, M14 of lot 5 treated with P2 (*Mucuna pruriens* ash) would be more protected against atherosclerosis than F3 of lot 4, F11 of lot 5, F7 of lot 6, F6 of lot 7, F14 of lot 8, F13 of lot 9 and F2 of lot 10".

The elevation of LDLc was seen in M8 of lot 6 treated with P3 (*Pongamia pinnata* ash) as well as in F3 of lot 4, F11 of lot 5, F7 of lot 6, F6 of lot 7, F14 of lot 8, F13 of lot 9, and F2 of lot 10 treated with P1 (ash of both plants and viper skull), P2 (*Mucuna pruriens* ash), P3 (*Pongamia pinnata* ash), P4 (viper skull ash), P5 (*Mucuna pruriens* and *Pongamia pinnata* ash), P6 (viper skull and *Mucuna pruriens* ash), and P7 (viper skull and *Pongamia pinnata* ash) respectively. [35,36], and [32], suggested that a 0.38 g/L reduction in LDLc would be associated with a one-third decrease in coronary mortality. Hence, M8 in batch 6, F3 in batch 4, F11 in batch 5, F7 in batch 6, F6 in batch 7, F14 in batch 8, F13 in batch 9, and F2 in batch 10 would be more prone to coronary mortality because of their increased.

The increase in sodium observed in M1, M6, and F6 of lot 7, M5, M12, and F14 of lot 8, M15, M2,

and F13 of lot 9, and M7, M4, and F2 of lot 10 treated with P4 (viper skull ash), P5 (*Mucuna pruriens* and *Pongamia pinnata* ash), respectively, P6 (viper skull ash and *Mucuna pruriens*) and P7 (viper skull ash and *Pongamia pinnata*) was in contradiction to F3 of lot 4, F11 and M14 of lot 5 and F7 of lot 6 treated with P1 (ash of both plants and viper skull), P2 (*Mucuna pruriens* ash) and P3 (*Pongamia pinnata* ash) respectively. This elevation and regression of sodium would determine according to [38] “respectively a hypernatremia and a hyponatremia. According to the same author, hypernatremia can be the consequence of a decline in the quantity of water, diarrhea, vomiting, significant sweating, significant water loss (diabetes insipidus, diabetes mellitus), intense exercise. So, M1 and M6 and F6 of lot 7, M5, M12 and F14 of lot 8, M15, M2 and F13 of lot 9, and M7, M4 and F2 of lot 10 would conform to this assertion. As for hyponatremia, it can be the consequence of a decrease in the amount of salt during digestive losses (vomiting, diarrhea), renal losses, skin losses (extensive burning, sweating), an increase in the amount of water during inappropriate secretions of anti-diuretic hormones, excessive water intake, renal failure, cardiac failure, hepatic failure. Therefore, F3 of batch 4, F11 and M14 of batch 5 and F7 of batch 6 would comply with these remarks”.

The lowering of chlorine was detected in F6 of lot 7 cared for by P4 (viper skull ash) in contrast to F3 of lot 4, F11 of lot 5, F7 of lot 6, M1 of lot 7, F14 of lot 8, F13 of lot 9 and F2 of lot 10 processed by P1 (ash of both plants and viper skull), P2 (ash of *Mucuna pruriens*) respectively, P3 (*Pongamia pinnata* ash), P4 (viper skull ash), P5 (*Mucuna pruriens* and *Pongamia pinnata* ash), P6 (viper skull and *Mucuna pruriens* ash) and P7 (viper skull and *Pongamia pinnata* ash) would, according to [38], announce hypochloremia and hyperchloremia respectively. Hypochloremia can be the consequence of a depletion of salt due to digestive losses (vomiting, diarrhea), renal losses, skin losses (extensive burns, sweating), an increment of water during inappropriate secretions of antidiuretic hormones, excessive water intake, renal failure, cardiac failure, liver failure. Whereas hyperchloremia can be the consequence of decreased water intake, diarrhea, vomiting, heavy sweating, heavy water loss (diabetes insipidus, diabetes mellitus), intense exercise. Thus F6 of batch 7 would approve the case of hypochloremia while the others would approve hyperchloremia.

Potassium depreciation noticed in F3 of lot 4, M9 and F11 of lot 5, M8, M11 and F7 of lot 6, M6 and F6 of lot 7, and M5 of lot 8 tended by P1 (ash of both plants and viper skull) respectively, P2 (ash of *Mucuna pruriens*), P3 (ash of *Pongamia pinnata*), P4 (ash of viper skull), and P5 (ash of *Mucuna pruriens* and *Pongamia pinnata*) was the reverse in M1 from lot 7, M12 and F14 from lot 8, M2 and F13 from lot 9, M4 and F2 from lot 10 processed with P4 (viper skull ash), P5 (*Mucuna pruriens* and *Pongamia pinnata* ash), P6 (viper skull and *Mucuna pruriens* ash), and P7 (viper skull and *Pongamia pinnata* ash) respectively. These variations would report hypokalemia and hyperkalemia respectively according to [38]. The same author reveals that dyskalemia (abnormalities: hypokalemia or hyperkalemia) are frequently the cause of cardiovascular disorders. However, hypokalemia can be encountered during digestive losses (vomiting, diarrhea), in hyperaldosteronism (corticoid treatment), renal tubular acidosis, hyperglycemia, metabolic alkalosis. Meanwhile, hyperkalemia can be in exogenous potassium intake, adrenal insufficiency (or antialdosterone treatment), hemolysis, crush syndrome, chemotherapy, intense exercise, chronic renal failure. Therefore, F3 of batch 4, M9 and F11 of batch 5, M8, M11 and F7 of batch 6, M6 and F6 of batch 7, and M5 of batch 8 would show hypokalemia, while M1 of batch 7, M12 and F14 of batch 8, M2 and F13 of batch 9, M4 and F2 of batch 10 would show hyperkalemia.

4.5 Results of Mineral Tests on Biochemical Parameters of Preventively Treated and Envenomed Rabbits

The death of all the animals of lot 4; 7 and 8, and the survival of a male and a female of lot 5, of all the animals of lot 6, of two males of lot 9 and of a male of lot 10 treated respectively by P1, P4 and P5 (for those who died) and, P2, P3, P6 and P7 (for those who survived). These results would indicate that in the first case P1, P4 and P5 could not have had a preventive action on the various animals constituted against the action of the venom. This state of affairs was at the origin of the death of all these animals. In the second case, P2, P3, P6 and P7 were able to prevent the life of one, two or even three animals, depending on the batch considered against the action of the venom.

If the comparison of the biochemical parameters of the batches treated as a preventive measure

and then envenomated with those of the untreated envenomated batch allowed considerable variability in their parameters, this would prove that the venom caused clinical and blood disorders. These clinical and blood disorders obtained during this study were confirmed by the work of [25] and [52]. Thus, according to the work of [12,13], and [14], “the toxins from the venom can intervene alone or in synergy with other components of the venom to cause serious disturbances of the biological systems of the organism. These disturbances are the cause of edema, hematological, neurological, renal, pulmonary and tissue necrosis characteristic of ophidian envenomation”.

4.6 Results of Mineral Tests on Biochemical Parameters of Rabbits Treated Curatively after Envenomation

The mortality of two females from batch 11 treated with P1 (i.e., approximately 66.67%) and one female from batch 12 treated with P3 (i.e., approximately 33.33%) would mean that these various potions could not have had a complete curative action against the venom on these respective animals. However, the survival of one female from lot 11 cured by P1 (i.e., about 33.33%) and two females treated by P3 (i.e., about 66.67%) would show that these various potions could have had a partial curative effect against the venom on these animals. Consequently, it is possible to say that P3 is more curatively active than P1 on the animals used.

If the comparison of the biochemical parameters of the envenomed and then curatively treated batches to those of the untreated envenomed batch allowed variable biochemical parameters from one batch to another, this would prove that the venom induced clinical and blood disorders. These are corroborated by the work of [25] and [52]. Similarly, the work of [12,13] and [14] have shown that “toxins from the venom can act alone or in synergy with other components of the venom to cause serious disruption of the biological systems of the body. These disturbances are the cause of edema, hematological, neurological, renal, pulmonary and tissue necrosis characteristic of ophidian envenomation”.

5. CONCLUSION

The general objective of this study was to determine the antivenom effects of *Mucuna*

pruriens and *Pongamia pinnata* minerals on some biochemical parameters in rabbits. These two plants are traditionally considered to have antivenom therapeutic properties. Thus, this study corroborates the objective of our course which is to develop the medicinal plants of our heritage. The results of this study showed that the venom has a dose-dependent action on red blood cells *in vitro*. As for the *in vivo* tests, they demonstrated that in general, the minerals from the potion P2 (*Mucuna pruriens*), P3 (*Pongamia pinnata*), P4, P5 and P6 stabilize certain biochemical parameters on a preventive basis. As a curative measure, P1 and P3 (*Pongamia pinnata*) more or less preserve certain parameters because some animals were able to survive in both cases of treatment against the action of the venom, with a greater activity for P3. Therefore, for the development of an antivenomous phytomedicine, it would be more advisable to use *Pongamia pinnata* since it has proven to be effective both as a preventive and curative agent.

ETHICAL APPROVAL

Animal Ethic committee approval has been taken to carry out this study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Chippaux JP, Goyffon M. Epidemiology of envenomations in the world. In: Mion G, Goyffon M, editors. Serious envenomation. Paris: Arnette. 2000;1-7.
2. Larréché S. Envenomation by vipers in the Republic of Djibouti from October 1994 to May 2006: retrospective study in the intensive care unit of the Bouffard medico-surgical group. Doctorate of state in medicine. Val-de-Marne, France: Faculty of Medicine of Créteil, University of Paris. 2007; 155. English.
3. Pollet L, Ponchel C, Carpentier JP. Cobraic envenomation. In: Mion G, Goyffon M, Serious envenomations, Paris: Arnette. 2000; 53-64.
4. Chippaux JP. Estimate of the burden of snakebites in sub-Saharan Africa: A meta analytic approach. *Toxicon*. 2011; 57(4): 586-99. DOI: 10.1016/j.toxicon.2010.12.022, PMID 21223975.

5. Chippaux JP. Snake venoms and envenomations. Paris, France: RID Editions Research Institute for Development, Didactic Collection. 2002; 293. English.
6. Dramé BSI. Envenomation accidents by snakebite in the Surgical Emergency Department of the Gabriel Touré Hospital. State Doctorate in Medicine. Faculty of Medicine, Pharmacy and Odontostomatology, University of Mali. 2000; 72. English.
7. Grema M, Koné PP. Effects of the venom of a snake (*Bitis arietans*) and of an antivenomous plant from the traditional African pharmacopoeia (*Annona senegalensis*) on the junction between the sciatic nerve and the gastrocnemius muscle of a toad (*Bufo regularis*). Review CAMES-series A. English. 2003; 02:79-85.
8. Mion G, Olive F. Viperid envenomation (in Africa). Resusc Trop Med, Paris, France. 1998; 349-65. English.
9. Kassogué A. Renal complications of snakebite envenomation at SUC HGT from January to December 2004. Doctorate of State in Medicine. Mali: Faculty of Medicine, Pharmacy and Odontostomatology, University of Bamako. 2006; 86. English.
10. Mion G, Larréché S. Cobra syndrome. Trop Med. 2008; 68:348-58. English.
11. Larréché S, Mion G, Clapson P, Debien B, Wybrecht D, Goyffon M. Ophidian neurotoxins. French Ann Anesth Resusc. 2008; 27(4):310-6. English.
12. Nelson BK. Snake envenomation Incidence. Clinical presentation and management. Med Toxicol Adverse Drug Experience. 1989;4(1):17-31. DOI: 10.1007/BF03259900, PMID 2468987.
13. Brown GV, Warrell DA. Venomous bites and stings in the tropical world. Med J Aust. 1993; 159(11-12):773-9. DOI:10.5694/j.1326-5377.1993.tb141345.x, PMID 8264466.
14. Hamza L. Biological and biochemical characterization of hemorrhagic and myonecrotic activities of *Vipera lebetina* venom. Magister in Biochemistry-Immunology, Faculty of Biological Sciences. Algeria: University of Sciences and Technology Houari Boumediene. 2001; 127. English.
15. Candan F, Unlu M, Tepe B, Daferera D, Polissiou M, Sökmen A et al. Antioxidant and antimicrobial activity of the essential oil and methanol extracts of *Achillea millefolium* subsp. *millefolium* Afan. J Ethnopharmacol. 2003; 87(2-3):215-20. DOI: 10.1016/s0378-8741(03)00149-1, PMID 12860311.
16. Lagnika L, Amoussa M, Adjovi Y, Sanni A. Antifungal, antibacterial and antioxidant properties of *Adansonia digitata* and *Vitex doniana* from Benin pharmacopeia. J Pharmacogn Phytother. 2012; 4(4): 44-52. DOI: 10.5897/JPP12.006
17. Dinzedi MR. Antibacterial activities of extracts of *Terminilia catappa* and *Thonningia sanguinea* on *Escherichia coli*, *Klebsiella pneumoniae* and multiresistant *Staphylococcus aureus* of human origin [university doctoral thesis] Specialty: Pharmacology of Natural Substances. Option: Biochemistry Microbiology. Côte d'Ivoire: Biosciences TRU, Félix Houphouët-Boigny Abidjan University. 2015; 133. English.
18. Okou OC, Yapo Sopié E-S. Antibacterial activity of *Solanum torvum* leaves: Extracts of *Solanum torvum* leaves (Solanaceae) on the in vitro growth of 3 strains of enterobacteriaceae. European University edition, Berlin, Germany. English. 2018; 1-43.
19. Naceiri Mrabti H. Pharmacological and toxicological study of *Arbutus unedo* L. in Morocco. Doctorate in Medicine Sciences. Morocco: Faculty of Medicine and Pharmacy of Rabat, Centers for Doctoral Studies of Life and Health Sciences, Mohammed V University of Rabat. 2018; 134. English.
20. Adjanohoun EJ, Aké-Assi L. Contribution to the census of medicinal plants in Côte d'Ivoire CRES. National floristic center, University of Côte d'Ivoire. English. 1979; 358.
21. Okou OC. Efficacy and spectrum of activity of *Mitracarpus scaber* Zucc extracts. Ex Schult + Scult.f. (Rubiaceae) and fusidic acid on Gram-Positive Cocci Bacteria [doctoral thesis in Biochemistry]-Pharmacology. Abidjan, Côte d'Ivoire: Félix Houphouët-Boigny University. 2012; 229. English.
22. Chippaux JP. Snake bite epidemiology in Benin (West Africa). Toxicol. 1989; 27: 127-35.
23. Koné PP. Toxicological, electrophysiological and pharmacological

- study of the venom of *Naja nigricollis* (Elapidae from Côte d'Ivoire) and of an antivenom substance from the African pharmacopoeia, the aqueous extract of *Securidaca longepedunculata* [Doctoral thesis], Sciences TRU. University of Côte d'Ivoire. 1980; 172. English.
24. Somé N, Poda JN, Guissou IP. Epidemiology and management of ophidian envenomations in the district of Dano health, province of Ioba (Burkina Faso) from 1981 to 2000. In: Envenomation and its treatment in Africa. Bulletin of the Exotic Pathological Society. 2002; 95(3):163-66. English
 25. Fumba G. Medicinal antivenom plants from Burundi. The presses of the future, Arlon, Belgium. 1983; 136. English
 26. Chippaux JP. Snakes of West and Central Africa. 3rd ed revised and expanded, Research Institute for Development RID edition. Paris, France. 2006; 329. English.
 27. Djerrou Z. Study of the pharmacotoxicological effects of medicinal plants from Algeria: Healing activity and harmlessness of the vegetable oil of *Pistacia lentiscus* L [Doctoral thesis in Sciences], Faculty of Nature and Life Sciences, Mentouri University from Constantine. Algeria. 2011; 131. English.
 28. Coulibaly FA, Coulibaly A, N'guessan JD, Kouamé KG, Djaman AJ, Guédé-Guina F. Study of biochemical serum parameters: the case of rabbits (*Neozelandais-cunistar*) from Côte d'Ivoire. Sci Nat. 2007; 4(1):37-43. English.
 29. Abenga JN, Anosa VO. Serum total proteins and creatinine levels in experimental Gambian trypanosomiasis of vervet monkeys. Afr J Biotechnol. 2005; 4:187-90.
 30. Hilali M, Abdel-Gawad A, Nassar A, Abdel-Wahab A. Hematological and biochemical changes in water buffalo calves (*Bubalus bubalis*) infected with *Trypanosoma evansi*. Vet Parasitol. 2006; 139(1-3): 237-43. DOI: 10.1016/j.vetpar.2006.02.013, PMID 16567045.
 31. Zabre ZM. Determination of the usual biochemical parameters in small ruminants from Burkina Faso and their variations in subjects naturally infected with trypanosomiasis [Veterinary thesis]. Dakar, Senegal: Faculty of Medicine, Pharmacy and Odonto-Stomatology, Cheikh Anta Diop University. 2013; 71. English.
 32. Obou Constantin Okou, Djako Sosthène Thierry Akéré, Gomé Michel Bleu, Assandé Francis Ehounou, Alain Dit Philippe Bidié. Study of the variation of some biochemical parameters in *Rattus norvegicus*. Asian J Res Biochem. 2020; 7(4):21-34.
 33. Al-Shinnawy M. Vitamin C as ameliorative agent against thiodicarb toxicated male albino rats (*Rattus norvegicus*). Egypt Acad J Biol Sci. 2008; 1(2):177-87.
 34. Djeffal A. Evaluation of the toxicity of a "methomyl" carbamate insecticide in Wistar rats: oxidative stress and exploration of the protective effects of selenium and/or vitamin C supplementation. 3rd Cycle Thesis. Option: Applied biochemistry. Algeria: Faculty of Sciences, Badji Mokhtar-Annaba University. 2014; 170. English.
 35. Lewington S, Whitlock G, Clarke R, Sherliker P, Emberson J et al. Blood cholesterol and vascular mortality by age, sex, and blood pressure: A meta-analysis of individual data from 61 prospective studies with 55,000 vascular deaths. Lancet. 2007; 370(9602):1829-39. DOI: 10.1016/S0140-6736(07)61778-4, PMID 18061058.
 36. Heuillet M. Development of reference methods for lipid profile biomarkers: Application to quality control in clinical biology [doctoral thesis], Discipline. Paris, France: Life Sciences, University of Burgundy. p. 199. English; 2013.
 37. Diaby V. Biochemical, hematological and histological studies of cadmium sulphate in Wistar rats [Doctoral thesis], Speciality: Functional and Molecular Biology. Côte d'Ivoire: Biosciences TRU, Félix Houphouët-Boigny Abidjan University. 2017; 130. English.
 38. Siby S. Study of the variation of biochemical and hematological parameters in the district of Bamako [State thesis in Medicine]. Mali: Faculty of Medicine, Pharmacy and Odonto-Stomatology, University of Bamako. 2008; 77. English.
 39. Carip C Chapter 4. Hepatobiliary pathology. In: Physiopathology: pathophysiology: pathophysiological basis of dietetics, Lavoisier TEC & doc, Paris, France. 3rd ed. 2014; 71-87. English.
 40. Ichai C. Dyskalaemias. 2012; 20. Available: https://sofia.medicalistes.fr/spip/IMG/pdf/Les_dyskaliemies.pdf. English.

41. Anonymous. Potassium balance abnormalities. English 2018:43-60
42. Eastham Rd. Compendium of biological constants. Paris, France: Masson. 1978; 248. English.
43. Isler C. Evolution of biochemical parameters during displacement to the left of the abomasum in dairy cows: study of four cases [veterinary thesis], National Veterinary School. Claude: Bernard University, Lyon I. France. 2007; 111. English.
44. Pitel Ph Moulin M, Valette JP, Dumontier S, Petit L, Fortier G, Courouc -Malblanc A. Approach to hematological and biochemical values in two donkey breeds. Equine Vet Pract. 2006; 38:19-25. English.
45. Pritchard JC, Burn CC, Barr ARS, Whay HR. Haematological and serum biochemical reference values for apparently healthy working horses in Pakistan. Res Vet Sci. 2009; 87(3):389-95. DOI: 10.1016/j.rvsc.2009.05.003, PMID 19552930.
46. Grenier-Michaud S, Lyne C, Pierre N. Understanding renal function is an essential component of paraclinical monitoring. Nurs Perspect. 2011; 30-5. English.
47. Dufour DR, Lott JA, Nolte FS, Gretch DR, Koff RS, Seeff LB. Diagnosis and monitoring of hepatic injury II. Recommendation for use of laboratory tests in screening, diagnosis and monitoring. Clin Chem. 2000; 46(12):2050-68. DOI: 10.1093/clinchem/46.12.2050, PMID 11106350.
48. Gom  MB, Kouakou K, Tour  A, Traor  F. Study of the acute and subchronic toxicity of the aqueous extract of *Passiflora foetida* Linn. (Passifloraceae) in rats and mice. Int J Biol Chem Sci. 2011; 5(5):1777-89. English.
49. Kaneko JJ, Harvey JW, Bruss ML. Clinical biochemistry of domestic animals. 5th ed. San Diego: Academic Press. 1997; 932.
50. Lahoz C, Mostaza JM. Atherosclerosis as a systemic Disease. Rev Esp Cardiol. 2007; 60(2):184-95. DOI: 10.1157/13099465, PMID 17338883.
51. Bidi  ADP, Adeoti FM, Yapo FA, Tiekpa JW, N'guessan JD, Djaman JA. Effect of the total aqueous extract of *Chrysophyllum perpulchrum* on hematological, biochemical parameters and weight growth of healthy Wistar rats. Ivorian J Sci Technol. 2016; 28:333-48. English.
52. Goyffon M, Chippaux JP. Poisonous terrestrial animals. Encyclopedia of medicine and surgery, Paris, France. English. 1990; 14.

  2023 Okou et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/97893>