



Characteristics of Whole Wheat, Red Kidney Bean and Defatted Coconut Flour Blends and Its Application in Bread Production

Vachefon Heuvey Forwoukeh ^{a*}, Julius Amove ^b
and M. I. Yusufu ^c

^a Centre for Food Technology and Research, CEFTER, Benue State University, Makurdi, Nigeria.

^b Department of Food Science and Technology, Joseph Sarwuan Tarka University, Makurdi, Nigeria.

^c Department of Food Science and Technology, Federal Polytechnic, Idah, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Author VHF did conceptualization, performed methodology, and helped in software development of the manuscript. Authors JA and VHF did data validation of the study. Author VHF wrote and prepared the original draft of the manuscript. Authors VHF, JA and MIY wrote, reviewed and edited the manuscript. Authors JA and MIY supervised the study. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AFSJ/2023/v22i9655

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/95440>

Original Research Article

Received: 12/11/2022

Accepted: 16/01/2023

Published: 13/07/2023

ABSTRACT

Aim: To evaluate the functional, proximate and anti-nutrient content of flour blends and sensory properties of bread produced from blends.

Methodology: Flour was prepared from whole wheat (WW), red kidney bean (RKB) and coconut fruit (CF). It was than blend as follows: S1 (100% refined wheat), S2 (100% WW), S3 (90% WW, 5% RKB and 5% CF), S4 (85% WW, 10% RKB and 5% CF), S5 (80% WW, 15% RKB and 5% CF), S6 (75% WW, 20% RKB and 5% CF). Blends were subjected to functional, proximate and anti

*Corresponding author: Email: forwoukehheuvey@gmail.com;

nutrient content analyses. Also the sensory attributes of breads produced from blends were determined.

Results: Water absorption, oil absorption, swelling, foaming, foaming stability and emulsification capacity increased respectively from 0.62 ± 0.06 g/ml (S1) to 0.92 ± 0.01 g/ml (S6), 0.96 ± 0.02 g/ml (S1) to 1.42 ± 0.03 g/ml (S6), 1.05 ± 0.01 (S1) to 1.73 ± 0.01 (S6), $11.47\pm 0.02\%$ (S1) to $13.20\pm 0.05\%$ (S6), $50.93\pm 0.01\%$ (S1) to $53.47\pm 0.01\%$ (S6), $41.43\pm 0.03\%$ (S1) to $43.36\pm 0.01\%$ (S6) while bulk density decreased from 0.84 ± 0.01 g/ml (S1) to 0.60 ± 0.02 g/ml (S6). Moisture, ash, fat, fiber, and protein content increased respectively from $12.23\pm 0.03\%$ (S1) to $13.44\pm 0.01\%$ (S6), $0.66\pm 0.01\%$ (S1) to $1.66\pm 0.02\%$ (S6), $0.22\pm 0.3\%$ (S1) to $2.57\pm 0.01\%$ (S6), $0.67\pm 0.26\%$ (S1) to $4.34\pm 0.02\%$ (S6), $10.70\pm 0.12\%$ (S1) to $13.06\pm 0.03\%$ (S6) while carbohydrate decreased from $75.52\pm 0.33\%$ (S1) to $64.93\pm 0.04\%$ (S6). Phytate, oxalate and tannins content increased respectively from 0.03 ± 0.00 mg/100g (S1) to 1.09 ± 0.01 mg/100g (S6), 0.02 ± 0.02 mg/100g (S1) to 0.14 ± 0.01 mg/100g (S6), 0.01 ± 0.01 mg/100g (S1) to 0.09 ± 0.01 mg/100g (S6). In term of overall acceptability sample S1 and S3 showed no significantly different ($p > 0.05$).

Conclusion: Addition of red kidney beans and coconut flour into whole wheat had significant ($P < 0.05$) impact on the functional, proximate characteristic of flour. There sensory scores showed S3 to be suitable for bread.

Keywords: Anti-nutrients; blends; flour; functional; proximate.

1. INTRODUCTION

Wheat (*Triticum aestivum*) is the world's third most important cereal crop after maize and rice that grows mostly in the temperate region [1]. Wheat is the major ingredient for the baking industries due to the presence of gliadins and glutenins collectively called gluten proteins wide variety of cereal [2]. "The demand for the production of refined and whole meal flour has greatly increased during the last decades" [3]. "Whole wheat is rich in nutritive compounds in comparison with refined flour" [4]. "In addition to carbohydrate and protein whole wheat also contains minerals, fiber, B vitamins and bioactive compounds having high health-promoting properties" [5]. "Thus, it helps to reduce the risk of several diseases, such as cardiovascular disorders, type 2 diabetes and some cancers, as well as improving intestinal health" [3]. Wheat which is the major ingredient in bread production is not readily cultivated in the tropics hence most of the wheat flour in developing countries are imported [6,7]. This has resulted to high cost of production of baked products especial bread with low protein content in the tropic [8,7]. Substitutions of wheat with other local raw materials in the bakery can be undertaken in order to reduced the cost of production and improve the nutritional value of bread. Research efforts have been geared towards finding a local and cheap substitute for wheat flour and this has led to the concept of composite flour [9,10].

"Red kidney bean (*Phaseolus vulgaris*) a grain legume, is one of the neglected tropical legumes

that can be used to fortify cereal-based diets especially in developing countries, because of its high protein content" [11]. "It is also a rich source of vitamin, minerals and relatively high in crude fibre" [10]. Red kidney bean is one such protein source, which when used in the fortification or enrichment of cereal – based diets could go a long way in improving their nutritional status [10]. "Low cost, protein-rich and high energy food formulation based on cereal legume mixtures have been suggested" [3]. "The enrichment or fortification of traditional cereal based diets with other protein sources such as oilseeds and legumes has received considerable attention. This is because oil seed and legumes proteins are rich in lysine, but deficient in sulphur containing amino acids" [4]. "Legumes generally contain relatively high amount of protein than other plant food stuffs. Cereals (wheat) have low protein content and are in general deficient in lysine but are adequate in sulphur containing amino acids. Legume proteins are mainly used in food formulations to complement the protein in cereal grains" [12].

"Coconut (*Cocos nucifera*) contains higher amounts of dietary fiber (60 g/100 g) and other nutrients" [13]. "Coconut contains low amount of digestible carbohydrates, and has no gluten" [14]. "The dietary fiber in coconut has significant health benefits. These include the inhibition of chronic diseases such as cancer, cardiovascular diseases and diabetes mellitus. Also, it has the ability to reduce weight" [2]. "Coconut has a glycemic lowering effect. Low glycemic index (GI: 35) food particularly such containing high dietary-

fiber, has been demonstrated to moderate post-prandial blood glucose and insulin responses enhancing blood- glucose and lipid concentrations in humans and patients having diabetes mellitus” [13]. “Moreover, coconut is known for the production of coconut milk, coconut oil and flour. Coconut flour can be utilized to produce several delicious pies, cookies, cakes, breads, snacks and desserts” [14] The use of coconut flour in food formulation does not only improve the nutritional value of the product but give unique aroma and taste [15]. Therefore, the objective of this study was to assess the characteristics of whole wheat, red kidney bean and defatted coconut flour blends and its suitability in bread making.

2. MATERIALS AND METHODS

2.1 Materials Procurement

Whole wheat grains and coconut fruits were purchased respectively from high- level and railway Markets Makurdi metropolis of Benue state Nigeria while red kidney bean was purchased from Ndop main market, North West region Cameroon.

2.2 Preparation of Materials

2.2.1 Preparation of whole wheat flour

Whole wheat grains were processed into flour following the method described by Ndife et al. [4]. The grains were cleaned from dirt by sorting out contaminants such as sticks, leaves and sand, washed in potable water and dried at 55°C for six hours in a conventional air oven. The grains were then milled in attrition mill, sieved to pass through 0.5mm mesh sieve, packaged in an air tight container.

2.2.2 Preparation of red kidney bean flour

The method described by Ukeyima et al. [10], was followed in the preparation of red kidney bean flour. “The kidney bean seeds were cleaned to remove foreign materials, defected seeds, insect etc. the healthy beans were washed and conditioned for 12 hours. After 12 hours of conditioning, the seeds were blanched at temperature of 85°C for 30 minutes. The water was drained and allowed to cool to prevent further cooking. The beans were dehulled to remove the skin. It then was washed and placed in a hot air oven to dry at 55°C for 20 hours. The dried beans were milled using an attrition mill into flour. It was then sieved using a 0.05 mm mesh sieve, packaged in an air tight container” [10].

2.2.3 Preparation of defatted coconut flour

The method described by Afoakwah et al. [16], was followed in the preparation of red kidney bean flour. “The coconuts were de-husked and the liquid drained. The meat was cut into smaller pieces using a paring knife. It was then transferred into a liquidizer containing 0.5 L of boiling water and was blended for 5 min, until the pulp was smooth. The milk was separated from the meal using cheesecloth. The meal was washed again in hot (100°C) water to reduce the oil content. The residue obtained was then weighed and dried using Hot Air Oven Dryer at 60°C until a constant weight was attained. The dried coconut was then milled into flour using plate mill. The coconut flour was sieved using a 0.5mm mesh sieve, packaged in an air tight container” [16]. All the flour was mix at different percentages to give composite as shown in Table 1.

Table 1. Blend Formulation of Composite Flour

Samples	Refined Wheat flour	Whole wheat flour	Red kidney bean flour	Coconut flour
S1 control	100	0	0	0
S2 control	0	100	0	0
S3	0	90	5	5
S4	0	85	10	5
S5	0	80	15	5
S6	0	75	20	5

2.2.4 Preparation of bread

The method used for the preparation of bread samples was the straight - dough method as described by Adelekan and Alamu [17]. "All the ingredients were weighed and poured into a mixing bowl of the mixing machine. The ingredients were mixed with the aid of a mixer using slow speed for 5 minutes. Little water was added as the mixing continued, using high speed for 15 minutes. The dough was then put on a moulding table and moulded into desired shape (by hand). After cutting and weighing, the moulded dough was placed inside a lubricated baking pan and covered with a lubricated lid. It was transferred into a proofing chamber for 1hour 30 minutes in order to enhance fermentation and dough development, it was then transferred into the oven at temperature 200-210°C for 25 minutes" [17]. The bread was removed from the oven and cooled Fig. 1.

2.3 Methods of Analysis

2.3.1 Functional properties of refined wheat, whole wheat, red kidney bean and defatted coconut flour blends

2.3.1.1 Water absorption capacity (WAC)

The water absorption capacity was determined using the method described by Chandra and Samsher [18]; Horsfall et al. [19]; Ojo et al. [20]. 10mL of distilled water was added to 1g of the composite flour sample in a weighed centrifuge tube. The tube was agitated on a vertex mixer for

two minutes and then centrifuged at 500 rpm for 30 minutes. The clear supernatant was decanted and discarded. The adhering drops of water are removed and then weighed. Water absorption capacity was expressed as the weight of water bound by 100 g of dried flour.

$$\text{Water absorption capacity (ml/g)} = \frac{\text{volume of water absorbed}}{\text{mass of sample (g)}}$$

Water absorbed (ml) = Volume of water added - Volume of water obtained after centrifugation.

2.3.1.2 Oil absorption capacity (OAC)

The oil absorption capacity was determined using the method described by Chandra and Samsher [18]; Ojo et al. [20]. One gram (1g) of the composite flour sample was mixed with 10 millilitres (ml) of refined vegetable oil and allowed to stand at ambient temperature (30±2°C) for 30 minutes. It was then centrifuged for 30 minutes at 500 rpm. The oil and adhering drops of oil was decanted and discarded. Oil absorption capacity was expressed as percent oil bound per gram flour.

$$OAC = \frac{W2 - W3}{W1}$$

W3= Weight of sample after centrifugation, W2= Weight before centrifugation, W1= Weight of original sample




Fig. 1. Flow chart for preparation of bread by Adelekan and Alamu [17]

2.3.1.3 Bulk density (BD)

Bulk density of the flour blends was determined using the method described by Ukeyima et al. [10]. The flour blend sample (5 g) was poured into a (10 ml) dry measuring cylinder. The bottom of the cylinder was tapped 50 times (until a constant volume achieved) on the laboratory table and the volume was recorded for packed bulk density.

$$\text{Bulk density } \left(\frac{\text{g}}{\text{ml}} \right) = \frac{\text{weight of sample}}{\text{volume of sample after trapping}}$$

2.3.1.4 Swelling capacity (SC)

The swelling capacity was determined using the method described by Ukeyima et al. [10]. Five (5) gram of the sample was weighed into a clean and dry graduated measuring cylinder. Fifty (50) ml of boiled distilled water was added and stirred with a glass rod. It was then allowed to stand for 15 minutes and change in volume of the sample was recorded. The ratio of the initial volume to the final volume as swelling Index was calculated as:

$$\text{swelling capacity} = \frac{\text{change in volume of sample (ml)}}{\text{original volume of sample (ml)}}$$

2.3.1.5 Foaming capacity (FC) and Foaming stability (FS)

The foam capacity (FC) and foam stability (FS) were determined as described by Ukeyima et al. [10]. The 1.0 g flour sample was added to 50 mL distilled water at $30 \pm 2^\circ\text{C}$ in a graduated cylinder. The suspension was mixed and shaken for 5 minutes to foam. The volume of foam at 30 sec after whipping was expressed as foam capacity using the formula:

$$\text{Foaming capacity (\%)} = \frac{\text{volume of foam AW} - \text{volume of foam BW}}{\text{volume of of foam BW}} \times 100$$

Where, AW = after whipping, BW = before whipping.

The volume of foam was recorded two minutes after whipping to determine foam stability as per percent of initial foam volume.

2.3.1.6 Emulsion activity

The emulsion activity was determined using the method described by Horsfall et al. [19] The emulsion, 1g of the composite flour sample, 10 ml refined vegetable oil and 10 ml distilled water is prepared in a calibrated tube. The emulsion was centrifuged at 200 rpm for 15 minutes. The ratio of the height of the emulsion layer to the total height of the mixture was calculated as the emulsion activity expressed in percentage.

$$\text{Emulsion activity} = \frac{\text{height of emulsion layer}}{\text{height of mixture}} \times 100$$

2.4 Proximate Composition of Whole Wheat, Red Kidney Bean and Defatted Flour Blends

2.4.1 Moisture content determination

Moisture content (MC) was determined by oven method described by Ukeyima et al. [10]. Two 2g of each sample was weighed into a weighed dried dish (moisture cans). It was then placed in an oven at 103°C for three hours. Then it was removed and transferred into desiccators at room temperature to cool. After cooling, the samples were weighed and replaced in the oven for another 1 hour. After this, the dish containing the sample was removed and cooled in desiccators and reweighed. The procedure was repeated until a constant weight was obtained. The loss in weight was reported as moisture content and calculated as:

$$\%MC = \frac{\text{initial mass of sample} - \text{mass after drying}}{\text{initial mass of sample}} \times 100$$

2.4.2 Ash content determination

Ash was determined by the method outlined by Ukeyima et al. [10]. The crucible was weighed and dried in an oven and cooled in the desiccators. 5g of the each sample was weighed in the empty porcelain crucible. The crucible containing the samples were placed in the muffle furnace and ignited at 550°C . This temperature was maintained for three hours after which the crucibles were then transferred into the desiccators, cooled and weighed. The ash content was calculated as follows;

$$\% \text{ ash} = \frac{\text{weight of crucible + ash} - \text{weight of empty crucible}}{\text{weight of sample}} \times 100$$

2.4.3 Crude fibre content determination

The crude fibre of the flour samples was determined according to the method describe by Ukeyima et al. [10]. "Two grams each of the samples were boiled under reflux for thirty minutes with 200 mL of the solution containing 1.25 g of H₂SO₄ per 100 ml of solution. The solution was filtered through linen on a flaunted funnel and washed with water until the washing is no longer acidic. The residue was then transferred to a beaker and boiled for 30 minutes with solution containing 1.25 g of NaOH per 100 ml of solution. The final residue was filtered through a thin but closer pad of washed and ignited asbestos in a Gosh crucible. The residue was then dried in an electric oven and weighed; the residue was ashed in a muffle furnace at 550 °C for 3hours, cooled and weighed" [10].

$$\% \text{ of crude fiber} = \frac{(W2 - W3)}{W1} \times 100$$

Where:

W1 = weight of sample used,
W2 = weight of crucible plus sample,
W3 = weight of sample crucible + ash.

2.4.4 Fat content determination

The fat content of the flour samples was determined using the method describe by Ukeyima et al. [10]. "Five (5) grams each of the flour samples in a filter paper and placed in a soxhlet reflux flask which is connected to a condenser on the upper side and a weighed oil extraction flask full with two hundred ml petroleum ether. The ether was brought to its boiling point, the vapour condensed into the reflux flask immersing the samples completely for extraction to take place on filling up the reflux flask siphons over carrying the oil extract back to the boiling solvent in the flask. The process of boiling, condensation, and reflux was allowed to go on for four hours before the defatted samples were removed. The oil extract in the flux was dried in the oven at 60°C for 30 minutes and then weighed" [10].

$$\% \text{ fat} = \frac{\text{mass of fat obtained}}{\text{dry mass of sample used}} \times 100$$

2.4.5 Protein content determination

Crude protein of the sample flours was determined using the Kjeldahl method as describe by Ukeyima et al. [10]. "One gram of the sample was introduced into the digestion flask. Kjedadahl catalyst (Selenium Tablets) was added to the sample. Twenty ml of concentrated sulphuric acid was added to the sample and fixed to the digester for eight hours until a clear solution was obtained. The cooled digest was transferred into one hundred ml volumetric flask and made up to the mark with distilled water. The distillation apparatus was set and rinsed for ten minutes after boiling. 20ml of 4% boric acid was pipetted into a conical flask. Five drops of methyl red were added to the flask as an indicator and the sample was diluted with seventy-five ml distilled water. Ten ml of the digest was made alkaline with twenty ml of NaOH (20%) and distilled. The steam exit of the distillatory was closed and the change of color of the boric acid solution to green was timed. The mixture was distilled for fifteen minutes. The filtrate was then titrated against 0.1 N HCl. The percentage total was calculated" [10]:

$$\% \text{ of total nitrogen} = \frac{(Vb - Va) \times 0.1N \times 0.0142}{W} \times 100$$

Where: Va = Volume of acid used in titration, Vb = Volume of base, W =weight of sample used.

$$\% \text{Crude protein} = \% \text{ of total Nitrogen} \times 6.25$$

2.4.6 Carbohydrate content determination

Total carbohydrate was determined by differences between 100 and total sum of the percentage of fat, moisture, ash, crude fibre and protein content [21].

$$\% \text{CHO} = 100 - (\% \text{Protein} + \% \text{Moisture} + \% \text{Fat} + \% \text{Ash} + \% \text{Fibre})$$

2.5 Anti-Nutritional Properties of Whole Wheat, Red Kidney Bean and Defatted Coconut Flour Blends

2.5.1 Phytate content determination

The phytate content was determined using the method described by Marolt and Kolar [22]. Two grams of composite flour sample was weighed into a 250ml conical flask and 100ml of 2% concentrated HCl was added. It was allowed to stand for 3hr and filtered. After filtration, 50ml of

the filtrate was pipetted into a 250ml beaker and 107ml of distilled water was added to improve acidity. Ten millimetre of 0.3% ammonium thiocyanate solution was added as an indicator. The solution was titrated with standard iron III chloride (FeCl_3) which contains 0.00195g iron/ml until a brownish yellow colour appear and persist for 5minutes. The phytic acid content was calculated using the formula below:

$$\text{Phytic acid (g/kg)} = \frac{0.00195 \times \text{volume of FeCl}_3 \text{ consumed} \times \text{DF}}{\times 1000 \text{ sample}}$$

Where DF= dilution factor

2.5.2 Oxalates content determination

Oxalate was determined by the method describe by Adeniyi et al. [23]. Two (2 g) of the sample was digested with 10 ml 6 M HCl for one hour and made up to 250 ml in a volumetric flask. The pH of the filtrate was adjusted with conc. NH_4OH solution until the colour of solution changed from salmon pink colour to a faint yellow colour. Thereafter, the filtrate was treated with 10 ml of 5% CaCl_2 solution to precipitate the insoluble oxalate. The suspension is now centrifuged at 2500 rpm, after which the supernatant was decanted and precipitate completely dissolved in 10 ml of 20% (v/v) H_2SO_4 . The total filtrate resulting from the dissolution in H_2SO_4 is made up to 300 ml. An aliquot of 125 ml of the filtrate was heated until near boiling point and then titrated against 0.05 M of standardized KMnO_4 solution to a faint pink colour which persisted for about 30 seconds after which the burette reading was taken. The oxalate content was evaluated from the titre value.

2.5.3 Tannin content determination

Tannin was determined by the method described by Nwokenkwo et al. [24] using Unico UV-2102 Spectrophotometer. Five hundred milligrams of the sample was weighed into a 100 ml plastic bottle and 50 ml of distilled water was added mixed thoroughly for 1 hour using a mechanical shaker. The sample was subsequently filtered into a 50 ml volumetric flask and made up to mark with distilled water. Then 5 ml of the filtrate was transferred into a tube and mixed with 3 ml of 0.1m FeCl in 0.1N HCl and 0.008M Potassium ferrocyanide (KCN) solution. The absorbance of the sample was measured spectrophotometrically at 120nm wavelengths for 10 minutes. A blank sample was prepared and the colour was developed and read at the same

wavelength. A standard was prepared using tannin acid to get 100ppm and measured. The total tannin content was calculated using the formula:

$$\text{tannin } \left(\frac{\text{mg}}{\text{cm}^3} \right) = \frac{\text{absorbance of sample}}{\text{absorbance of standard}} \times \frac{\text{conc. of standard}}{1}$$

2.6 Sensory Evaluation of Bread Produced from Whole Wheat, Red Kidney and Defatted Coconut Flour Blends

Sensory evaluation was carried out using a preference test as described by Hashim et al. [25]. A 9-point hedonic scale where 1 represents “extremely dislike” and 9 represents “extremely like” was used for this study. The organoleptic evaluation of the bread samples was carried out for consumer acceptance and preference using 20 semi- trained panelists. The properties evaluated were appearance texture, aroma, taste, and overall acceptability.

2.7 Data Analysis

Data obtained were analysed using the one-way ANOVA and mean separated using Duncan’s Multiple Range Test (DMRT) at 5% limit of significant using Statistical package for social science (SPSS) version 26.

3. RESULTS AND DISCUSSION

3.1 Functional Properties of Whole Wheat, Red Kidney Bean and Defatted Coconut Flour Blends

The results for the functional properties of the flour blends are presented in Table 2.

3.1.1 Water absorption capacity (WAC)

The water absorption capacity (WAC) of the flour blends ranged from $0.62 \pm 0.06 \text{g/ml}$ (S1) to $0.92 \pm 0.01 \text{g/ml}$ (S6). The results shows increased in WAC of flour blends with the addition of red kidney bean. The WAC of sample S1 (control 1) was significantly ($p < 0.05$) lower than that of the other samples. Sample S2 (control 2) was significantly ($p < 0.05$) lower than samples S4 to S6. The major chemical compositions that enhance the water absorption capacities of flours are proteins, fiber and carbohydrates since these constituents contain

hydrophilic parts such as polar or charged side chains. Therefore the increased in the WAC of the flour is due to increased in the protein and fiber content of the flour as the quantity of red kidney bean flour increased. The lower WAC of refined wheat flour could be attributed to the presence of lower amount of hydrophilic constituents in wheat such as polysaccharides [26]. This report agrees with results presented by Chandra and Samsher [18]; Kaushal et al. [27], who noted that flour with high protein and fiber content had high WAC. The values presented here are higher due to addition of coconut flour which increased both protein and fiber content of the flour samples. Very low or excessive water absorption can negatively affect the quality of food products. Water absorption can influence the following parameters of bread and baking: Proofing, Fracture stress of bread crumb, Loaf volume, and Bread yield [28]. Under-absorption of water result to Firm and dense internal structure (crumb) and Low volume bread while over-absorption of water result to open crumb grain with large cells, Large volume and bread which is prone to mold [28].

3.1.2 Oil absorption capacity (OAC)

Oil absorption capacity (OAC) of the flour blends ranged from 0.96 ± 0.02 g/ml (S1) to 1.42 ± 0.03 g/ml (S6). The oil absorption capacity increased from sample S1 to S6 with the addition of red kidney flour. The OAC of S2 to S6 were significantly ($p<0.05$) higher than S1 (control 1). The presence of high fat content in flours might have affected adversely the OAC of the composite flours. Therefore, the possible reason for increase in the OAC of composite flours after incorporation of red kidney bean and coconut flour is the variations in the presence of non-polar side chain, which might bind the hydrocarbon side chain of the oil among the flours [29]. The primary chemical component affecting OAC is protein that is composed of both hydrophilic and hydrophobic parts. Non-polar amino acid side chains can form hydrophobic interaction with hydrocarbon chains of lipids [28]. Increasing red kidney beans quantity was found to increase the protein content of the flour sample, resulting to increase in OAC. Similar findings were observed by chandra and Samsher [18]; Kaushal et al. [27], who noted flour of high protein retain more oil than flour with low protein content. The OAC makes the flour suitable in facilitating, enhancement in flavor and mouth feel when used in food preparation. Due to these properties, the protein probably could be used as

functional ingredient in foods such as bread and cakes [28].

3.1.3 Bulk density (BD)

The bulk density (BD) of the flour blends ranged from 0.84 ± 0.01 g/ml (S1) to 0.60 ± 0.02 g/ml (S6). There was a decreased in BD of the flour blends from sample S1 to sample S6 with the addition of red kidney bean. The BD of sample S1 (control 1) was significantly ($p<0.05$) higher than samples S2 to S6. BD depends on the particle size and moisture content of flours. Bulk density can be improved when the particles are smaller. The use of whole wheat, with the addition of red kidney bean and coconut flour increased the particles size, resulting to decrease in BD from sample S2 to S6. The variation in bulk density of foods could also be attributed to the variation in starch content of the foods. The higher the starch content the more likely the increase in bulk density [28][30]. This observation fails to agrees with works presented by [10] due to the addition of coconut flour which increased the particles size of the flour but agrees with report by Awuchi et al. [28], who observed that flour with fine particle had high BD than those with larger particles sizes. Bulk density reflects the relative volume or capacity of the required packaging material[28].

3.1.4 Swelling capacity (SC)

The swelling capacity (SC) of the flour blends ranged from 1.05 ± 0.01 (S1) to 1.73 ± 0.01 (S6). From the results, there was an increased in swelling index of flour blends from sample S1 to S6 with the addition of red kidney bean. Sample S1 (control 1) was significantly ($P<0.05$) lower than other samples. Samples S4 to S6 also show significant difference ($p<0.05$) from sample S2 (control 2). Swelling capacity of composite flours increased with increase in the level of incorporation ratio of red kidney bean flour. The swelling capacity (index) of flours are influenced by the particle size, species variety and method of processing or unit operations [18]. The increased in SC may also be attributed to the increased in the fiber and pre-gelatinized starch content of the composite flour (since red kidney bean was parboiled) [29]. "High starch content increases swelling capacity (index) of foods and flours, especially in starch with higher amount of the branched amylopectin. Starch is made up of amylose (linear chain) and amylopectin (branched chains), both are chains of glucose units. The amount and proportion of amylose and

amylopectin found in starch vary according to the plant source. This explains why different flours from different (plant) sources and species have different swelling capacities" [18]. Similar observation were made by Suresh et al. [29]; Moses and Olanrewaju [31]; Adejuyitan et al. [32], who stated that composite flour rich in fiber have high swelling capacity.

3.1.5 Foaming capacity (FC)

The foaming capacity (FC) of the flour blends ranged from $11.47 \pm 0.02\%$ (S1) to $13.20 \pm 0.05\%$ (S6). There was increased in foaming capacity of the flour blends from sample S1 to sample S6 with the addition of red kidney bean. The FC of S1 (control 1) was significantly ($p < 0.05$) lower than other samples. S2 (control 2) was significantly ($p < 0.05$) lower than S4 to S6. Protein is mainly responsible for foaming. Foaming capacity and stability generally depend on the interfacial film formed by the proteins, which maintains the suspension of air bubbles and slows down the coalescence rate [28]. Increase in level of substitution of red kidney bean flour result to increase in the foaming capacity of the flour samples. This observation agrees with reports by Ukeyima et al. [10]; Abdelghafor et al. [33], who reported that legumes have high foaming capacity due to the high protein content of the flour.

3.1.6 Foam stability (FS)

The Foam stability (FS) of the flour blends ranged from $50.93 \pm 0.01\%$ (sample S1) to $53.47 \pm 0.01\%$ (S6). There was increased in foaming stability (FS) of the flour from sample S1 to S6 with the addition of red kidney bean flour. There was significant difference ($p < 0.05$) between samples S3 to S6 and the controls (S1 and S2). Increased in foam stability is due to increase in the protein content of the flour as the quantity of red kidney bean and flour increased. Flours are capable of producing foams due to surface active proteins. "Foaming capacity and stability generally depend on the interfacial film formed by the proteins, which maintains the suspension of air bubbles and slows down the coalescence rate" [28]. "Protein in the dispersion may cause a lowering of the surface tension at the water air interface, thus always been due to protein which forms a continuous cohesive film around the air bubbles in the foam" [27]. Stable foams are known to occur when low surface tension and high viscosity occur at the interface forming vacuoles in the foam [28]. This

observation agrees with reports by Ukeyima et al. [10]; Abdelghafor et al. [33], who noted that Stable foams are known to occur when low surface tension and high viscosity occur at the interface forming a continuous cohesive film around the air vacuoles in the foam. But according to Kapil et al. [26], "there is always an inverse relationship between the foaming capacity and the foam stability. Flours with high foaming capacity may form large air bubbles encircled by thinner less flexible protein film. Consequently, this air bubbles may collapse easily and consequently lower the foam stability". This is true for flour with low protein content but not true in this study because of high protein content of red kidney bean which increased the protein value of flour creating a stable cohesive film around the air bubbles as also reported by Hyacinthe et al. [34].

3.1.7 Emulsification capacity (EC)

Emulsification capacity (EC) of the flour blends ranged from $41.43 \pm 0.03\%$ (sample S1) to $43.36 \pm 0.01\%$ (sample S6). Emulsification capacity (EC) of the flour blends increased from sample S1 to S6 with the addition of red kidney bean. The EC of sample S1 was significantly ($p < 0.05$) lower than samples S2 to S6. Sample S2 (control 2) was significant difference ($p < 0.05$) from samples S3 to S6. The Increased in emulsion capacity of the flour as the quantity of red kidney bean and coconut flour increased is due to increased in the protein content of the flour. "Emulsifiers like protein are compounds that typically have a non-polar (that is, lipophilic or hydrophobic) part and a polar or hydrophilic (water-soluble) part. Due to this, emulsifiers tend to be more or less soluble either in oil or in water. Emulsifiers which are more soluble in oil (and conversely, less soluble in water) will generally form water-in-oil emulsions, while emulsifiers that are more soluble in water, and less soluble in oil, will form oil-in-water emulsions. Emulsifying particles and emulsifiers (protein) tend to promote the dispersion of the food phases in which they are not well dissolved. For example, protein dissolves better in water than it does in oil, and as a result tend to form oil-in-water emulsions (i.e., protein promotes the dispersion of droplets of oil throughout the continuous phase of water)" [28]. This report agrees with work presented by Kausha et al. [27], who noted that increased in protein content of composite flour resulted to increased in emulsification capacity.

Table 2. Functional Properties of Refined Wheat, Whole Wheat, Red Kidney Bean and Defatted Coconut Flour Blends

Flour blends	WAC (ML/G)	OAC (ML/G)	BD (G/ML)	SC (ML/ML)	FC (%)	FS (%)	EC (%)
S1	0.62 ^D ±0.06	0.96 ^C ±0.02	0.84 ^A ±0.01	1.05 ^E ±0.01	11.47 ^D ±0.02	50.93 ^D ±0.01	41.43 ^E ±0.03
S2	0.71 ^C ±0.02	1.28 ^B ±0.03	0.78 ^B ±0.01	1.62 ^D ±0.03	12.98 ^C ±0.01	51.95 ^C ±0.06	43.05 ^D ±0.02
S3	0.75 ^C ±0.05	1.29 ^B ±0.01	0.75 ^B ±0.01	1.63 ^D ±0.01	12.99 ^C ±0.00	52.88 ^B ±0.02	43.09 ^C ±0.01
S4	0.83 ^B ±0.04	1.30 ^B ±0.09	0.69 ^C ±0.02	1.65 ^C ±0.01	13.09 ^B ±0.01	52.89 ^B ±0.01	43.24 ^B ±0.03
S5	0.88 ^A ±0.03	1.36 ^A ±0.03	0.63 ^D ±0.03	1.70 ^B ±0.00	13.11 ^B ±0.02	53.43 ^A ±0.02	43.33 ^A ±0.03
S6	0.92 ^A ±0.01	1.42 ^A ±0.03	0.60 ^D ±0.02	1.73 ^A ±0.01	13.20 ^A ±0.05	53.47 ^A ±0.01	43.36 ^A ±0.01

Values are means of triplicate determinations ± S.D. Means followed by different superscript letters in the same column indicate significant difference at (p<0.05).

Water absorption capacity (WAC), Oil absorption capacity (OAC), Bulk density (BD), Swelling Capacity (SC), Foaming Capacity (FC), Foam Stability (FS), Emulsification Capacity (EC)

3.2 Effect of Red Kidney Bean and Defatted Coconut Flour on the Proximate Composition of Whole Wheat Flour

The results for the proximate composition of the flour blends are presented in Table 3.

3.2.1 Moisture content

The moisture content of the flour blends ranged from $12.23 \pm 0.03\%$ (S1) to $13.44 \pm 0.01\%$ (S6). There was an increase in moisture content of the flour blends with the addition of red kidney bean. The increase in moisture content could be attributed to an increase in the fiber content of the flour. Whole wheat flour and mostly whole grains absorb more liquid than other flours such as white flour. In whole wheat flour, the whole grain is used, the grain is rich in fiber that loves to trap water and therefore the flour will be able to retain more moisture. This result agrees with the report presented by Okoye et al. [12], who reported that high fiber flour turns to hydrophilic. The results presented here are higher (but within the acceptable limits) due to the addition of coconut flour which further increased the hydrophilic property of the flour [35]. "The moisture content of flour is important for two reasons. First, the higher the amount of moisture content, the less the amount of dry solids in the flour. Secondly, flour with moisture content higher than 14-15% is not stable at room temperature as this is prone to microbial spoilage. The moisture content of all the blends were within the acceptable range of flour specification according to" Adelekan and Alamu [17].

3.2.2 Ash content

The ash content of the flour blends ranged from $0.66 \pm 0.01\%$ (S1) to $1.66 \pm 0.02\%$ (S6). There was an increase in the ash content of the flour blends from sample S1 to sample S6 with the addition of red kidney bean. There was a significant difference ($p < 0.05$) between the control (S1) and the other samples. Comparing sample S2 (Control 2) with samples S4 to S6, there was a significant difference ($p < 0.05$) with ash content. The increase in ash content of the flours could be due to the addition of red kidney bean and coconut flour, which are rich sources of minerals. This report is similar to findings presented by Okoye et al. [12]; Stoin et al. [36], who reported on improving the ash content of composite flour using legume. The values reported here are slightly higher due to

the addition of coconut flour, which has been reported as a good source of mineral [37][17].

3.2.3 Fat content

The fat content of the flour blends ranged from $0.22 \pm 0.3\%$ (S1) to $2.57 \pm 0.01\%$ (S6). There was an increase in the fat content of the flour blends from sample S1 to sample S6 with the addition of red kidney bean. The fat content of sample S1 was significantly ($p < 0.05$) lower than samples S2 to S6. Sample S2 (control 2) was significantly ($p < 0.05$) lower than the other samples (S3 to S6). The increase in fat content could be due to the use of whole wheat flour and the addition of defatted coconut flour, which may contain a certain percentage of oil. Red kidney bean, though very low in fat, could also contribute to an increase in fat content. This report fails to agree with findings reported by Okoye et al. [12]; Sachithra et al. [37], who reported that the addition of kidney bean reduces the fat content of the composite flour. This could be attributed to the addition of defatted coconut flour, which increased the fat content, contrary to their findings.

3.2.4 Fiber content

The fiber content ranged from $0.67 \pm 0.26\%$ (S1) to $4.34 \pm 0.02\%$ (S6). There was an increase in the fiber content of the flour blends with the addition of red kidney bean. There was a significant difference ($p < 0.05$) between samples S1 (control 1) and the other samples. The fiber content of sample S1 was significantly ($p < 0.05$) lower than the other samples. The increase in fiber content could be due to the high fiber content of whole wheat, red kidney bean, and coconut flour. This report is similar to work presented by Manonmani et al. [38]; Okoye et al. [12]; Adelekan and Alamu [17], who reported on the use of legumes and nuts for the development of functional foods due to their fiber content. Nutritionally, consumption of these experimental dough meals would be of benefit to the consumers, since evidences have shown that dietary fibers promote good health by preventing degenerative diseases like diabetes and hypertension. Fiber in food helps in burning of fat and boosting of the immune system. It could also provide bulk in the diet, enhance gastrointestinal function, and prevent constipation [39].

3.2.5 Protein content

The protein content ranged from $10.70 \pm 0.12\%$ (S1) to $13.06 \pm 0.03\%$ (S6). There was an

increased in the protein content of the flour blends with increased in red kidney bean addition. There was a significant difference ($p < 0.05$) between sample S1 (control 1) and the other samples. The protein content of samples S2 to S6 were significantly ($p < 0.05$) higher than sample S1. Sample S2 (control 2) and sample S3 did not show any significant difference ($p > 0.05$). Samples S4 to S6 show significant difference ($p < 0.05$) with sample S2 (control 2). The increased in protein content as the quantity of red kidney bean increased might be due to the high protein content of the red kidney bean as report by Manonmani et al. [38]; Kambabazi et al. [40]. This report is in agreement with report presented by Okoye et al. [12]; Akinjayeju et al. [41]; Adelekan and Alamu [17]; ukeyima et al. [10], who reported that combination of two or more varieties of cereals and legumes usually increased the protein content of the end products.

3.2.6 Carbohydrate content

The carbohydrate content of flour decreased from $75.52 \pm 0.33\%$ (S1) to $64.93 \pm 0.04\%$ (S6). The carbohydrate content of flour decreased from sample S1 to sample S6. The decreased could be due to increased in the addition of red kidney which is a low carbohydrate food. Also increase in other proximate components' result to decrease in carbohydrate since it is obtain by difference. This report agrees with finding by Okoye et al. [12]; Ndife et al. [4]; Ukeyima et al. [10], who reported a decreased in the carbohydrate content of their products when fortified with legumes.

3.3 Effect of Red Kidney Bean and Defatted Flour on the Anti-Nutrients Composition Whole Wheat Flour

The results for the anti nutrient composition of the flour blends is presented in Table 4.

3.3.1 Phytate content

The phytate content of the flour blends increased from $0.03 \pm 0.00 \text{mg}/100\text{g}$ (S1) to $1.09 \pm 0.01 \text{mg}/100\text{g}$ (S6). There was an increased in the phytate content of the flour blends from sample S1 to S2 with the addition of red kidney bean. The phytate content of sample S1 (control 1) was significantly ($p < 0.05$) lower than other samples. The increased in the phytate content of the flour samples could be due to the addition of red kidney beans flour which are known to

content phytate [42][22]. The phytate content of sample S1 was far lower due to refining (Refining of wheat grain remove part of the bran which contains highest percentage of phytate). The result of this study is similar to findings by Ohini et al. [43]. The values reported here are lower due to use of combined (soaking, fermentation dehulling and boiling) methods of reducing anti nutrient in food. These treatments methods have been reported to greatly reduced or eliminate phytate content of kidney bean. But the values are higher than those presented by Ayele et al. [44]. This might be due to the use of whole wheat flour in the research as part of the composite flour with legume. "Phytate are known to form complexes with iron, zinc, calcium and magnesium making them less available and thus inadequate in food samples especially for children. However, the phytate content in this study is far lower than the minimum amount of phytic acid reported to hinder the absorption of iron and zinc. It is known that 10-50 mg phytate per 100g will not cause a negative effect on the absorption of zinc and iron" [43].

3.3.2 Oxalate content

The oxalate content of the flour blends increased from $0.02 \pm 0.02 \text{mg}/100\text{g}$ (S1) to $0.14 \pm 0.01 \text{mg}/100\text{g}$ (S6). There was an increased in the oxalate content of the flour blends from sample S1 to sample S6 with the addition of red kidney bean. The oxalate content of sample S1 was significantly ($p < 0.05$) lower than that of samples S2 to S6. Comparing control 2 (sample S2) with sample S3 there was no significant difference ($p > 0.05$) but when compared with samples S4 to S6 there was a significant difference ($p < 0.05$). The increased in oxalate content could be due to the addition of red kidney bean flour which have been reported to contain oxalate [22][23]. The oxalate content of sample S1 was far lower equally due to refining (Refining of wheat grain remove part of the bran which contains the highest percentage of oxalate). This report agrees with findings by Adeniyi et al. [23]; Rasha et al. [42], but the values here are far lower due to soaking, fermenting and boiling of the raw material before milling to flour. These treatments have been reported to greatly reduced anti nutrients (oxalate) content of red kidney bean [45]. Oxalate is known to form complexes with calcium to form insoluble calcium-oxalate salt. According to ohini et al. [43] reported a safe range of 4- 9 mg/100g for oxalates. This implies that, the oxalates content of the flour samples was within the safe range.

3.3.3 Tannins content

The tannins content of the flour blends increased from $0.01 \pm 0.01 \text{ mg}/100 \text{ g}$ (S1) to $0.09 \pm 0.01 \text{ mg}/100 \text{ g}$ (S6). There was a significant ($p < 0.05$) increase in tannins content of flour from sample S1 to S6. The increase in the tannins content of the flour samples could be due to the addition of red kidney flour which has been reported to contain tannins [45]. The study agrees with findings by Mamiro et al. [46]; Pandit and Kaur [47]. But the values are lower due to soaking, fermenting and boiling of the raw material before milling to flour. These treatment methods have been reported to greatly reduce oxalate content of wheat red kidney bean [45]. On the other hand, the values are higher compared to those presented by Ayele et al. [44], due to the use of whole wheat flour in this research. The permissible limit of tannins in food is $20 \text{ mg}/\text{g}$, implying that the tannins content of the flour blends was within the acceptable limit.

3.4 Effects of Red Kidney Bean and Defatted Coconut Flour on the Sensory Attributes of Whole Wheat Bread

The results of the sensory attributes of the bread are presented in Table 5.

3.4.1 Appearance

The appearance scores ranged from 8.50 ± 0.61 like very much (S1) to 6.60 ± 0.60 like slightly (S6). Sample S1 has the best appearance while sample S6 has the least appearance with the addition of red kidney bean. There was a significant difference ($p < 0.05$) between sample S1 (control 1) and the other samples. Appearance was evaluated in terms of the crust and crumb colour and honey comb structure of the bread loaves. The colour of the samples changes from

white to dark as it moves from sample S1 (control 1) to sample S2 (control 2) and becomes darker as the substitution of red kidney beans flour increases. The dark bran of wheat flour is responsible for the dark colour of sample S2 (control 2). Also, coconut and red kidney beans flours are dark in colour and together with the dark bran of whole wheat flour are responsible for the darker appearances of samples S3 to S6, as compared to refined wheat flour. Again, the difference in colour could be as a result of the Millard browning reaction caused by the reaction between the wheat protein and sugars [48]. The darker colour of the crumbs of whole wheat bread and fortified breads has been reported by Adelekan and Alamu [17]; Ndife et al. [4]; Serrem et al. [49]. Sample S1 was found to develop the best honey comb (sponge) structure while sample S6 was found to develop the least. The reduction in the sponge structure of the bread samples was due to the dilution of the gluten content of the flour with non-wheat flour [50].

3.4.2 Aroma

The aroma scores ranged from 6.45 ± 0.76 like slightly (S2) to 8.55 ± 0.69 like very much (S3). The sensory scores for samples S3 to S6 were significantly ($p < 0.05$) higher than those of S1 and S2 (controls). This is because the incorporation of coconut flour resulted in very good flavour/aroma. This result did not agree with work done by Ndife et al. [4], who reported that an increase in the level of substitution of wheat with bean (legumes) flour results in a beany flavour and aroma from the beans in the composite breads. The result was different due to soaking, fermentation and blanching which help to get rid of the beany flavour as earlier reported by Ukeyima et al. [10], using white kidney bean in bread production. Also, coconut flour was reported by Adelekan and Alamu [17] to add good aroma to baked products.

Table 3. Proximate Composition (%) of Refined wheat, Whole Wheat, Red Kidney Bean and Defatted Coconut Flour Blends

Flour blends	Moisture	ASH	FAT	Fiber	Protein	Carbohydrate
S1	$12.23^D \pm 0.03$	$0.66^D \pm 0.01$	$0.22^E \pm 0.03$	$0.67^D \pm 0.26$	$10.70^D \pm 0.12$	$75.52^A \pm 0.33$
S2	$13.33^C \pm 0.03$	$1.51^C \pm 0.03$	$2.40^D \pm 0.01$	$4.07^C \pm 0.02$	$11.15^C \pm 0.06$	$67.54^B \pm 0.11$
S3	$13.34^C \pm 0.03$	$1.53^C \pm 0.03$	$2.44^C \pm 0.03$	$4.13^C \pm 0.03$	$11.21^C \pm 0.02$	$67.34^B \pm 0.02$
S4	$13.38^B \pm 0.01$	$1.58^B \pm 0.02$	$2.50^B \pm 0.01$	$4.20^B \pm 0.01$	$12.07^B \pm 0.01$	$66.27^C \pm 0.02$
S5	$13.40^{AB} \pm 0.02$	$1.60^B \pm 0.02$	$2.52^B \pm 0.02$	$4.30^A \pm 0.02$	$12.56^B \pm 0.01$	$65.62^C \pm 0.00$
S6	$13.44^A \pm 0.01$	$1.65^A \pm 0.02$	$2.57^A \pm 0.01$	$4.34^A \pm 0.02$	$13.06^A \pm 0.03$	$64.93^D \pm 0.04$

Values are means of triplicate determinations \pm S.D. Means followed by different superscript letters in the same column indicate significant difference at ($p < 0.05$)

Table 4. Anti-Nutrient Composition (mg/100g) of Refined Wheat, Whole Wheat, Red Kidney Bean and Defatted Coconut Flour Blends

Flour blends	Phytate	Oxalate	Tannins
S1	0.03 ^F ±0.00	0.02 ^D ±0.02	0.01 ^D ±0.01
S2	1.05 ^E ±0.00	0.06 ^C ±0.00	0.05 ^C ±0.01
S3	1.06 ^D ±0.00	0.09 ^C ±0.01	0.06 ^C ±0.01
S4	1.07 ^C ±0.00	0.11 ^B ±0.00	0.07 ^B ±0.00
S5	1.08 ^B ±0.00	0.12 ^B ±0.01	0.08 ^A ±0.01
S6	1.09 ^A ±0.01	0.14 ^A ±0.01	0.09 ^A ±0.01

Values are means of triplicate determinations ± S.D. Means followed by different superscript letters in the same column indicate significant difference at ($p < 0.05$).

Table 5. Sensory Attributes of Bread Produced from Refined Wheat, Whole Wheat, Red Kidney Bean and Coconut Flour Blends

Bread samples	Appearance	Aroma	Taste	Texture	Overall acceptability
S1	8.50 ^A ±0.61	6.55 ^B ±0.69	8.20 ^A ±0.59	8.40 ^A ±0.68	8.45 ^A ±0.39
S2	7.50 ^B ±0.51	6.45 ^B ±0.76	7.25 ^B ±0.91	7.25 ^B ±0.55	6.75 ^C ±0.91
S3	7.75 ^B ±0.72	8.55 ^A ±0.69	8.25 ^A ±0.87	6.60 ^C ±1.23	8.12 ^A ±0.96
S4	6.80 ^C ±0.77	8.45 ^A ±0.60	7.30 ^C ±0.86	6.40 ^C ±0.94	7.15 ^{BC} ±1.08
S5	6.70 ^C ±0.73	8.40 ^A ±0.68	6.90 ^{CD} ±0.91	5.75 ^D ±1.01	6.25 ^D ±1.29
S6	6.60 ^C ±0.60	8.30 ^A ±0.73	6.55 ^D ±0.69	5.55 ^D ±0.76	6.05 ^D ±1.39

Values are means of triplicate determinations ± S.D. Means followed by different superscript letters in the same column indicate significant difference at ($p < 0.05$).

3.4.3 Texture

The texture scores (softness and chewiness) of bread sample ranged from 8.40±0.68 like very much (S1) to 5.55±0.76 neither like nor dislike (S6). There was decreased in acceptability in terms of texture with increase in non-wheat flour substitution. The sensory scores for sample S3 to S6 was significantly ($p < 0.05$) lower than that of samples S1 and S2 (controls). Between the controls there was a significant difference ($P < 0.05$) in texture. Hard crumb texture of samples could be due to fibre from wheat bran and dilution in the gluten content of the flour [4]. This result agrees with works done by Eimam et al. [51]; Ukeyima et al. [10], who reported that the addition of non-wheat flour (legumes) into wheat flour increased the hardness of the bread due to dilution of gluten protein.

3.4.4 Taste

The taste scores ranged from 8.25±0.87 like very much (S3) to 6.55±0.69 like slightly (S6). Sample S3 was the most preferred and sample S6 was the least preferred in terms of taste. This could be due to the fact that coconut flour is known to introduce a characteristic taste to baked products as reported by Adelekan and Alamu [17]. But as the quantity of non-wheat flour (red kidney bean)

increased, the coconut taste is subdued resulting to a significant ($p > 0.05$) drop in sensory scores samples S4 to S6. This was because of the high fiber content of the samples which give the bread a blend taste. Samples S3, S1 shows no significant difference ($P > 0.05$).

3.4.5 Overall acceptability

The overall acceptability scores range from 8.45±0.39 like very much (S1) to 6.05±1.39 like slightly (S6). Samples S1, S3 shows no significant difference ($P > 0.05$) and were the most preferred. Sample S6 was the least preferred. Therefore it was concluded that good bread of acceptable sensory attributes can be produced from whole 90%, red kidney bean 5% and coconut flour 5% (sample S3).

4. CONCLUSION

The study has shown that flour of improved functional properties and proximate composition could be produced from blends of whole wheat, red kidney bean and defatted coconut flours blends. Also the result showed the anti-nutrients contents of composite flour were within the permissible level making the composite flour safe for human consumption. More so, the sensory scores indicate that bread of acceptable sensory

quality can be produced from blends (sample S3), since it shows no significant difference ($p>0.05$) with sample S1 (control).

ACKNOWLEDGEMENTS

The authors are grateful to chemistry department, Benue state university for supplying laboratory equipment used in this research work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Shewry PR. Darwin wheat review. J Exp Bot. 2009;60(6):1537-53. DOI: 10.1093/jxb/erp058, PMID 19386614.
2. Offia-Olua Bl. Chemical, functional and pasting properties of wheat (*Triticum spp*) - Walnut (*Juglans regia*) flour. Food Nutr Sci. 2014;05:1591-604. DOI: 10.4236/fns.2014.516172
3. Hassoon WH, Dziki D. Mi's, A. and Biernacka B. Wheat Grinding Process with Low Moisture Content: A New Approach for Wholemeal Flour Production, J urnal of Process. 2020;9(32):1-11.
4. Ndife J, Abdulraheem LO, Zakari UM. Evaluation of the nutritional and sensory quality of functional breads produced from whole wheat and soya bean flour blends. Afr J Food Sci. 2011;5(8): 466-72.
5. Călinoiu DC, Vodnar LF. Whole grains and phenolic acids: a review on bioactivity, functionality, health benefits and bioavailability. Cereal Chem. 2018;10 (1615).
6. Kasprzak M, Rzedzicki Z. Effect of pea seed coat admixture on physical properties and chemical composition of bread. Int Agrophys. 2010;24:149-56.
7. Ohimain EI. The prospects and challenges production in Nigeria Global Journal of Human-social Science. 2014;14(3).
8. Laze A, Arapi V, Ceca E, Gusho K, Pezo L, Brahusi F et al. Chemical composition and amino acid content in different genotypes of wheat flour. Period Polytech Chem Eng. 2019;63(4):618-28. DOI: 10.3311/PPch.13185
9. Alebiosu MO, Akinbode BA, Oni IS, Oladele JO. Quality evaluation of cookies produced from wheat, sorghum and defatted coconut flour blends. Asian Food Sci J. 2020;17(3):11-21. DOI: 10.9734/AFSJ/2020/v17i330192
10. Ukeyima MT, Dendegh TA, Isusu SE. Quality characteristics of bread produced from wheat and white kidney bean composite flour. Eur J Nutr Food Saf. 2019;10(4):263-72. DOI: 10.9734/EJNFS/2019/v10i430120
11. Nwagbaoso O, Okoronkwo KA, Awah AI. Investigation into the functional and sensory properties of two varieties of local black beans (*Phaseolus vulgaris*) flour. IOSR J Environ Sci Toxicol Food Technol. 2018;12(7):39-45. DOI: 10.9790/2402-1207023945
12. Okoye JI, Nkwocha AC, Agbo AO. 'Chemical composition and functional properties of kidney bean/wheat flour blends,' Continental J. Food Science and Technology. 2008;2:27-32.
13. Trinidad TP, Mallillin AC, Valdez DH, Loyola AS, Askali-Mercado FC, Castillo JC et al. Dietary fiber from coconut flour: A functional food. Innov Food Sci Emerg Technol. 2006;7(4):309-17. DOI: 10.1016/j.ifset.2004.04.003
14. Ramaswamy L. Coconut flour: A low carbohydrate, gluten free flour. Int J Ayurvedic Herb Med. 2014:1426-36.
15. Raczyk M, Kruszewski B, Michałowska D. Effect of coconut and chestnut flour supplementations on texture, nutritional and sensory properties of baked wheat based bread. Molecules. 2021;26(15). DOI: 10.3390/molecules26154641, PMID 34361794
16. Afoakwa NA, Owusu J, Owusu V. Characteristics of coconut (*Cocos nucifera*) flour and its application in cake. Asian Food Sci J. 2019;13(1):1-11. DOI: 10.9734/afsj/2019/v13i130095
17. Adelekan AO, Alamu T. Effect of coconut (*Cocos nucifera*) flakes substitution on some quality parameters of wheat bread, african Journal of food. Agric Nutr Dev. 2021;21(6):18349-67. DOI:10.18697/ajfand.102.20170.
18. Chandra S, Samsher. Assessment of functional properties of different flours. Afr J Agric Res. 2013;8(38):4849-52. DOI: 10.5897/AJAR2013.6905
19. Mepba HD, Eboh L, Nwaojigwa SU. Chemica composition, functional and baking properties of; wheat-plantain composite flours, african Journal of food. Agric Nutr Dev. 2007;7(1).

20. Ojo MO, Ariahu CC, Chinma EC. Proximate, functional and pasting properties of cassava starch and mushroom (*Pleurotus pulmonarius*) flour blends. *Am J Food Sci Technol*. 2017;5(1):11-8. DOI: 10.12691/ajfst-5-1-3.
21. Makinde FM, Oluwaseun M. Influence of margarine and/or coconut oil on quality of bread prepared from wheat flour. *Asian J Plant Sci Res*. 2017;7(1):23-30.
22. Marolt G, Kolar M. Analytical methods for determination of phytic acid and other inositol phosphates: a review. *J Moeculesl*. 2021;26(174). DOI: 10.3390/molecules26010174
23. Adeniyi SA, Orjiekwe CL, Ehiagbonare JE. Determination of alkaloids and oxalates in some selected food samples in Nigeria. *Afr J Biotechnol*. 2009;8(1):110-2.
24. Nwokenkwo EC, Nwosu JN, Onuegbu NC, Olawuni IA, Ofoedum AF. Evaluation of the Antinutrients, amino acid profile and Physicochemical Properties of Hura crepitans Seed. *Arch Curr Res Int*. 2020;20(5):1-17. DOI: 10.9734/ACRI/2020/v20i530192
25. Hashim IB, Khalil AH, Afifi HS. Quality characteristics and consumer acceptance of yoghurt fortified with dietary fiber. *J Dairy Sci*. 2009;92(11):5403-7. DOI: 10.3168/jds.2009-2234, PMID 19841201
26. Kumar K, Chandra S, Chauhan N, Singh J, Kumar M. Functional properties of food commodities (wheat, kidney bean, cowpea, turnip, cauliflower) flours. *Int J Chem Stud*. 2017;5(6):1199-202.
27. Kaushal P, Kumar V, Sharma HK. Comparative study of physicochemical, functional, anti-nutritional and pasting properties of taro (*Colocasia esculenta*), rice (*Oryza sativa*), pigeon pea (*Cajanus cajan*) flour and their blends. *J Food Sci Technol*. 2012;48:59-68.
28. Awuchi CG, Igwe VS, Echeta CK. The functional properties of foods and flours. *Int J Adv Acad Res*. 2019;5(11): 139-60.
29. Chandra S, Singh S, Kumari D. Evaluation of functional properties of composite flours and sensorial attributes of composite flour biscuits. *J Food Sci Technol*. 2015;52(6):3681-8. DOI: 10.1007/s13197-014-1427-2, PMID 26028751.
30. MH, MN, HN, Zainol MK, Jahurul MH. Functional properties of composite flour: a review. *Food Res*. 2020;4(6):1820-31. DOI: 10.26656/fr.2017.4(6).419
31. Moses MO, Olanrewaju MJ. Evaluation of functional and pasting properties of different corn starch flours. *Int J Food Sci Nutr*. 2018;3(26):95-9.
32. Adejuyitan JA, Otunola ET, Adesola MO, Onalapo OE. Production and quality evaluation of short bread biscuit using wheat and fermented unripe plantain flour. *Eur J Nutr Food Saf*. 2020;12(4):30-42. DOI: 10.9734/ejnfs/2020/v12i430216
33. Abdelghafor AMH, Mustafa RF, Ibrahim AI, Krishnan PG. Quality of bread from composite flour of sorghum and hard white winter wheat. *Adv J Food Sci Technol*. 2011;3:9-15.
34. Hyacinthe AA, Bedel FJ, Gisèle KA, Hermann A, Atta F, Patrice KL. Functional properties of composite flours produced with Ivorian taro (*Colocasia esculenta* L. Cv Fouê) corms flour and wheat (*Triticum aestivum* L.) flour, GSC Biological and Pharmaceutical Sciences. 2021;15(03): 164–176.
35. Pathirana HPDTH, Lakdusinghe WMK, Yalagama LLWC, Chandrapeli CATD, Madusanka JAD. Evaluation of nutritional composition of defatted coconut flour incorporated biscuits. *Cord*. 2020;36:33-9. DOI: 10.37833/cord.v36i.427
36. Stoin D, Petrovich LI, Velciov AB, Trasca T, Ravis A, Jianu C. Red kidney bean and rice flours: potential ingredients in the production of gluten-free bread with functional quality. *J Agroalimnt Processes Technol*. 2019;25(3):147-52.
37. Mihiranie S, Jayasundera M, Perera N. Development of snack crackers incorporated with defatted coconut flour. *J Microbiol Biotechnol Food Sci*. 2017;7(2):153-9. DOI: 10.15414/jmbfs.2017.7.2.153-159
38. Manonmani D, Soumya BS, Bosco JD. Effect of red kidney bean (*Phaseolus vulgaris* L.) flour on bread quality. *Open Access Libr J*. 2014;01:1-6. DOI: 10.4236/oalib.1100366
39. Nirmala NPV, Joye IJ. Dietary fibre from whole grains and their benefits on metabolic health. *J Nutrients*. 2020; 12(3045). DOI: 10.3390/nu12103045, PMID 33027944

40. Kambabazi MR, Okoth MW, Ngala S, Njue L, Vasanthakaalam H. Evaluation of nutrient content in red kidney beans, amaranth leaves, sweet potato roots and carrots cultivated in Rwanda. *AJFAND*. 2021;21(4):17801-14.
DOI: 10.18697/ajfand.99.21095
41. Akinjayeju O, Ijarotimi OS, Awolu OO, Fagbemi TN. Nutritional composition, glycaemic properties and antidiabetic potentials of cereal-based soy-fortified flours for functional dough meal in diabetic induced rats. *J Food Sci Nutr Research*. 2020;3(2):102-20.
DOI: 10.26502/jfsnr.2642-11000042.
42. Rasha MK, Esmat AAA, Gibriel AY, Nagwa MHR, Ferial MAS. Effect of legume processing treatments individually or in combination on their phytic acid content. *Afr J Food Sci Technol*. 2011;2(2):36-46.
43. Ohini OP, Ferdinand OM, Betsy OO. Chemical, anti-nutritional factors and sensory properties of maize-kidney bean flours. *Food Sci Nutr Technol*. 2019;4(6):1-9.
DOI: 10.23880/fsnt-16000203
44. Ayele HH, Bultosa G, Abera T, Astatkie T. Nutritional and sensory quality of wheat bread supplemented with cassava and soybean flours. *Cogent Food Agric*. 2017;3(1):1-13.
DOI: 10.1080/23311932.2017.1331892
45. Samtiya M, Aluko RE, Dhewa T. Plant food anti-nutritional factors and their reduction strategies: an overview. *Food Prod Process Nutr*. 2020;2(1):1-14.
DOI: 10.1186/s43014-020-0020-5
46. Mamiro PS, Mwanri AW, Mongi RJ, Chivaghula TJ, Nyagaya M, Ntwenya J. Effect of cooking on tannin and phytate content in different bean (*Phaseolus vulgaris*) varieties grown in Tanzania. *Afr J Biotechnol*. 2017;16(20):1186-91.
DOI: 10.5897/AJB2016.15657
47. Pandit M, Kaur N. Physico-chemical characteristics and anti-nutritional factors of wheat, soybean, oats and pumpkin leaves. *Chem Sci Rev Lett*. 2020;9(34):260-7.
DOI: 10.37273/chesci.CS20510126
48. Mohsen SM, Fadel HHM, Bekhit MA, Edris AE, Ahmed MYS. Effect of substitution of soy protein isolate on aroma volatiles, chemical composition and sensory quality of wheat cookies. *Int J Food Sci Technol*. 2009;44(9):1705-12.
DOI: 10.1111/j.1365-2621.2009.01978.x
49. Serrem C, Kock H, Taylor J. Nutritional quality, sensory quality and consumer acceptability of sorghum and bread wheat biscuits fortified with defatted soy flour. *International J Food Sci Technol*. 2011;46:74-83.
50. Nwanekezi EC. composite Flours for Baked Products and Possible Challenges – a review. *Off J Niger Inst Food Sci Technonology*. 2013;31(2):8-17.
DOI: 10.1016/S0189-7241(15)30071-0.
51. Hassan E, M. Awad Al A, Moniem I. A. Effect of Fermentation and particle size of wheat bran on the antinutritional factors and bread quality. *Pak J Nutr*. 2008;7(4):521-6.
DOI: 10.3923/pjn.2008.521.526.

© 2023 Forwoukeh et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/95440>