



Incidence of Wilt Disease on Fenugreek Caused by *Fusarium oxysporum* Schlecht in Rajasthan, India

Vikash Kumar ^{a*}, A. L. Yadav ^a, Rakesh Kumar ^a and Anuj ^a

^a Department of Plant Pathology, College of Agriculture, SKRAU, Bikaner -334006, Rajasthan, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JEAI/2023/v45i122282

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/110641>

Original Research Article

Received: 14/10/2023

Accepted: 19/12/2023

Published: 30/12/2023

ABSTRACT

Fenugreek is an annual legume crop with small seeds and self-pollinated plant, belonging to the family *Fabaceae*. It ranks as the third largest seed spice in India following coriander and cumin. The crop suffers from several diseases, among them wilt caused by *Fusarium oxysporum* is one of the major economically important diseases in many fenugreek growing states of India including Rajasthan. The survey for fenugreek wilt disease during *Rabi* (winter) season 2021-22 and 2022-23 revealed that wilt disease caused by *Fusarium oxysporum* is an imperative pathological problem. Average per cent disease incidence was 24.03 per cent recorded in surveyed districts of Rajasthan. Highest average disease incidence (29.31%) was reported in Nagaur followed by Sikar (24.85%), while lowest in Bikaner (17.93%) district.

Keywords: Disease; fenugreek; *Fusarium oxysporum*; Rajasthan; wilt.

*Corresponding author: E-mail: vikashsihag029@gmail.com;

1. INTRODUCTION

Fenugreek commonly known as '*Methi*', is a self-pollinated, small seeded, annual legume crop which belongs to the family *Fabaceae*. It is an annual forage legume and spice crop and is widely known for its medicinal, pharmaceutical and nutraceutical properties [1]. Fenugreek is native to an area extending from Iran to northern India [2], but it is now grown as a spice in most parts of the world. Fenugreek seed is used as a spice, is one of the main ingredients in curry powder [3] and also has a long history of use as a medicinal herb, being used extensively in both Indian ayurvedic and traditional chinese medicine [4]. It is regarded as the oldest known medicinal plant in recorded history [5].

Fenugreek is an important multipurpose winter season seed spice crop. It ranks as the third largest seed spice in India following coriander and cumin. In India, the major fenugreek growing states are Rajasthan, Madhya Pradesh, Gujarat, Haryana, Uttarakhand and West Bengal. In India, it is cultivated on approximately 168716 hectare of land with an annual production of about 252063 tonnes and a productivity of 1.49 tonnes ha⁻¹. Rajasthan leads in both area and production of fenugreek in India with 90469 hectare area under cultivation, an annual production of 110869 tonnes and a productivity of 1.23 tonnes ha⁻¹ [6]. In Rajasthan, the major fenugreek growing districts are Bikaner, Jodhpur, Pratapgarh, Nagaur, Churu and Sikar etc. Bikaner takes the first position in terms of area and production of fenugreek in Rajasthan with 25533 ha area and 27372 tonnes production along with 1.07 tonnes ha⁻¹ productivity [7].

The production and productivity of fenugreek is highly affected by fungal diseases such as root rot, powdery mildew, downy mildew, wilt, rust, leaf spot and charcoal rot diseases [8]. Among them, wilt caused by *Fusarium oxysporum* Schlecht is one of the major economically important diseases in many fenugreek growing states of India including Rajasthan. The disease was first reported from Rajasthan by Shivpuri and Bansal [9]. *Fusarium* wilt of fenugreek can lead to annual losses up to 56 per cent [10]. Wilt of fenugreek, exhibited various types of symptoms at any stage of the crop. Infections at the seedling stage result in complete drying, while infected plants exhibit various types of symptoms such as stunting of the plants, gradual yellowing of lower leaves, drooping, epinasty, drying of leaves and in some cases partial

wilting. Dried leaves remain attached to the wilted plants. When wilted plants split longitudinally, they reveal brown vascular discoloration. Ultimately, wilted plants lose vigor, leading to the death of the entire plant [11]. A proper investigation needs a perfect scrutiny. So, the survey was carried out to monitor the disease incidence in major fenugreek growing districts of Rajasthan.

2. MATERIALS AND METHODS

Survey for the occurrence of wilt of fenugreek was conducted during *Rabi* (winter) 2021-22 & 2022-23 to assess the prevalence and incidence in major fenugreek growing areas of Rajasthan. Survey was conducted in Bikaner, Sikar and Nagour districts of Rajasthan. Three tehsils (Blocks) under each district and four (4) villages under each tehsil were included. Under each village four (4) farmer's fields were assessed. In each field, five (5) spots of 1 m² area were marked diagonally at randomly to cover entire field. Diseased and healthy plants were counted in each spot and the per cent disease incidence was calculated by using the following formula [12].

$$\text{Percent Disease Incidence} = \left(\frac{\text{Number of wilted plants}}{\text{Total number of observed plants}} \right) \times 100$$

The mean per cent of disease incidence was calculated using Excel.

3. RESULTS AND DISCUSSION

Survey was conducted in major fenugreek growing districts of Rajasthan during *Rabi* (winter) season 2021-22 and 2022-23 which showed the wilt disease caused by *Fusarium oxysporum* is an imperative pathological problem. Average per cent disease incidence was 24.03 per cent recorded in surveyed districts of Rajasthan. Highest average disease incidence (29.31%) was reported in Nagaur followed by Sikar (24.85%) while lowest in Bikaner (17.93%) district (Table, Fig. & Plate 1).

The survey in Bikaner district revealed that the average disease incidence ranged from 15.13 to 21.57 per cent during the *Rabi* seasons of 2021-22 and 2022-23. The maximum per cent disease incidence was observed in fenugreek fields at Punrasar village of Shri Dungargarh *tehsil* that is upto 21.57 per cent and minimum per cent disease incidence (15.13%) in Parwa village of

Nokha *tehsil*. In the Sikar district, the average disease incidence was recorded from 22.86 to 26.23 per cent. The highest per cent disease incidence was observed at Gokulpura (26.23%) village of Sikar *tehsil* and minimum per cent disease incidence in Kayamsar (22.86%) village of Fatehpur *tehsil*. The survey conducted in Nagaur district demonstrated an average disease incidence ranging from 27.30% to 31.32%. The

highest per cent disease incidence was observed at Kuchera (31.32%) village of Nagaur *tehsil* and minimum per cent disease incidence in Dangavas (22.86%) village of Merta *tehsil*. The elevated disease incidence could be attributed to factors such as the previous crop, monocropping system and the complex nature of the pathogen, which may have exacerbated the disease situation [10]. Similar finding was reported by

Table 1. Per cent disease incidence of fenugreek wilt disease in major growing districts of Rajasthan

Districts	Tehsils	Villages	Disease incidence (%)*		Mean PDI
			2021-22	2022-23	
Bikaner	Bikaner	Deshnok	15.25	18.25	15.25
		Himtasar	17.63	20.28	17.63
		Barsingsar	19.85	17.55	19.85
	Nokha	Lalamdesar	15.69	17.86	15.69
		Rasisar	18.46	16.74	18.46
		Parwa	15.13	18.00	15.13
		Bhamatsar	17.65	19.67	17.65
		Manyana	17.55	20.10	17.55
		Sheruna	19.17	17.45	19.17
	Shri Dungargarh	Sanwatsar	19.92	21.12	19.92
		Ridi	17.32	18.38	17.32
		Punrasar	21.57	20.65	21.57
		Average		17.93	18.84
Sikar	Sikar	Gokulpura	27.67	24.78	26.23
		Shyampura	24.43	27.65	26.04
		Bajor	27.45	23.68	25.57
	Lachhmangarh	Radha Kishanpura	25.64	26.75	26.20
		Patoda	22.78	24.10	23.44
		Ghassu	26.25	24.00	25.13
		Dantujala	23.89	25.93	24.91
		Hameerpura	26.12	24.55	25.34
	Fatehpur	Ramgarh	25.36	22.85	24.11
		Bhojdesar	21.85	24.25	23.05
		Hetamsar	26.65	24.14	25.40
		Kayamsar	22.49	23.23	22.86
	Average		25.05	24.66	24.85
Nagaur	Nagaur	Kuchera	32.40	30.23	31.32
		Mundwa	29.32	31.17	30.25
		Bhadana	30.70	28.20	29.45
		Gothra	27.89	29.00	28.45
	Merta	Dangavas	28.24	26.35	27.30
		Netadiya	31.65	28.60	30.13
		Dholerao Khurd	29.56	30.45	30.01
		Raliyawata	31.20	27.32	29.26
		Langod	26.90	28.12	27.51
	Degana	Rewat	29.76	28.20	28.98
		Tamroli	28.50	30.37	29.44
		Jalsoo Khurd	30.43	28.87	29.65
		Average		29.71	28.91
Total Average		24.23	24.13	24.03	

*Average of four fields in each village

Singh et al. [13], who assessed the pathogenic disease incidence of foot rot of fenugreek caused by *Fusarium moniliforme* and recorded

occurrence of disease incidence ranged from 44-58 per cent in Rohilkhand region of Uttar Pradesh.




Fig. 1. chart shows Per cent disease incidence of fenugreek wilt disease in major growing districts of Rajasthan



Plate 1. Survey of wilt disease of fenugreek in major growing districts of Rajasthan

Our findings are also in accordance with the earlier reports of Rani and Hegde [14], who conducted survey for wilt of fenugreek in three districts viz., Dharwad, Belagavi and Haveri. Highest per cent disease incidence was recorded in Dharwad (1.00 to 56.19%) followed by Haveri (11.31% to 42.59%) and least disease incidence was observed in Belagavi (9.71% to 29.22%). Similarly, Bhimani et al. [15] isolated *Fusarium oxysporum* schlecht. in fenugreek from fifteen different locations of Saurashtra regions of Gujarat. Among them, one isolate (Khadpipali) found highly pathogenic with 93.33 per cent disease incidence [16].

4. CONCLUSION

A proper investigation needs a perfect scrutiny. So, the survey was carried out to monitor the disease incidence in major fenugreek growing districts of Rajasthan. The survey revealed that wilt disease caused by *Fusarium oxysporum* is an imperative pathological problem. Average disease incidence was 24.03 per cent recorded in surveyed districts of Rajasthan. Highest average disease incidence (29.31%) was reported in Nagaur followed by Sikar (24.85%), while lowest in Bikaner (17.93%) district.

ACKNOWLEDGEMENT

The authors are very much thankful to Department of Plant Pathology, College of Agriculture, Swami Keshwanand Rajasthan Agricultural University, Bikaner for providing facilities.

COMPETING INTERSTS

Authors have declared that no competing interests exist.

REFERENCES

1. Bawadi HA, Maghaydah SN, Tayyem RF. The postprandial hypoglycemic activity of fenugreek seed and seeds extract in type 2 diabetics: A pilot study. *Pharma. Mag.* 2009;4(18):134-138.
2. Acharya SN, Srichamroen A, Basu S, Ooraiku B, Basu T. Improvement in the nutraceutical properties of fenugreek (*Trigonella foenum-graecum* L.). *J. Sci. Tech.* 2006;28(1):1-9.
3. Srinivasan K. Fenugreek (*Trigonella foenum-graecum*): A review of health beneficial physiological effects. *Food Rev. Int.* 2006;22:203-224.
4. Tiran D. The use of fenugreek for breast feeding woman. *Comp. Ther. Nurs. Midwifery.* 2003;9(3):155-156.
5. Lust, J. B. (1986) *The herb book*. Bantam books Inc. New York.1-55.
6. Anonymous. Directorate of Arecanut and Spices Development, Kozhikode, Kerala, Govt. of India; 2021-22a.
7. Anonymous. Rajasthan Agricultural Statistics at a glance, Department of Agriculture, Govt. of Rajasthan; 2021-22b.
8. Prasad R, Acharya S, Erickson S, Thomas J. Identification of cercospora leaf spot resistance among fenugreek accessions and characterization of the pathogen. *Australian J. Crop. Sci.* 2014; 8(6):822-830.
9. Shivpuri A, Bansal PK. *Fusarium wilt of Trigonella foenum-graecum* L. *Indian J. Mycolo. Pl. Pathol.* 1987;26:749-751.
10. Rani N. Studies on root rot/wilt of fenugreek with special reference to *Fusarium oxysporum* Schlecht. Doctoral dissertation, University of Agricultural Sciences Dharwad; 2015.
11. Rani N, Hegde YR, Nargund VB. Symptomatology, proving pathogenicity and standardization of inoculum density of *Fusarium oxysporum* causing wilt of fenugreek. *J. Pharma. Phytochem.* 2017a; 6(6):505-509.
12. Jangir H, Ram D. and Choudhary A. Management of wilt disease of cumin caused by *F. oxysporum f. sp. cumini* with fungicides in vitro and in vivo. *The J. Res. Pjtsau.* 2022;50(1):101-107.
13. Singh CP, Mishra US, Pandey UC. Occurrence and incidence of foot rot disease on fenugreek caused by *Fusarium moniliforme* in Rohilkhand region of Uttar Pradesh, India. *J. Environ. Sci. Technol.* 2014;2(2): 5-28.
14. Rani N, Hagde YR. Survey for the incidence of root rot/wilt of fenugreek in Northern Karnataka, India. *Int. J. Curr. Microbiol. App. Sci.* 2017;6(5):1564-1569.
15. Bhimani MD, Akbari LF, Golakiya BA. Characterization of Indian isolates of *Fusarium oxysporum* Schlecht causing fenugreek wilt. *Int. J. Chem. Studies.* 2018a;6(2):1167-1172.

16. Mary J. Fenugreek seed benefits the digestive system and lowers blood pressure; 2009. Available:<http://EzineArticles.com>

© 2023 Kumar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/110641>