



Growing Degree Days Requirement and Yield of Wheat Cultivars as Influenced by Irrigation Scheduling and Time of Sowing

**Arjun Lal Prajapat ^{a*}, Rani Saxena ^a, Mahesh Sharma ^a,
Rameshwr Lal Mandeewal ^a, Babu Lal ^a and Bhuwanesh Didal ^a**

^a Sri Karan Narendra Agriculture University, Jobner, Rajasthan-303329, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i330843

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/83009>

Original Research Article

Received 05 December 2021
Accepted 10 February 2022
Published 14 February 2022

ABSTRACT

The crop production characteristics are directly related with the temperature and relative humidity of atmosphere during crop growing season. Therefore, the main aim of this study was to highlights thermal requirement (GDD) of wheat varieties for sustainable yield. The field experiment were conducted during *Rabi* season 2016 and 2017 at Research farm, Rajasthan Agricultural Research Institute, Sri Karan Narendra Agriculture University, Durgapura, Jobner, Rajasthan to find out the response of wheat cultivars to irrigation scheduling under different sowing dates. Thirty six treatment combinations were investigated. Treatments comprises four irrigation levels: I₁ (0.6 ETc), I₂ (0.8 ETc), I₃ (1.0 ETc) and I₄ (1.2 ETc), three cultivars: C₁ (Raj-4120), C₂ (Raj-4079) and C₃ (Raj-4238) and three dates of sowing: D₁ (15th Nov.), D₂ (30th Nov.) and D₃ (15th Dec.) in Split plot design. The pooled analysis of data revealed that irrigation scheduling treatment I₄ (1.2 ETc) recorded significantly higher value of yield and Growing degree days (GDD), By cultivar Raj 4079 observed the higher value of yield and GDD. Further, results showed that the wheat crop sown on 15th November obtained the maximum yield and growing degree days value with phenological stages. The number of days required to attain different phenological stages decreased with late sowing condition. This indicated that irrigation scheduling at 1.2 ETc should adopted in cultivar Raj 4079 at the sowing on 15th November to achieve maximum yield under hot climate of Rajasthan.

*Corresponding author: E-mail: prajapatasu@gmail.com;

Keywords: Growing degree days; wheat and yield.

1. INTRODUCTION

Wheat is a long-day crop that is sensitive to temperature. The temperature has a significant impact on its growth and productivity. Wheat is sown late, exposing pre-anthesis phenological events to high temperatures, which affect grain development and, eventually, yield [1]. The buildup of heat or temperature units above the threshold or base temperature is related to phenological development from sowing to maturity (below which no growth occurs). To reach a specific phenophase, a quantifiable value of heat or temperature units is necessary. Various meteorological elements influence plant physiological and morphological development, such as temperature, which is an essential environmental factor regulating crop growth, development, and yield. During growth and development of a cereal crop several growth stages are distinguishable in which important physiological processes occur [2]. Plants have a definite temperature requirement before they attain certain phenological stages. Several research findings noticed that temperature below (<10 °C) or above (>25 °C) the optimum (12-25 °C) alter phenology, growth and development and finally reduce the yield of wheat cultivars [3]. Influence of temperature on phenology and yield of crop plants can be studied under field condition through accumulated heat units system [4]. Temperature stress intensity is severe under late sowing, causing reduction in the duration of later growth phases resulted in acquisition of less days to mature. Air temperature based agrometeorological indices viz., growing degree days (GDD), has been used to describe changes in phenological behavior and growth parameters [5]. The values of accumulated GDD for each phenophases are relatively constant and independent of sowing date but vary in a crop from cultivar to cultivar [6]. Agronomic application of temperature effect on plant is the concept of heat unit or growing degree days and heat stress in wheat can be mitigated in two ways: heat management or the development of heat-tolerant cultivars. Evaluation of high yielding, stable genotypes having good quality are considered pre-requisite for increasing crop production in any region. So, there is a need to broader genetic base and to replace old cultivars with new and improved ones [7]. Recommendation of several varieties will also help farmers to select best one and also adequate supply of improved varieties to farmers. Especially in wheat choice of

varieties depends on the time of higher, unfavorable air temperature, where difference in crop agronomy lead to difference in sowing dates, longer duration cultivars for earlier sowing and shorter duration cultivars for late sowing are recommended. So this experiment was conducted to study the effect of irrigation scheduling and dates of sowing on the phenology, heat unit requirement and heat use efficiency of wheat cultivars.

2. MATERIALS AND METHODS

2.1 Experimental Site

The experiments were carried out during *Rabi* seasons 2016-17 and 2017-18 at Research farm, Rajasthan Agricultural Research Institute, Sri Karan Narendra Agriculture University, Jobner, Rajasthan. The experimental site's geographically situated at 75° 47' East longitudes, 26° 51' North latitude and at altitude of 390 m above mean sea level. The region's climate is classified as semi-arid eastern plain zone of Rajasthan with characterized by aridity of the atmosphere and extremity of temperature both in summer (45.5°C) and winter (4°C) with annual rainfall of 500-700 mm. The weather data for the entire cropping season observed from the RARI, SKNAU, Jobner meteorological observatory. The soil of experimental field was loamy sand in texture, slightly alkaline in reaction. Soil samples from 0-30 and 30-60 cm deep layers were collected from the experimental field prior to experimentation using core sampler and later, the analysis for various physico-chemical properties was carried out. The details of the initial soil properties of the experimental sites are presented in Table 1.

2.2 Experimentation and Crop Husbandry

The experiments were laid out in Split plot design with three replications added thirty-six treatment combinations. Treatments comprises four irrigations viz., I1 (0.6 ETc), I2 (0.8 ETc), I3 (1.0 ETc) and I4 (1.2 ETc), three cultivars: C1 (Raj-4120), C2 (Raj-4079) and C3 (Raj-4238) and three dates of sowing: D1 (15th Nov.), D2 (30th Nov.) and D3 (15th Dec.). In the recommended irrigation treatments applied at different irrigation intervals according to ETC level with the help of water meter. The crop was raised by following all agronomic and plant protection measures except irrigation schedule.

2.3 Measurement and Sampling

Observations on number of days taken to various phenological stages *viz.*, germination, CRI, tillering, jointing, flag leaf and maturity stage were in one meter row length. The readings on various phenophases were recorded once the plants have reached 50% of that particular stage in each replication. Crop was harvested manually in the end week of March and first week of April when 80% of the grains turned to golden colour. Grain and biological yield were recorded at the harvest. Thus, the data recorded during the course of investigation were subjected to statistical analysis.

2.4 Growing Degree Days (GDD)

Growing degree days were calculated by simple arithmetic accumulation of daily mean temperature above the base temperature value of 5°C considered for the wheat crop. The different indices for each stage were calculated as suggested by [8].

$$\text{Growing degree days (}^{\circ}\text{C)} = \frac{T_{\max} + T_{\min}}{2} - T_b$$

where,

T_{\max} = Daily maximum temperature ($^{\circ}\text{C}$)

T_{\min} = Daily minimum temperature ($^{\circ}\text{C}$)

T_b = Base temperature (5°C)

3. RESULTS AND DISCUSSION

3.1 Crop Phenology

Data pertaining to crop phenology and from emergence to maturity duration presented in Table 2. Irrigation did not have differential effect on duration from seeding to tillering because of absence of variation in moisture availability and temperature up to tillering stage. However, onset and duration of phenophases beyond maximum tillering increased with increase variation in moisture availability, possibly owing to retention of more physiologically active leaf area for a longer period [9]. Increasing water availability required more heat units to shift from one phenophases to another. The crop raised with 1.2 ET_c irrigation scheduling required more days to attain various phenophases and heat units. The three wheat cultivars showed a significant difference in number of days taken to different phenophases. Cultivar Raj 4079 than other cultivars more days taken to complete different

phenophases. This is obvious because of genetic behavior of each cultivar of a crop had specific requirement of heat units for initiation and completion of its phenological stages under particular environmental conditions. Similar results were also reported by Dixit et al. (2014). Further, results revealed that different phenological stage (Emergence, Crown root initiation, Tillering, Jointing, Flage leaf, Maturity) were significantly influenced by dates of sowing in wheat crop. The crop sown on November 15 took significantly higher number of days (123) to attain maturity as compared to 30 November (117) and 15 December (106) sown crops. The phenological studies revealed that delayed sowing on 15 December required significantly less number of days for initiation and completion of the different phenophases in comparison to the sowing on 15 November and 30 November. Conversely under late sowing, reproductive growth as well as total crop duration was drastically reduced. It is an established fact that the crop phenology is largely dependent on genetic and environmental factor *viz.*, temperature, solar radiation, rainfall *etc* [10].

3.2 Growing Degree Day (GDD)

The results were revealed that the agrometeorological index *i.e.* growing degree day (GDD) from emergence to tillering stage not influenced significantly. However, onset and duration of phenophases beyond maximum tillering increased with increase variation in moisture availability, possibly owing to retention of more physiologically active leaf area for a longer period [9]. Jointing to maturity accounted higher GDD with I₄ (1.2 ET_c) than other irrigation level while lowest GDD observed when crop irrigated with I₁ (0.6 ET_c) and I₂ (0.8 ET_c) irrigation level. At maturity of the crop highest GDD (1797.59°C day) was accounted with I₄ irrigation level than I₃ (1780.97°C day) and I₂ while lowest GDD were recorded in case of I₁ (1688.84°C day) irrigation level (Table 3.). At maturity stage, higher GDD was recorded in case of cultivar Raj 4079 followed by cultivar Raj 4238. In pooled analysis maximum GDD (1730.73 °C day) was recorded for Raj 4079 followed by Raj 4238 (1720.32 °C day) (Table 3.). GDD requirement was maximum in November 15 (1772.85 °C day) sowing among the sowing dates and decreased with delay in sowing followed by November 30 (1713.27 °C day) and December 15 (1633.57 °C day) at maturity stage. The requirement of heat unit was higher for

Table 1. Initial physico-chemical properties of experimental site

| S.No. | Parameters | Value | | Methods of Analysis Employed |
|-----------|------------------------------|---------|---------|---|
| A. | Mechanical Properties | 2016-17 | 2017-18 | |
| i. | Coarse Sand % | 27.5 | 25.5 | Bouyoucous Hydrometer Method |
| ii. | Fine Sand % | 53.8 | 51.3 | |
| iii. | Silt % | 10.7 | 11.0 | |
| iv. | Clay % | 7.8 | 8.1 | |
| | Textural class | | | |
| B. | Chemical Properties | | | |
| i. | pH | 8.3 | 8.1 | 1:2.5 Soil water suspension measured with Glass electrode pH |
| ii. | EC (ds/m) | 0.13 | 0.18 | 1:2.5 Soil water suspension measured with Systronics conductivity meter |
| iii. | Organic Carbon (%) | 0.27 | 0.24 | Walkley and Black method |
| iv. | Available N (kg/ha) | 134.2 | 139.2 | Alkaline potassium permanganate method |
| v. | Available P (kg/ha) | 12.6 | 12.8 | Sodium bicarbonate method |
| vi. | Available K (kg/ha) | 134.9 | 136.4 | Ammonium acetate method |

Table 2. Effect of irrigation scheduling, varieties and dates of sowing on phenology of wheat

| Treatments | Emergence | CRI | Tillering | Jointing | Flag leaf | Maturity |
|--|-----------|-----|-----------|----------|-----------|----------|
| Irrigation scheduling | | | | | | |
| I ₁ (Etc 0.6) | 7 | 22 | 41 | 56 | 67 | 108 |
| I ₂ (Etc 0.8) | 7 | 22 | 41 | 63 | 74 | 115 |
| I ₃ (Etc 1.0) | 7 | 22 | 42 | 66 | 78 | 119 |
| I ₄ (Etc 1.2) | 7 | 22 | 42 | 67 | 79 | 121 |
| Cultivars | | | | | | |
| V ₁ (Raj 4120) | 7 | 21 | 40 | 61 | 73 | 114 |
| V ₂ (Raj 4079) | 7 | 24 | 42 | 64 | 77 | 118 |
| V ₃ (Raj 4238) | 7 | 23 | 41 | 63 | 75 | 117 |
| Date of sowing | | | | | | |
| D ₁ (15 th NOV.) | 6 | 24 | 44 | 68 | 80 | 123 |
| D ₂ (30 th NOV.) | 7 | 23 | 41 | 64 | 75 | 117 |
| D ₃ (15 th DEC.) | 8 | 21 | 38 | 58 | 68 | 106 |

Tables 3. Effect of irrigation scheduling, cultivars and varying sowing dates on growing degree days ($^{\circ}\text{C}$ days) at different phenological stages of wheat

| Treatment | GDD at emergence stage | GDD at CRI stage | GDD at tillering stage | GDD at jointing stage | GDD at flag leaf stage | GDD at maturity stage |
|--|------------------------|------------------|------------------------|-----------------------|------------------------|-----------------------|
| Irrigation scheduling | | | | | | |
| I ₁ (Etc 0.6) | 91.57 | 313.00 | 530.40 | 708.70 | 835.80 | 1558.88 |
| I ₂ (Etc 0.8) | 91.81 | 314.17 | 530.90 | 799.98 | 941.96 | 1688.84 |
| I ₃ (Etc 1.0) | 92.19 | 315.69 | 533.30 | 832.22 | 999.06 | 1780.97 |
| I ₄ (Etc 1.2) | 92.37 | 316.89 | 535.90 | 839.96 | 1012.12 | 1797.59 |
| SEm \pm | 0.33 | 1.34 | 2.91 | 3.68 | 4.99 | 10.57 |
| CD(P= 0.05) | NS | NS | NS | 11.33 | 15.37 | 32.58 |
| Cultivars | | | | | | |
| V ₁ (Raj 4120) | 91.70 | 307.99 | 521.70 | 774.41 | 910.13 | 1668.64 |
| V ₂ (Raj 4079) | 92.21 | 319.18 | 540.80 | 807.87 | 969.52 | 1730.73 |
| V ₃ (Raj 4238) | 92.04 | 317.63 | 535.50 | 803.36 | 962.04 | 1720.32 |
| SEm \pm | 0.20 | 0.93 | 1.91 | 2.27 | 2.73 | 7.72 |
| CD(P= 0.05) | NS | 2.68 | 5.51 | 6.54 | 7.86 | 22.23 |
| Date of sowing | | | | | | |
| D ₁ (15 th NOV.) | 97.77 | 331.07 | 568.80 | 899.53 | 1053.88 | 1772.85 |
| D ₂ (30 th NOV.) | 95.36 | 315.64 | 531.90 | 795.54 | 942.29 | 1713.27 |
| D ₃ (15 th DEC.) | 82.82 | 298.10 | 497.20 | 690.57 | 845.53 | 1633.57 |
| SEm \pm | 0.15 | 0.74 | 1.27 | 1.40 | 1.97 | 6.41 |
| CD(P= 0.05) | 0.43 | 2.08 | 3.57 | 3.92 | 5.54 | 17.99 |

Table 4. Effect of irrigation scheduling, varieties and dates of sowing on Yield and Heat Use Efficiency of wheat

| Treatment | Grain yield | Biological yield |
|--|-------------|------------------|
| Irrigation scheduling | | |
| I ₁ (Etc 0.6) | 3442 | 7445 |
| I ₂ (Etc 0.8) | 4600 | 10130 |
| I ₃ (Etc 1.0) | 4986 | 10967 |
| I ₄ (Etc 1.2) | 5136 | 11250 |
| SEm± | 49 | 102 |
| CD(P= 0.05) | 151 | 315 |
| Cultivars | | |
| V ₁ (Raj 4120) | 3788 | 7957 |
| V ₂ (Raj 4079) | 4974 | 11051 |
| V ₃ (Raj 4238) | 4861 | 10835 |
| SEm± | 39 | 80 |
| CD(P= 0.05) | 113 | 229 |
| Date of sowing | | |
| D ₁ (15 th NOV.) | 5201 | 11411 |
| D ₂ (30 th NOV.) | 4780 | 10520 |
| D ₃ (15 th DEC.) | 3641 | 7912 |
| SEm± | 31 | 58 |
| CD(P= 0.05) | 86 | 164 |

timely sown crop than late sown crop due to longer period for all the phenological stages in the timely sown crop. The late sowing decreased the duration of phenophases which caused a decrease in agrometeorological indices during various phenophases forced the crop to attain early maturity. This decline in agrometeorological indices accumulation was due to prevailing low temperature during vegetative phases and high temperature during reproductive phases of development in late-sown crop. Our results were similar to those of [11]. Paul and Sarker [12] also reported that the requirement of heat units decreased for different phenological stages with delay in sowing.

3.3 Wheat Yield

Data on grain yield (Table 4) indicated that it was markedly influenced due to different irrigation schedules. The treatment I₄ (Irrigations at 1.2 ETc) recorded significantly higher grain and biological yield of wheat (5136 and 11250 kg ha⁻¹) in pooled analysis, respectively over I₁ and I₂ and at par with I₃ (Irrigations at 1.0 ETc). While significantly lowest grain and biological yield (3442 and 7445 kg ha⁻¹) in pooled analysis was recorded under the treatment I₁ (Irrigations at 0.6 ETc). Higher grain and biological yield under the treatment I₄ and I₃ might be the result of cumulative effect of improvement in growth and yield attributes such as effective tillers, number of grains spike⁻¹ test weight, spike length, number

of spikes per unit area, number of spikelets spike⁻¹. Plant nutrients, particularly nitrogen, phosphorus, and potassium, were found to be more available with sufficient moisture in the soil profile under the I₄ irrigation schedule, and may have translocated to produce more dry matter. Second, higher yields with higher irrigation levels may be owing to irrigation's vital function in root development by reducing soil mechanical resistance, increasing transpiration, increasing nutrient uptake, and increasing photosynthesis due to plant metabolic activities [13]. The other reason of yield increase might be that irrigation scheduling at 1.2 ETc and 1.0 ETc throughout growth and reproductive phase created longer reproductive period with larger photosynthetic surface and reproductive storage capacity to attain higher allocation of net photosynthates to yield. The results obtained by [14-18] also confirm the findings of present investigation. Wheat yield is a complicated process that is influenced by interactions between the source (photosynthesis and assimilate availability) and sink components (storage organs). As a result of the significant improvement in both of these regulatory processes as shown by higher accumulation of biomass and nutrients as well as yield components under cultivar Raj 4079, grain and biological yield increased significantly. Further, the yield of wheat is dependent on two most important components namely spikes per unit area and weight of grains (test weight). Thus, due to a greater number of grains by virtue

of increased number of spikes and more test weight under Raj 4079, increased the grain yield over Raj 4238 and Raj 4120, and remained at par with cultivar Raj 4238. Since, biological yield is a sum of grain and straw yield produced by the crop, the increased grain yield under Raj 4079 might have resulted in higher biological yield in this cultivar. The marked variation in various yield components and yield between cultivars was observed by [19-23]. Grain, biological yield of wheat increased significantly when sowing of wheat on 15th November. Grain and biological yield decreased significantly as sowing was delayed from 15th November. This might be due to cumulative effect of poor expression of vegetative growth and yield contributing characters *i.e.*, number of spikes, ear length, grains spike⁻¹ and test weight under late sown conditions accompanied with high temperature and hot winds which leads toward forced maturity of the crop and ultimately resulted in lower grain and biological yield. The early sown crop, on the other hand, having favorable cool weather conditions for longer duration recorded better growth and yield attributes resulted in greater productivity [24].

4. CONCLUSION

Based on the results of two year experimentation, it may be inferred that irrigation scheduling treatment I₄ (1.2 ETC) recorded significantly higher value of yield and Growing degree days (GDD), By cultivar Raj 4079 observed the higher value of yield and GDD. Further, results showed that the wheat crop sown on 15th November obtained the maximum yield and growing degree days value with phenological stages. The number of days required to attain different phenological stages decreased with late sowing condition.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Nagarajan S, Anand A, Chaudhary HB, Response of spring wheat (*Triticum aestivum* L.) genotypes under changing environment during grain filling period. Indian Journal of Agricultural Sciences. 2008;78: 177-179.

2. Sikder S. Accumulated heat unit and phenology of wheat cultivars as influenced by late sowing heat stress condition. Agricultural Rural Development. 2008;7: 57-64.
3. Hakim MA, Hossain A, Teixeira da Silva JA, Zvolinsky VP, Khan MM. Yield, protein and starch content of 20 wheat (*Triticum aestivum* L.) genotypes exposed to high temperature under late sowing conditions. Journal of Scientific Research. 2012;4:477-489.
4. Bishnoi OP, Singh S, Niwas R. Effect of temperature on phenological development of wheat (*Triticum aestivum* L.) crop in different row orientations. Indian Journal of Agricultural Sciences. 1995;65:211-214.
5. Kumar R, Ramesh K, Singh RD, Prasad R. Modulation of wild marigold (*Tagetes minuta* L.) Phenophases towards the varying temperature regimes - A field study. Journal of Agrometeorology. 2010;12:234-240.
6. Phadnawis BN, Saini AD. Photothermal effects on tiller survival of wheat. Ann. Plant Physiol. 1992;6: 262-267.
7. Jat BT, Dhakar LL, Poonia TC. Phenological and heatunit accumulation of wheat (*Triticum aestivum* L. emend Fiori and Paol) varieties under varying sowing dates and seed rates. Agriculture Science Digest, 2003;23:84-87.
8. Nuttonson MY. Wheat climatic relationship and use of phenology in ascertaining the thermal and photothermal requirements of wheat. Am. Inst. Crop Ecol., Washington D.C;1955.
9. Pal SK, Kaur S, Thakur R, Verma VN, Singh MK. Effect of irrigation seeding dates and fertilizer on growth and yield of wheat (*Triticum aestivum* L.). Indian Journal of Agronomy. 1996;41:366-369.
10. Praveen KV, Patel SR, Choudhary JL, Bhelawe S. Heat unit requirement of different rice varieties under Chhattisgarh plain zones of India. Earth Sciences Climatic Change. 2013;5(1):1-4.
11. Khichar ML, Niwas R. Thermal effect on growth and yield of wheat under different sowing environments and planting systems. Indian Journal of Agriculture Research. 2007;41:92-96.
12. Paul NK, Sarker DK. Accumulated heat units and phenology relationships in wheat as influenced by sowing dates. Bangladesh Journal of Botany. 2000; 29:49-54.

13. Bhunia SR, Chauhan RPS, Yadav BS, Bhati AS. Effect of phosphorus, irrigation and rhizobium on productivity, water use and nutrient uptake in fenugreek (*Trigonella foenum graecum* L). Indian Journal of Agronomy. 2006;51:239-241.
14. Sharma KD, Pannu RK. Physiological response of wheat (*Triticum aestivum* L.) to limited irrigation. Journal of Agrometeorology. 2008;10:113-117.
15. Sarwar N, Mawqsood M, Mubeen K, Shehzad M, Bhullar MS, Qamar R, Akbar N. Effect of different levels of irrigation on yield and yield components of wheat cultivars. Pakistan Journal of Agricultural Sciences 2010;47:371-374.
16. Kumar B, Dhar S, Vyas AK, Paramesh V. Impact of irrigation schedules and nutrient management on growth, yield and root traits of wheat (*Triticum aestivum* L.) varieties. Indian Journal of Agronomy. 2015; 60:87-91.
17. Mishra G, Kushwaha HS. Winter wheat yield and soil physical properties responses to different tillage and irrigation. European Journal of Biological Research. 2016;56:530-537.
18. Rohj MK, Bhunia SR, Shivran H, Bhawariya H, Bawaliya SC, Ramniwas, Mandeewal RL. Effect of irrigation levels and intervals on groundnut (*Arachis hypogaea* L.) cultivars under drip system. International Journal of Plant & Soil Science. 2021;33(6):41-45.
19. Amrawat T, Solanki NS, Sharma SK, Jajoria DK, Dotaniya ML. Phenology growth and yield of wheat in relation to agrometeorological indices under different sowing dates. African Journal of Agricultural Research. 2013;49:6366-6374.
20. Pandey, I.B, Thakur, S.S and Singh, S.K. Response of timely sown wheat (*Triticum aestivum* L.) varieties to seed rate and fertility level. Indian Journal of Agronomy, 1999;44:745-749.
21. Nainwal, K. and Singh, M. Varietal behavior of wheat (*Triticum aestivum* L.) to dates of sowing under Tarai region of Uttar Pradesh. Indian Journal of Agronomy, 2000;45:107-113.
22. Sardana V, Sharma SK and Randhawa AK. Performance of wheat cultivars under different sowing dates level of nitrogen under rainfed conditions. Annals of Agricultural Research, 1999;20: 60-63.
23. Singh, R.K., Singh, S.K. and Singh, L.B. Integrated nitrogen management in wheat (*Triticum aestivum* L.). Indian Journal of Agronomy. 2007; 52:124-126.
24. Kulhari, S.C.; Sharma, S.L.; Kantwa, S.R. Effect of varieties, sowing dates and nitrogen levels on yield, nutrient uptake and quality of durum wheat (*Triticum durum* Desf.). Annals of Agricultural Research. 2003;24:332-336.

© 2022 Prajapat et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/83009>